


Nuclear modification factor of inclusive charged particles in Au+Au collisions at $\sqrt{s_{NN}} = 27$ GeV with the STAR experiment.

Alisher Aitbayev^{1,2,3} (for the STAR collaboration)

¹Joint Institute for Nuclear Research, Dubna - Russia,

²The Institute of Nuclear Physics, Almaty - Kazakhstan

³Al-Farabi Kazakh National University, Almaty - Kazakhstan



Supported in part by




U.S. DEPARTMENT OF
ENERGY

Office of
Science

Abstract

The Quantum Chromodynamics (QCD) phase diagram, often represented using coordinates of temperature (T) and baryonic chemical potential (μ_B), includes a transition from a hadronic gas phase to a quark-gluon plasma (QGP) phase. The Beam Energy Scan (BES) program at Relativistic Heavy Ion Collider (RHIC) varies the gold-gold collision energy aiming to explore the phase diagram and pinpoint the critical point. BES's initial phase (2010-2014) revealed intriguing results, including the suppression of high transverse momentum particle production ($p_T > 2$ GeV/c) at collision energies from $\sqrt{s_{NN}} = 62.4$ to 200 GeV that is quantified by the nuclear modification factor (R_{CP}). In 2018, STAR at RHIC collected a large-statistics dataset at $\sqrt{s_{NN}} = 27$ GeV, ten times larger than BES-I. This poster introduces new BES-II measurements of inclusive charged particles at 27 GeV, extending BES-I findings across a wider transverse momentum range with better precision. The relevant physics implications including the potential jet quenching effects at low energy collisions will also be discussed.

Introduction



QCD Phase Diagram

- ✦ Cross-over transition expected at low baryon chemical potential (μ_B)
- ✦ First-order transition expected at high μ_B
- ✦ Critical point is the end point of the first order phase transition

Beam Energy Scan (BES)

- ✦ Explore the QCD matter by colliding gold ions at different energies - and search for the potential QCD critical point
- ✦ Seeking to map onset of deconfinement, and the predicted QCD critical point

Motivation




Figure 1: Dependence of the nuclear modification factor on the transverse momentum of produced particles (BES-I)

The suppression effect of charged particle production with high transverse momenta ($p_T > 2$ GeV/c) is one of the most interesting results observed at the Solenoidal Tracker At RHIC (STAR) experiment during the BES-I program. This effect has been interpreted as the increase in energy loss of partons in the quark-gluon plasma produced at high energy heavy ion-collisions. It is commonly referred to as jet quenching in dense

partonic matter and was predicted as a sign of the formation of the QGP phase, where simple model of hadron scattering cannot describe the observations. This effect can be quantified using the nuclear modification factor R_{CP} :

- Nuclear modification factor:

$$R_{CP} = \frac{\langle N_{coll} \rangle_{Peripheral}}{\langle N_{coll} \rangle_{Central}} \frac{\left(\frac{d^2 N}{dp_T d\eta} \right)_{Central}}{\left(\frac{d^2 N}{dp_T d\eta} \right)_{Peripheral}} \quad (1)$$




Figure 2: Comparison of statistics between *BES-I* (2010-2017) and *BES-II* (2019-2021)

Results 1

The transverse momentum particle spectra for Au+Au collisions at energy of $\sqrt{s_{NN}} = 27$ GeV for inclusive charged particles in different centrality classes are shown in figure 3.




Figure 3: Transverse momentum distribution of inclusive charged particles for collision energy of 27 GeV. Each spectrum corresponds to a certain centrality class and is multiplied by coefficient from $1 - 10^5$ for visibility. The vertical error bars correspond to statistical uncertainties and the colored boxes to the systematic uncertainties.

From figure 3, it can be noticed that in the BES-II program, the spectra have a greater coverage in terms of transverse momentum p_T for all centrality classes, which enables a more comprehensive investigation of the nuclear modification factor.

Results 2




Figure 4: R_{CP} for inclusive charged particles at $\sqrt{s_{NN}} = 27$ GeV collision energy. The error band at unity on the right side of the plot corresponds to the p_T independent uncertainty on N_{bin} scaling. The vertical error bars correspond to statistical uncertainties and the colored boxes to the point-to-point systematic uncertainties.

$$R_{CP} = \frac{\langle N_{coll} \rangle_{Peripheral}}{\langle N_{coll} \rangle_{Central}} \frac{d^2 N / dp_T d\eta_{0-5\%}}{d^2 N / dp_T d\eta_{60-80\%}} \quad (2)$$

The growth of R_{CP} is seen at low values of p_T (up to $p_T \approx 2$ GeV/c), which is affected by effects such as Cronin enhancement, radial flow, and the relative dominance of coalescence over fragmentation during hadronization. However, as p_T increases, R_{CP} reaches a plateau and then demonstrates suppression of hadrons produced in central collisions with respect to peripheral collisions.

Conclusion

New data from the BES-II allow to extend investigation of the particle production modification in medium to the region of high transverse momenta p_T . First measurement of the nuclear modification factor R_{CP} at the collision energy of 27 GeV has shown a behavior similar to what was previously obtained at higher energies with a plateau and a decays at transverse momenta $p_T > 2$ GeV/c. An energy dependent study of the R_{CP} on data from BES-II should allow to better map the position of the phase transition from hadronic to partonic degrees of freedom in nuclear matter.

Acknowledgments: Supported by Russian Science Foundation under grant N 22-72-10028.