Local parity violation in the dense hadronic medium

O. O. Novikov in collaboration with A. Zakharova

Saint Petersburg State University

VII International Conference Models in Quantum Field Theory (MQFT-2022)

《曰》 《聞》 《臣》 《臣》 三臣

Local \mathcal{CP} violation in QCD

When **finite volume** is considered a metastable axial topological charge may arise in hot QCD

$$T_5 = \frac{1}{8\pi^2} \int d^3 x \epsilon_{jkl} \operatorname{Tr} \left[G^j \partial^k G^l - i \frac{2}{3} G^i G^k G^l \right]$$

Such fluctuations of the topological charge may appear in a fireball produced in the heavy-ion collisions. The lifetime for its fluctuation is comparable to the lifetime of the fireball and estimated to be $\Delta t \sim \tau_f \sim 5 - 10 fm/c.$

In the infinite volume limit this fluctuation is associated with a sphaleron (localized lump connecting vacua with different topological number) transitions



Chiral imbalance

The topological charge change is,

$$\Delta T_{5} = \frac{1}{8\pi^{2}} \int_{0}^{\tau_{f}} dt \int_{Vol} d^{3}x \operatorname{Tr}[G_{\mu\nu}\tilde{G}^{\mu\nu}] = \frac{1}{2\pi^{2}} \int_{0}^{\tau_{f}} \int_{Vol} d^{3}x \partial_{\mu}K^{\mu}$$

Where $\tilde{G}^{\mu\nu} = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} G_{\alpha\beta}$. The effects of this local topological charge may be mimicked by introduction of the θ -term to the Lagrangian with $\theta(t, x)$, $\partial_t \theta = \mu_5$

For the massless quarks $(m_{u,d} \ll \tau_f^{-1})$ the quark axial current $J_5^{\mu} = i\bar{q}\gamma_5\gamma^{\mu}q$ satisfies the conservation law broken by the chiral anomaly

$$\partial_{\mu}J_{5}^{\mu} = \frac{N_{f}}{2\pi^{2}}\partial_{\mu}K^{\mu} \Rightarrow \frac{d}{dt}\left(Q_{5} - 2N_{f}T_{5}\right) = 0$$

Thus, the local fluctuations of the topological charge should also produce the local fluctuations of $Q_5 = N_R - N_L$ - chiral imbalance

Chiral magnetic effect

In the peripheral heavy ion collisions these fluctuations result in a **chiral magnetic effect** [Kharzeev, McLerran, Warringa, 2008]

In the peripheral collision a large magnetic field is produced along the angular momentum.

Positively charged right-handed and negatively charged left-handed quarks tend to orient their spins and momenta along the magnetic field. Negatively charged left-handed and positively charged right-handed orient in the opposite directions.

As result, in the presence of the chiral imbalance the electric current is produced along the direction of the magnetic field

What are effects in the central collisions?

< ロ > < 同 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

Effective meson description on a chiral background

In the hadronic phase we may use the effective field theory description with the appropriate modifications

• Scalar and pseudoscalar mesons

Chiral perturbation theory with the modified covariant derivative,

$$D_{lpha} \mapsto D_{lpha} - i\{\mu_5 \delta_{lpha,0}, \cdot\}$$

Vector mesons

Vector dominance model with the extra term,

$$\epsilon^{\mu\nu\alpha\beta}\operatorname{Tr}\left[\zeta_{\mu}V_{\nu}V_{\mu\nu}\right]$$

where $\zeta^{\mu} = \mu_5 \delta^{\mu 0}$

• Bottom-up holographic models - large charge difference for A_L^a and A_R^b 5d fields dual to the J_L^{μ} and J_R^{μ} currents, Chern-Simons interaction

Manifestations of the chiral imbalance in the hadronic medium

Using these effective field theories a number of phenomena inside the fireball was predicted by A.A.Andrianov and his coauthors

- \bullet For scalar mesons: parity violating decays $\eta,\eta'\to\pi\pi$
- Exotic parity violating decays of \tilde{a}_0 , $\tilde{\pi}$ meson states inside the fireball
- Asymmetry of the photon polarizations in $\pi^\pm\gamma\to\pi^\pm\gamma$ scattering
- For vector mesons: the splitting of the longitudinal and transverse polarization masses
- An observable: anomalous yield of dileptons from the vector meson decays

イロト 不得下 イヨト イヨト 二日

Lowest radiatively induced lagrangian for QED and ρ and ω mesons

Vector meson dominance model [V.Kovalenko,A.Andrianov,V.Andrianov]

$$\begin{split} \mathcal{L}_{\text{eff}} &= -\frac{F_{\mu\nu}F^{\mu\nu}}{4} - \frac{\omega_{\mu\nu}\omega^{\mu\nu}}{4} - \frac{\rho_{\mu\nu}\rho^{\mu\nu}}{4} \\ &+ \frac{5m_V^2e^2A_\mu A^\mu}{9g^2} - \frac{m_V^2eA^\mu\omega_\mu}{3g} - \frac{m_V^2eA^\mu\rho_\mu}{g} + \frac{b^2N_cV^\nu V_\nu}{24\pi^2m^2} \\ &- \frac{N_c\epsilon_{\delta\gamma\mu\nu}b^\mu V^\nu V^{\delta\gamma}}{8\pi^2} + \frac{V_{\gamma\lambda}N_cV_\mu^\lambda b^\gamma b^\mu}{12\pi^2m^2} + \frac{1}{2}m_V^2\omega_\mu\omega^\mu + \frac{1}{2}m_V^2\rho_\mu\rho^\mu \\ &V_\mu \equiv -eA_\mu Q + \frac{1}{2}g_\omega\omega_\mu \mathsf{I}_q + \frac{1}{2}g_\rho\rho_\mu\lambda_3 \end{split}$$

O. O. Novikov (SPbU)

< ロ > < 同 > < 回 > .

Image: Image:

Simplified model

To get qualitative understanding of the behavior of the vector mesons in the hadronic medium, let us consider a simplified model.

A massive electrodynamics with θ -term

$$S = \int d^{4}x \Big(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{4} \theta(x) F_{\mu\nu} \tilde{F}^{\mu\nu} + \frac{1}{2} m^{2} A^{\mu} A_{\mu} + A^{\mu} \partial_{\mu} B + \frac{1}{2} B^{2} \Big)$$

The auxiliary field B is introduced to make a massive model self-consistent by the Stückelberg mechanism. It gives,

$$(\Box + m^2)(\partial_\mu A^\mu) = 0$$

Such model was used in [A.A.Andrianov, V.A.Andrianov, D. Espriu, S.S.Kolevatov] and [A.A.Andrianov, S.S. Kolevatov, R.Soldati] with,

$$\partial_{\mu}\theta \sim \zeta_{\mu}x^{\mu}\theta\Big(-\zeta_{\alpha}x^{\alpha}\Big)$$

O. O. Novikov (SPbU)

MQFT-2022

Bubble boundary

Such model was used to study two processes:

- The transition of the effective meson states to the ordinary mesons after the fireball decay
- The propagation of the meson states through the spatial boundary of the fireball

However the ansatz considered may be associated only with the very edge of the fireball boundary. Also μ_5 corresponds to $\partial_t \theta$.



Near the boundary

Assume that the boundary thickness is much smaller than the fireball

$$egin{aligned} &A^{\mu}(t,x,y,z)=\int d^2kA^{\mu}(t,z)\exp(ik_xx+ik_yy),\ &A_{\pm}=A^x\pm iA^y,\quad k_{\pm}=k_x\pm ik_y \end{aligned}$$

Then the equations take the form,

$$(\Box + m^2)A^t = -\frac{1}{2}(\partial_z \theta) \Big[k_- A_+ - k_+ A_- \Big],$$
$$(\Box + m^2)A^z = \frac{1}{2}(\partial_t \theta) \Big[k_- A_+ - k_+ A_- \Big],$$
$$(\Box + m^2)A_{\pm} = \pm k_{\pm} \Big[(\partial_t \theta)A^z + (\partial_z \theta)A^t \Big] \pm i \Big[(\partial_t \theta)\partial_z A_{\pm} - (\partial_z \theta)\partial_t A_{\pm} \Big]$$

For simplicity: momentum directed perpendicularly to the boundary i.e. $k_{\pm} = 0$. Then A_{\pm} decouple both from the equations on A^t and A^z .

O. O. Novikov (SPbU)

< ロ > < 同 > < 回 > < 回 > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ > < □ >

10/17

3

Circular polarizations

$$(\partial_t^2 - \partial_z^2 + m^2)A_{\pm} = \pm i \Big[(\partial_t \theta) \partial_z A_{\pm} - (\partial_z \theta) \partial_t A_{\pm} \Big]$$

 A_{-} satisfies the same equation but in the reverse time.

Assume that $\partial_t \theta$ is small compared to $\partial_t A_+$ (adiabatic regime)

This would correspond to the chemical potential being small compared to the spatial change near the boundary

$$A_+ = a(t,z) \exp\left(-i\int dt\,\omega(t)
ight)$$

a and $\omega(t)$ are slowly varying at the same rate as θ .

Adiabatic regime

In the leading approximation $i(\partial_t \theta)$ neglected,

$$a \simeq a_0, \quad \omega \simeq \omega_0$$

 $-\partial_z^2 a_0 + \left[m^2 - \omega_0^2 + \omega_0(\partial_z \theta)\right] a_0 = 0$

Schrödinger equation though with the potential proportional to the spectral parameter $\omega_{\rm 0}$

Sample shape for $\boldsymbol{\theta}$ - the transition between two homogeneous regions

$$heta(t,z) = ar{ heta}(t) + \Delta heta(t) \cdot anh\left[\mu(t)\Big(z - z_0(t)\Big)
ight],$$

 $heta \xrightarrow[z o -\infty]{d} - \Delta heta, \quad heta \xrightarrow[z o +\infty]{d} + \Delta heta,$

O. O. Novikov (SPbU)

MQFT-2022

イロト 不得 トイヨト イヨト 二日

Sample potential

For this shape of θ the potential becomes the Pöschl-Teller one,

$$\left[-\partial_z^2 + m^2 - \omega_0^2 + \frac{\omega_0 \mu \cdot \Delta \theta}{\cosh^2 \mu (z - z_0)}\right] a_0 = 0, \qquad (1)$$

The potential has the continuous spectrum of the wavelike solutions for $\varepsilon > 0$ i.e. $|\omega_0| > m$.

In the band $|\omega_0| < m$ the bound states may exist. This is true for $\omega\mu\cdot\Delta\theta<$ 0,

$$\varepsilon_n = -\mu^2 (\lambda - n - 1)^2, \quad \lambda(\lambda - 1) = \left| \frac{\omega_0 \Delta \theta}{\mu} \right|$$

where *n* is integer from 0 to the largest integer $n_{max} \leq \lambda - 1$,

$$\lambda^4 - 2\lambda^3 + \left(1 + \Delta\theta^2\right)\lambda^2 - 2(n+1)\Delta\theta^2\lambda + \Delta\theta^2\left((n+1)^2 - \frac{m^2}{\mu^2}\right) = 0$$

Bound states spectrum



(a) $m/\mu = 0.5$ (b) $m/\mu = 1.0$

Disappearance and crossing of the levels - nonadiabaticity!

O. O. Novikov (SPbU)

Time dependence influence

At small times we may approximate the influence of the $\partial_t \theta(t, z) \equiv v(t, z)$ term with the adiabatically changing perturbation that changes the frequencies and profile functions,

$$\omega = \omega_0 + \omega_1, \quad a = a_0 + a_1, \quad \omega_1, a_1 \sim |v|$$

The perturbation is non-Hermitian and results in,

$$\omega_1 = -\frac{i}{2} \frac{\int_{-\infty}^{+\infty} dz \, (\partial_z v) a_0^2}{\int_{-\infty}^{+\infty} dz \, (2\omega_0 + \partial_z \theta) a_0^2}$$

For state n = 0 in our sample potential,

$$\omega_{1,n=0} = -\frac{i}{2} \frac{\frac{\lambda-1}{2\lambda-1}\mu(\partial_t \Delta \theta)}{\omega_0 + \frac{\lambda-1}{2\lambda-1}\mu \Delta \theta}$$

This means that the θ time-dependence results in the damping or amplification of the bound states

O. O. Novikov (SPbU)

Local parity violation

MQFT-2022

- We argue that the bound currents near the transition region of the fireball exist
- The present analysis was done in the adiabatic approximation which is likely to be violated in the actual fireball
- Some non-adiabatic processes are evident already in our approximation from the disappearance and crossing of the levels
- Impact on the signatures for the chiral imbalance? Resonances in the transition probabilities? Explosion of boundary currents? Work in progress.

Thank you for your attention

O. O. Novikov (SPbU)

< ∃ →

э

< □ > < /□ >