

VII International Conference "Models in Quantum Field Theory"
(MQFT-2022)



Contribution ID: 137

Type: **not specified**

Arising of mass in scalar quantum field theories.

Monday, 10 October 2022 18:00 (25 minutes)

We investigate massive models of quantum field theory of scalar field in logarithmic dimensions in Euclidean space. The Schwinger-Dyson equation and non-trivial solution for mass are considered in the paper.

The Schwinger-Dyson equation has the form:

$$D^{-1} = \Delta^{-1} - \Sigma$$

where D is a full propagator, Δ is a bare propagator, Σ is a self-energy operator. In the minimal subtraction (MS) scheme it holds:

$$\Delta(p) = 1/p^2$$

where p is a momentum. The inverse full propagator has the following characteristic:

$$r \int D^{-1}(p) p^2 = -m^2 = 0 \quad (\partial \partial (p^2) D^{-1}(p)) p^2 = -m^2 = 1/A.$$

In the main approximation of perturbation theory it holds:

$$D(p) = A/p^2 + m^2$$

where A is an amplitude, m is a mass. We investigate the scalar models ϕ_3 , ϕ_4 and ϕ_6 . For the theories ϕ_3 and ϕ_4 mass appears in the first order of perturbation theory whereas for the ϕ_6 -theory the mass does not appear in the first order.

Presenter: PISMENSKY, Artem (Saint Petersburg Electrotechnical University "LETI")

Session Classification: Section C

Track Classification: Section C: Field theoretical methods in statistical physics