

Anomaly sum rules for the case of a singlet current and their features

Tuesday, 10 November 2020 16:30 (15 minutes)

We generalize the dispersive approach to axial anomaly by A.D. Dolgov and V.I. Zakharov to a non-Abelian case with arbitrary photon virtualities. We derive the anomaly sum rule for the singlet current and obtain the $\pi^0, \eta, \eta' \rightarrow \gamma\gamma^{(*)}$ transition form factors. Using them, we established the behavior of a non-perturbative gluon matrix element $\langle 0 | G\tilde{G} | \gamma\gamma^{(*)} \rangle$ for the case of a real photons and for the case of a one virtual photon in both space-like and time-like regions. We found a significant contribution of the non-Abelian axial anomaly to the processes with one virtual photon, comparable to that of the electromagnetic anomaly. The duality between the axial and the vector channels was observed: the values of duality intervals and mixing parameters in the axial channel were related to vector resonances' masses and residues. The possibility of a light pseudoscalar glueball-like state is conjectured.

Primary authors: Mr OGANESIAN, Armen; TERYAEV, Oleg; Mr KHLEBTSOV, Sergei; KLOPOT, Yaroslav

Presenter: Mr KHLEBTSOV, Sergei

Session Classification: Theoretical Physics

Track Classification: Theoretical Physics