




Фазовая диаграмма КХД от LHC к NICA (эксперимент)

*Рогачевский Олег
for MPD collaboration*


Физика тяжелых ионов:
от LHC к NICA
3 февраля 2017
Дубна

QCD Phase diagram


N. Cabibbo and G. Parisi,
Phys. Lett. B59 (1975) 6769




STAR BES program




FAIR - CBM




arXiv:1502.0273/nuc-ex



NICA - MPD - BM@N



Fukushima, Hatsuda
Rep. Prog. Phys. 74 (2011) 014001




Hagedorn temperature

Statistical Thermodynamics
of Strong Interactions at High Energies.

R. HAGEDORN
CERN - Geneva

(ricevuto il 12 Marzo 1965)




Experimental data:

1411 states known in 1967;
4627 states of mid '90s.

$$\bar{\rho}_{\text{exp}}(m) = \frac{1}{\sqrt{2\pi\tau^2}} \sum_{i=1}^N \nu_i \exp\left[-\frac{(m - m_i)^2}{2\tau^2}\right], \quad \tau = 200 \text{ MeV},$$

J. CLEYMANS and D. WORKU
Modern Physics Letters A
Vol. 26, No. 16 (2011) 1197–1209



$$\rho_h(m) = \frac{c}{(m^2 + m_0^2)^{5/4}} \exp\left(\frac{m}{T_H}\right)$$




Parameter	
$c \text{ (GeV)}^{3/2}$	0.16 ± 0.02
$m_0 \text{ (GeV)}$	0.5
$T_H \text{ (GeV)}$	0.174 ± 0.011

where the sum goes from the pion mass to the highest known resonances.

Accelerators for Relativistic Nuclear Physics

Accelerator	Place	Ion periods	Energy	Projectiles
Synchro-Phasatron	JINR Dubna	1971 - 1985	3.6 AGeV	d, He, C
Bevalac	LBNL Berkeley	1972 - 1984	< 2AGeV	C,Ca,Nb, Ni,Au,...
AGS	BNL, Brookhaven	1986 - 1994	14,5/11,5 AGeV	Si, Au
SPS	CERN, Geneva	1986 - 2002	200/158 AGeV	O,S,In,Pb
SIS 18	GSI,Darmstadt	1992 - today	2 AGeV	Kr,Au
Nuclotron	JINR Dubna	1993 - today	< 4.5 AGeV	p, d, He,C,Li, Mg, Kr
RHIC	BNL, Brookhaven	2000 - today	$\sqrt{s_{NN}} = 200 \text{ GeV}$	Cu, Au
LHC	CERN, Geneva	2010	$\sqrt{s_{NN}} = 5.5 \text{ TeV}$	Pb
NICA	JINR Dubna	2019	$\sqrt{s_{NN}} = 4 - 11 \text{ GeV}$	p - Au
SIS 100	GSI,Darmstadt	2025	2 – 11 AGeV	Au

Relativistic nuclear physics in JINR LHE



$$\alpha(x_d) = \frac{d^2\sigma(d + \text{Cu} \rightarrow \pi^- + \dots)}{d^2\sigma(p + \text{Cu} \rightarrow \pi^- + \dots)}$$

Fixed Target Experiments at Relativistic Energies

Beam energies: 100A MeV \equiv 2A GeV

Pioneering experiments

Synchrophasotron – Dubna (1971 – 1985) DISK, 2-m B.C.

BEVALAC: Plastic Ball and Streamer Chamber (1972 - 1986)

2-nd generation experiments

SIS-100 GSI: FOPI, KAOS (finished),
HADES (1990 – today)

BEVALAC: EOS-TPC, DLS (1990 – 1992)

Physics:

Collective effects => Discovery and investigation of flow effects


Equation of state (EOS) => Study of compressibility of dense nuclear matter

In-medium modifications => Kaons, low mass di-leptons

Basic result:

Nuclear matter can be compressed and high energy densities can be achieved

Alternating Gradient Synchrotron



BNL-AGS (1986 – 2002)


(1986 – 1991):

^{16}O & ^{28}Si , $E_{\text{lab}}^{\text{max}} = 14.5 \text{ A GeV}$

1991: AGS Booster, to have more intense proton beams and heavy ions at the AGS

(1992 – 1994): "heavy" Au ions

^{197}Au , $E_{\text{lab}}^{\text{max}} = 11.5 \text{ A GeV}$




Heavy Ion Experiments at the AGS

5 large experiments: E802/866/917, E810, E814/877, E864,E895.

Experiment	Beam	Technology	Observables
E802	Si	Single arm magnetic spectrometer	Spectra (π , p , K^\pm), HBT
E810		TPCs in magnetic field	Strangeness (K_s^0 , Λ)
E814		Magnetic spectrometer + calorimeters	Spectra (p) + E_t
E859		E802 + 2 nd level PID trigger	Strangeness (Λ)
E866	Au	2 magnetic spectrometers (TPC, TOF)	Strangeness (Kaons)
E877		Upgrade of E814	
E891		Upgrade of E810	
E895		EOS TPC	Spectra (π , p , K^\pm), HBT
E896		Drift chamber + neutron detector	H^0 Di-baryon, Λ
E910		EOS TPC + TOF	p+A Collisions
E917		Upgrade of E866	


E895/910 experiment

- EOS TPC; developed for Bevalac experiment
- Spectra (π^\pm , p , K^\pm), particle correlation, HBT



CERN accelerator complex

CERN-SPS (1986 – 2004): $\sqrt{s} = 17$ GeV, Pb + Pb collisions
7 large experiments: WA80/98, NA35/49/61, NA38/50/60,
NA44, NA45/CERES, WA97/NA57, NA52.





Heavy Ion Experiments at the SPS

Experiment	Beam	Technology	Observables
NA34	$^{16}\text{O}, ^{32}\text{S}$	Muon spectrometer + calorimeter	Di-leptons, ρ , π , K , γ
NA35		Streamer chamber	π , K_s^0 , Λ , HBT
NA36		TPC	K_s^0 , Λ
NA38		Di-muon spectrometer (NA10)	Di-leptons, J/ψ
WA80/WA93		Calorimeter + Plastic Ball	γ , π^0 , η
WA85		Mag. spectrometer with MWPCs	K_s^0 , Λ , Ξ
WA94		WA85 + Si strip detectors	K_s^0 , Λ , Ξ
NA44	$^{16}\text{O}, ^{32}\text{S}, ^{208}\text{Pb}$	Single arm magnetic spectrometer	π , K^\pm , ρ
NA45		Cherenkov + TPC	Di-leptons (low mass)
NA49	^{208}Pb	Large volume TPCs	π , K^\pm , ρ , K_s^0 , Λ , Ξ , Ω , ...
NA50		NA38 upgrade	Di-leptons, J/ψ
NA52		Beamline spectrometer	Strangelets
WA97		Mag. spectrometer with Si tracker	h^- , K_s^0 , Λ , Ξ , Ω
WA98		Pb-glass calorimeter + mag. spectrom.	γ , π^0 , η
NA57	WA97 upgrade	h^- , K_s^0 , Λ , Ξ , Ω	
NA60	^{114}In	NA50 + Si vertex tracker	Di-leptons, J/ψ

Onset of deconfinement (NA49/61)


Statistical Model of the Early Stage

Gazdzicki M. Gorenstein M.
Acta. Phys. Pol., B30: 2705 1999




NA49 scan

arXiv:nucl-ex/0612007



Horn
vanishing




The scaled variance of the multiplicity distribution of negatively charged hadrons in the projectile hemi-sphere

RHIC

BNL-RHIC (from 2000):

$\sqrt{s} = 200 \text{ GeV}$, Au + Au collisions

4 large experiments: BRAHMS, PHENIX, PHOBOS, STAR.



Heavy Ion Experiments at RHIC

Experiment	Technology	Observables
STAR	TPC and Si vertex tracker (+ EMCAL, TOF)	π , K^\pm , p , K_s^0 , Λ , Ξ , Ω , ...
PHENIX	Drift chambers, calorimeter, RICH, TOF, muon spectrometer	γ , π^0 , η , J/ψ , K^\pm , p , ...
BRAHMS	2 arm magnetic spectrometer	π , K^\pm , p (large acceptance)
PHOBOS	Magnetic spectrometer with Si tracker	charged particles (large acceptance)

The Quark-Gluon-Plasma is Found at RHIC

BNL -73847-2005
Formal Report

CONTENTS

Hunting the Quark Gluon Plasma

RESULTS FROM THE FIRST 3 YEARS AT RHIC

ASSESSMENTS BY THE EXPERIMENTAL COLLABORATIONS

April 18, 2005



Forward.....	1
Quark Gluon Plasma and Color Glass Condensate at RHIC? The Perspective from the BRAHMS Experiment.....	1
Formation of Dense Partonic Matter in Relativistic Nucleus-Nucleus Collisions at RHIC: Experimental Evaluation by the PHENIX Collaboration	33
The PHOBOS Perspective on Discoveries at RHIC	159
Experimental and Theoretical Challenges in the Search for the Quark Gluon Plasma: The STAR Collaboration's Critical Assessment of the Evidence from RHIC Collisions	253

The early measurements have revealed compelling evidence for the existence of a new form of nuclear matter at extremely high density and temperature – a medium in which the predictions of QCD can be tested, and new phenomena explored, under conditions where the relevant degrees of freedom, over nuclear volumes, are expected to be those of quarks and gluons, rather than of hadrons. This is the realm of the quark gluon plasma, the predicted state of matter whose existence and properties are now being explored by the RHIC experiments.

Relativistic Heavy Ion Collider (RHIC) • Brookhaven National Laboratory, Upton, NY 11974-5000

STAR BES program (2005)

BES-Short-v8.3_0

Experimental Study of the QCD Phase Diagram and
Search for the Critical Point:
Selected Arguments for the Run-10 Beam Energy Scan
at RHIC

The STAR Collaboration (B. I. Abelev et al.)

Introduction & Summary

We present an overview of the main ideas that have emerged from discussions within STAR for the Beam Energy Scan (BES). The formulation of this concise and abridged document is facilitated by the existence of a much longer and more comprehensive companion document entitled Experimental Exploration of the QCD Phase Diagram: Search for the Critical Point [1].:

A. A search for turn-off of new phenomena already established at higher RHIC energies; QGP signatures are the most obvious example, but we define this category more broadly. If our current understanding of RHIC physics and these signatures is correct, **a turn-off must be observed in several signatures, and such corroboration is an essential part of the “unfinished business” of QGP discovery [2].**

STAR BES QGP signatures

The particular observables that STAR has identified as the essential drivers of our run plan are:

- (A-1) Constituent-quark-number scaling of v_2 , indicating partonic degrees of freedom;
- (A-2) Hadron suppression in central collisions as characterized by the ratio R_{cp} ;
- (A-3) Untriggered pair correlations in the space of pair separation in azimuth and pseudorapidity, which elucidate the ridge phenomenon;
- (A-4) Local parity violation in strong interactions, an emerging and important RHIC discovery in its own right, is generally believed to require deconfinement, and thus also is expected to turn-off at lower energies.


A search for signatures of a phase transition and a critical point. The particular observables that we have identified as the essential drivers of our run plan are:

- (B-1) Elliptic & directed flow for charged particles and for identified protons and pions, which have been identified by many theorists as highly promising indicators of a “softest point” in the nuclear equation of state;
- (B-2) Azimuthally-sensitive femtoscopy, which adds to the standard HBT observables by allowing the tilt angle of the ellipsoid-like particle source in coordinate space to be measured; these measurements hold promise for identifying a softest point, and complements the momentum-space information revealed by flow measurements, and
- (B-3) Fluctuation measures, indicated by large jumps in the baryon, charge and strangeness susceptibilities, as a function of system temperature – the most obvious expected manifestation of critical phenomena.

STAR BES I results


Number of constituent quarks scaling

Phys. Rev. C88, (2013), 014902




Ridge effect

B. Abelev et al., Phys. Rev. C80, 064912 (2009).




High P_T suppression

Stephen Horvat Quark Matter 2015




Chiral Magnetic Effect




STAR BES I results


PRL 112 (2014) 162301




R. Lacey, PRL 114, 142301 (2015)



STAR, PRL 112, 032302 (2014)





Selected observables that show interesting non-monotonic behavior as functions of collision energy around $\sqrt{s_{NN}} = 15-40$ GeV.

Top panel: The difference $R_{out}^2 - R_{side}^2$ – between the squared radii in the outward and sideways directions measured via two-pioninterferometry vs. $\sqrt{s_{NN}}$ using STAR, PHENIX, and ALICE data.



$R_{out}^2 - R_{side}^2$ – reflects the lifetime of the collision fireball and was predicted to reach a maximum for collisions in which a hydrodynamic fluid forms at temperatures where the equation of state is softest.

Middle panel: The rapidity-slope of the net proton directed flow v_1 , dv_1/dy . This quantity is sensitive to early pressure gradients in the medium.

Bottom panel: The kurtosis of the event-by-event distribution of the net proton (i.e. proton minus antiproton) number per unit of rapidity, normalized such that Poisson fluctuations give a value of 1. In central collisions, published results in a limited kinematic range [347] show a drop below the Poisson baseline around $\sqrt{s_{NN}} = 27$ and 19.6 GeV. New preliminary data over a larger pT range [348], although at present still with substantial error bars, hint that the normalized kurtosis may, in fact, rise above 1 at lower $\sqrt{s_{NN}}$, as expected from critical fluctuations [349]. The grey band shows the much reduced uncertainties anticipated from BES-II in 2018-2019, for the 0-5% most central collisions.


QCD Phase diagram (experiment)

Grazyna
Odyniec
Journal of Physics: Conference Series
455 (2013) 012037



The dependence of T_{ch} on μ_B , fitted with the Grand Canonical approach in THERMUS Model

SN0598 : Studying the Phase Diagram of QCD Matter at RHIC




Contents

1 Introduction	3
2 Review of BES-I Results and Theory Status	5
2.1 Region of the Phase Diagram Accessed in BES-I	5
2.2 Search for the Critical Point	8
2.3 Search for the First-order Phase Transition	10
2.3.1 Directed Flow (v_1)	10
2.3.2 Average Transverse Mass	12
2.4 Search for the Threshold of QGP Formation	13
2.4.1 Elliptic Flow	13
2.4.2 Nuclear Modification Factor	18
2.4.3 Dynamical Charge Correlations	21
2.4.4 Chiral Transition and Dileptons	24
2.5 Summary of BES-I	27
3 Proposal for BES Phase-II	30
3.1 Physics Objectives and Specific Observables	30
3.1.1 R_{CP} of identified hadrons up to $p_T = 5 \text{ GeV}/c$	32
3.1.2 The v_2 of ϕ mesons and NCQ scaling for identified particles	33
3.1.3 Three-particle correlators related to CME/LPV	34
3.1.4 The centrality dependence of the slope of $v_1(y)$ around midrapidity	36
3.1.5 Proton-pair correlations	38
3.1.6 Improved $k\sigma^2$ for net-protons	40
3.1.7 Dilepton production	41
3.2 Beam request	43
3.3 The Fixed-Target Program	45
3.4 The Importance of $p+p$ and $p+A$ Systems	45
3.5 Collider Performance	47
3.6 Detector Upgrades	47
3.6.1 <i>iTPC</i>	48
3.6.2 <i>EPD</i>	49
4 Summary	50


STAR BES II program

$\sqrt{s_{NN}}$ (GeV)	μ_B (MeV)	MinBias Events (10^6)	Time (weeks)	Year
7.7	420	4.3	4	2010
11.5	315	11.7	2	2010
14.5	260	24.0	3	2014
19.6	205	35.8	1.5	2011
27.0	155	70.4	1	2011
39.0	115	130.4	2	2010
62.4	70	67.3	1.5	2010

$\sqrt{s_{NN}}$ (GeV)	μ_B (MeV)	Needed Events (10^6)
7.7	420	100
9.1	370	160
11.5	315	230
14.5	260	300
19.6	205	400





Year	System and Energy	Physics/Observables	Upgrade
2017	<ul style="list-style-type: none"> p+p @ 500 GeV Au+Au @ 62.4 GeV 	<ul style="list-style-type: none"> Spin sign change diffractive Jets 	FMS post-shower, EPD (1/8 th), eTOF prototype
2018	<ul style="list-style-type: none"> Zr+Zr, Ru+Ru @ 200 GeV Au+Au @ 27 GeV 	<ul style="list-style-type: none"> CME, di-leptons CVE 	Full EPD? eTOF prototype
2019	Au+Au @ 14.5-20 GeV + fixed target	<ul style="list-style-type: none"> QCD critical point Phase transition CME, CVE,... 	Full iTPC, eTOF, and EPD
2020	Au+Au @ 7-11 GeV + fixed target	<ul style="list-style-type: none"> QCD critical point Phase transition CME, CVE,... 	
2020+	<ul style="list-style-type: none"> Au+Au @ 200 GeV p+A/p+p @ 200 GeV 	<ul style="list-style-type: none"> Unbiased jets, open beauty PID FF, Drell-Yan, longitudinal correlations 	<ul style="list-style-type: none"> HFT+ FCS, FTS




The timeline for future RHIC and LHC heavy ion running

Fixed Target Program with STAR




Au+Au FXT at 3.9 GeV

- Extend energy reach to overlap/complementary AGS/FAIR/JPARC
- Real collisions taken in run 14 and results (K. Meehan @ QM15 & WWND16)
- Upgrades (iTTPC+eTOF+EPD) crucial
- Unprecedented coverage and PID for Critical Point search in BES-II
- Spectra, flow, fluctuations and correlations



CBM Phase-0 Exp: eTOF at STAR




Install, commission and use 10% of the CBM TOF modules, including the read-out chains at STAR, starting in 2019


CBM participating in RHIC Beam Energy BES-II in 2019-2020:

- Complementary to part of CBM's physics program: $\sqrt{s_{NN}} = 3, 3.6, 3.9, 4.5, 7.7$ GeV especially for the physics of **B**- & **s**-production and fluctuations
- Operating of ~30 CBM TOF modules and electronics (~10 m², 10k channels)
- Experiencing with detector system, online calibration and monitoring tools
- Developing analysis strategies for particle identification and efficiency extraction


BES-III: CBM@BNL




- 1) Study QCD phase structure
- 2) Maintain heavy ion community
- 3) CBM@eRHIC is an add on cost



2022 - 2025
 CBM@AGS
 $\sqrt{s_{NN}} \leq 5.4 \text{ GeV}$




2025 - ...
 CBM@eRHIC
 $\sqrt{s_{NN}} \leq 14 \text{ GeV}$



Nu Xu


Resent & future experiments for HIC

Facility	SPS	RHIC BES II	Nuclotron M	NICA	SIS/100 (300)	J-PARK HI
Laboratory	CERN Geneva	BNL Brookhaven	JINR Dubna	JINR Dubna	FAIR GSI Darmstadt	J-PARK
Experiment	NA61 SHINE	STAR PHENIX	BM@N	MPD	HADES CBM	JHITS
Start of data taking	2011	2017	2015	2019	2020/25	2025
$\sqrt{s_{NN}}$ (GeV)	4.9 – 17.3	7.7 – 200	< 3.5	4 - 11	2.7 – 8.2	2.0 – 6.2
Physics	CP & OD	CP & OD	HDM	OD & HDM	OD & CP	OD & HDM





CP — critical point
 OD — onset of deconfinement, mixed phase, 1st order phase transition
 HDM — hadrons in dense matter
 PDM — properties of deconfined matter

Nuclotron based Ion Collider facility





BM@N experiment at NICA

AuAu $E_{\text{beam}} = 4 \text{ GeV}$




year	2016	2017 spring	2017 autumn	2019	2020 and later
beam	d(↑)	C, Ar	Kr	Au	Au, p
max.inten- sity, Hz	1M	1M	1M	1M	10M
trigger rate, Hz	10k	10k	20k	20k	50k
central tracker status	6 GEM half pl.	8 GEM half pl.	10 GEM half pl.	8 GEM full pl.	12 GEMs or 8 GEMs + Si planes
experim. status	techn. run	techn. run	physics run	stage 1 physics	stage 2 physics



MPD experiment at NICA



AuAu $\sqrt{s} = 11$ GeV





CBM experiment at FAIR



- $10^5 - 10^7$ Au+Au reactions/sec
- Hit densities $10/\text{cm}^2/\text{event}$ (at STS-1)
- High hit rates $1-100 \text{ MHz}/\text{cm}^2$
- Fast and radiation hard detectors
- Free-streaming readout electronics
- Online event selection
- High precision vertex reconstruction
- Identification of leptons and hadrons

$700 \pi^+$, $160 \pi^-$, $53 K^+$, $32 K^0$, $27 K_s^0$, $\sim 1 \Xi^-$, $0.022 \Omega^-$

CBM & NICA Physics



**Thank you for
attention**

