


Study of the resonances in heavy-ion collisions at NICA energies using the MPD detector

V. Riabov* for the MPD Collaboration



* This work was supported by RFBR according to the research project № 18-02-40038

Outline

- ❖ Motivation for resonance studies in heavy-ion collisions
- ❖ Expectations for resonance properties in heavy-ion collisions at NICA energies
- ❖ Feasibility studies for particle reconstruction at NICA-MPD
- ❖ Summary

Resonances in heavy-ion collisions

- ❖ Wide variety of resonances in the PDG, most often/easily measured are:



Particle	Mass (MeV/c ²)	Width (MeV/c ²)	Decay	BR (%)
ρ^0	770	150	$\pi^+\pi^-$	100
K^{*+}	892	50.3	$\pi^+K_s^0$	33.3
K^{*0}	896	47.3	π^0K^+	66.7
ϕ	1019	4.27	K^+K^-	48.9
Σ^{*+}	1383	36	$\pi^+\Lambda$	87
Σ^{*-}	1387	39.4	$\pi^-\Lambda$	87
$\Lambda(1520)$	1520	15.7	K^-p	22.5
Ξ^{*0}	1532	9.1	$\pi^+\Xi^-$	66.7


- ❖ Vacuum properties of the resonances are well defined (m, τ , BR etc.)
- ❖ Copiously produced in heavy-ion collisions at \sim GeV energies, large branching ratios in hadronic decay channels \rightarrow possible to measure
- ❖ Probe reaction dynamics and particle production mechanisms vs. system size and $\sqrt{s_{NN}}$:
 - ✓ hadron chemistry and strangeness production, ϕ with hidden strangeness is one of the key probes
 - ✓ reaction dynamics and shape of particle p_T spectra, p/K^* , p/ϕ vs. p_T
 - ✓ lifetime and properties of the hadronic phase
 - ✓ flow, comparison with e^+e^- measurements, jet quenching, background for other probes etc.

Hadronic phase and medium modifications

increasing lifetime \longrightarrow

	$\rho(770)$	$K^*(892)$	$\Sigma(1385)$	$\Lambda(1520)$	$\Xi(1530)$	$\phi(1020)$
$c\tau$ (fm/c)	1.3	4.2	5.5	12.7	21.7	46.2
σ_{rescatt}	$\sigma_{\pi}\sigma_{\pi}$	$\sigma_{\pi}\sigma_K$	$\sigma_{\pi}\sigma_{\Lambda}$	$\sigma_K\sigma_p$	$\sigma_{\pi}\sigma_{\Xi}$	$\sigma_K\sigma_K$

- ❖ Resonances have small lifetimes of $c\tau \sim 1 - 45$ fm, part of them decays in the fireball
- ❖ Reconstructed resonance yields in heavy ion collisions are defined by:
 - ✓ resonance yields at chemical freeze-out
 - ✓ hadronic processes between chemical and kinetic freeze-outs:
 - rescattering:** daughter particles undergo elastic scattering or pseudo-elastic scattering through a different resonance \rightarrow parent particle is not reconstructed \rightarrow loss of signal
 - regeneration:** pseudo-elastic scattering of decay products ($\pi K \rightarrow K^0$, $KK \rightarrow \phi$ etc.) \rightarrow increased yields




- SPS/RHIC/LHC observed multiplicity dependent suppression of ρ/π , K^*/K , Λ^*/Λ ratios, resonances with $c\tau \leq 20$ fm/c. Ratios of longer lived resonances are not affected
- Results support the existence of a hadronic phase that lives long enough to cause a significant reduction of the reconstructed yields of short lived resonances
- Hadronic phase lifetime, $\tau \sim 10$ fm/c*
- NICA: $\langle dN_{ch}/d\eta \rangle^{1/3} \sim 6^{**} \rightarrow$ RHIC/LHC report modifications at such multiplicities

* ALICE, Phys.Lett.B 802 (2020) 135225, Phys.Rev.C 99 (2019) 024905

** PHENIX, Phys.Rev.C 93 (2016) 2, 024901


Hadronic phase and particle ratios

- ❖ Models with hadronic cascades (UrQMD, PHSD, AMPT) → properties of hadronic phase
- ❖ Models predict centrality dependent ρ/π , K^*/K , ϕ/K and Λ^*/Λ , Σ^*/Λ , Ξ^*/Ξ ratios in AuAu@11
- ❖ Ratios are suppressed going from peripheral to central collisions for resonances with small τ
- ❖ Modifications occur at low momentum as expected for the hadronic phase effects



- ❖ Models predict yield modifications qualitatively similar to those obtained at SPS/RHIC/LHC:
 - ✓ lifetime and density of the hadronic phase are high enough
 - ✓ modification of particle properties in the hadronic phase should be taken into account when model predictions for different observables are compared to data
 - ✓ study of short-lived resonances is a unique tool to tune the hadronic phase simulations

- ❖ Phase 1: **TPC, TOF, FFD, FHCAL** и **ECAL**
- ❖ Startup in 2022
- ❖ Simulate AuAu@4-11 collisions using different event generators
- ❖ Propagate particles through the MPD, ‘mpdroot’:
 - ✓ Geant (v.3 or v.4) for particle transport
 - ✓ realistic simulation of subsystem response (raw signals)
 - ✓ track/signal reconstruction and pattern recognition
- ❖ Basic event and track selections:
 - ✓ event selection: $|Z_{\text{vtx}}| < 50$ cm
 - ✓ track selection:
 - number of TPC hits > 24
 - $|\eta| < 1.0$
 - $|DCA \text{ to PV}| < 3\sigma$ for primary tracks
 - V0 topology cuts for weakly decaying secondaries
 - $p_T > 50$ MeV/c
 - TPC-TOF combined $\pi/K/p$ PID
 - ✓ combinatorial background:
 - event mixing ($|\Delta Z_{\text{vtx}}| < 2$ cm, $|\Delta_{\text{Mult}}| < 20$, $N_{\text{ev}} = 10$)



TPC: $|\Delta\phi| < 2\pi$, $|\eta| \leq 1.6$






TOF, EMC: $|\Delta\phi| < 2\pi$, $|\eta| \leq 1.4$

FFD: $|\Delta\phi| < 2\pi$, $2.9 < |\eta| < 3.3$

FHCAL: $|\Delta\phi| < 2\pi$, $2 < |\eta| < 5$

Reconstruction efficiency: $\rho(770)$, $K^*(892)$, $\phi(1020)$, $\Lambda(1520)$

❖ Typical reconstruction efficiencies ($A \times \epsilon$) in AuAu @ **4**, **7.7** and **11** GeV, $|y| < 1$








❖ Reasonable efficiencies in the wide p_T range, $|y| < 1$

❖ Modest multiplicity (and/or $\sqrt{s_{NN}}$) dependence

Mass resolution: $\rho(770)$, $K^*(892)$, $\phi(1020)$, $\Lambda(1520)$

❖ Detector mass resolution ($m_{\text{reconstructed}} - m_{\text{generated}}$) in AuAu @ **4**, **7.7** and **11** GeV, $|y| < 1$









❖ Acceptable mass resolution

❖ Modest multiplicity (and/or $\sqrt{s_{NN}}$) dependence

$\phi(1020)$, reconstructed peaks


- ❖ UrQMD v.3.4: AuAu@11 (10M events), AuAu@7.7 (5M events), AuAu@4 (5M events)
- ❖ Full chain simulation and reconstruction, $p_T = 0.2-0.4$ GeV/c, $|y| < 1$



- ❖ Mixed-event combinatorial background is scaled to foreground at high mass and subtracted
- ❖ Distributions are fit to Voigtian function + polynomial
- ❖ Signal can be reconstructed at $p_T > 0.2$ GeV/c, high- p_T reach is limited by available statistics
- ❖ S/B ratios deteriorates with increasing centrality and collision energy

$K^*(892)$ and $\Lambda(1520)$, reconstructed peaks


❖ UrQMD v.3.4: AuAu@11 (10M events), AuAu@7.7 (5M events), AuAu@4 (5M events), $|y| < 1$



- ❖ Signal can be reconstructed from zero momentum, high- p_T reach is limited by statistics
- ❖ S/B ratios deteriorates with increasing centrality and collision energy

MC closure tests: ρ , $K^{*0,\pm}$, ϕ , Λ^*


- ❖ UrQMD v.3.4: AuAu@11 (10M events), AuAu@7.7 (5M events), AuAu@4 (5M events)
- ❖ Full chain simulation and reconstruction, p_T ranges are limited by the possibility to extract signals, $|y| < 1$




- ❖ Reconstructed spectra match the generated ones within uncertainties
- ❖ Measurements are possible starting from \sim zero momentum, sample p_T spectra in a wide range
- ❖ Maximum raw yields (smallest stat. uncertainties) are extracted at \sim 300 MeV/c

More complex decays: $\Sigma(1385)^\pm, \Xi(1530)^0$



$$\Sigma(1385)^\pm \rightarrow \pi^\pm \Lambda \quad (\Lambda \rightarrow p \pi)$$



$$\Xi(1530)^0 \rightarrow \pi^+ \Xi^- \quad (\Xi^- \rightarrow \Lambda \pi^-, (\Lambda \rightarrow p \pi^-))$$




❖ Typical reconstruction efficiencies ($A \times \epsilon$) in AuAu @ 11 GeV, $|y| < 1$




Reconstruction: $\Sigma(1385)^\pm, \Xi(1530)^0$

- ❖ UrQMD v.3.4: AuAu@11 (10M events), full chain simulation and reconstruction, $|y| < 1$



- ❖ $\Sigma(1385)^\pm$ signal can be reconstructed starting from zero momentum, high- p_T reach is limited by statistics
- ❖ Monte Carlo closure test is passed



- ❖ For $\Xi(1530)^0$ observe a hint of a signal at $p_T > 0.4 \text{ GeV}/c$, statistics-hungry measurement
- ❖ Larger data sample and embedded simulations are required

Summary

- ✓ Measurement of resonances is important for the MPD physical program
- ✓ Models predict high sensitivity of resonances to the properties of the partonic/hadronic medium produced in heavy-ion collisions at NICA energies
- ✓ Resonances can be reconstructed/measured using the MPD detector from zero momentum to ~ 3 GeV/c with 10^7 minimum bias events sampled, x10 events is needed for multiplicity dependent studies
- ✓ Measurements of resonances is a plausible task for year-1 operation

BACKUP





- ❖ Modernization of the existing Nuclotron facility
- ❖ Fixed target experiment: BM@N
- ❖ Construction of collider complex to collide:
 - ✓ heavy ions up to Au, $\sqrt{s_{NN}} = 4-11 \text{ GeV}$, $\mathcal{L} \sim 10^{27} \text{ cm}^{-2}\text{s}^{-1}$
 - ✓ polarized p and d, $\sqrt{s_{NN}} = 27 \text{ GeV}$, $\mathcal{L} \sim 10^{32} \text{ cm}^{-2}\text{s}^{-1}$ (pp)
- ❖ Collider experiments: MPD, SPD
- ❖ NICA, MPD – start of operation in 2022



Strangeness enhancement in pp, p-A and A-A

Nature Phys. 13 (2017) 535



- ❖ Observed in heavy-ion collisions at AGS, SPS, RHIC and LHC;
- ❖ For the first time observed in pp and p-A collisions by ALICE at the LHC
- ❖ Observed as for ground-state hadrons as for resonances (ϕ/π , Σ^*/π , Ξ^*/π)
- ❖ Strangeness production in A-A collisions is reproduced by statistical hadronization models. Canonical suppression models reproduce results in pp and p-A except for ϕ
- ❖ ϕ with hidden strangeness is not subject to canonical suppression $\rightarrow \phi$ is a key observable !!!

Hadronization at intermediate momenta

❖ Baryon puzzle - increased baryon-to-meson (p/π , Λ/K_s^0 , Λ_c^+/D) ratios in heavy-ion collisions at RHIC and the LHC


❖ Driving force of enhancement is not yet fully understood:

- ✓ particle mass (hydrodynamic flow)?
- ✓ quark count (baryons vs. mesons)?


❖ ϕ and K^{*0} are well suited for tests as mesons with masses very close to that of a proton:

- ✓ $\Delta m_\phi \sim 80 \text{ MeV}/c^2$, $\Delta m_{K^{*0}} \sim -45 \text{ MeV}/c^2$

Phys.Lett. B736 (2014) 196-207





Phys.Rev. C88 (2013) 2, 024906



Model predictions for resonances at NICA




- ❖ UrQMD, PHSD, AMPT, EPOS ...
- ❖ General predictions:
 - ✓ resonances are still copiously produced and can be used to study physics of heavy-ion collisions
 - ✓ models predict enhanced production of particles with strangeness and different interplay of mechanisms responsible for shaping of the particle p_T spectra.



- ❖ Eventually, model predictions (integrated yields, $\langle p_T \rangle$, particle ratios etc.) should be compared to data to differentiate different model assumptions

Hadronic phase and particle ratios

❖ Modifications occur at low momentum as expected for hadronic phase effects









❖ Models predict yield modifications for resonances qualitatively similar to those observed at higher collision energies:

- ✓ lifetime and density of the hadronic phase are high enough
- ✓ modification of particle properties in the hadronic phase should be taken into account when model predictions for different observables are compared to data
- ✓ study of short-lived resonances is a unique tool to tune hadronic phase simulations

$\rho(770)$, reconstructed peaks

- ❖ UrQMD v.3.4: AuAu@11 (10M events), AuAu@7.7 (5M events), AuAu@4 (5M events)
- ❖ Full chain simulation and reconstruction, $p_T = 0.2-0.4$ GeV/c, $|y| < 1$



- ❖ Mixed-event background subtraction, fits to Voigtian function + polynomial
- ❖ Contributions from K_s , ω , K^{*0} , f_0 and f_2 are subtracted (need to be measured in advance)*
- ❖ Signal can be reconstructed at $p_T > 0$ GeV/c, high- p_T reach is limited by available statistics
- ❖ S/B ratios deteriorates with increasing centrality and collision energy

$\rho(770)$, signal extraction – practice tests

Phys.Rev. C99 (2019) no.6, 064901




Fig. 1: (Color online) Invariant mass distributions for $\pi^+\pi^-$ pairs after subtraction of the like-sign background. Plots on the left and right are for the low and high transverse momentum intervals, respectively. Examples are shown for minimum bias pp, 0–20% and 60–80% central Pb–Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. Solid red curves represent fits to the function described in the text. Colored dashed curves represent different components of the fit function, which includes a smooth remaining background as well as contributions from K_S^0 , ρ^0 , $\omega(782)$, $K^*(892)^0$, $f_0(980)$ and $f_2(1270)$. See text for details.