

Inha University, Republic of Korea

Nucleons in Nuclear Matter and Properties of Nuclei

Ulugbek Yakhshiev

Talk@APCTP-BLTP JINR-PNPI NRC KI-SPbU Joint Workshop "Modern Problems in Nuclear and Elementary Particle Physics" July 24-28, 2017, Petergof, St.Petersburg

Motivation

Q. <<"Hadronic vs Nuclear" models>> vs. <<"Hadronic+Nuclear" models>>? A. Depends on a choice which has an underlying purpose...

Our (choice) aim: To construct a model which describes at same footing

the single nucleon properties

- in free space considering it as a structure-full system
- in nuclear medium (possible structure changes)
- as well as the properties of the whole nucleonic systems
 - infinite nuclear matter properties (EOS, volume and symmetry energy properties)
 - matter under extreme conditions (e.g. neutron stars)
 - few/many (ordinary/exotic) nucleon systems (symmetric nuclei, mirror nuclei, rare isotopes, halo nuclei,...)
 - nucleon knock-out reactions (lepton-nucleus scattering experiments)
 - possible changes in in-medium NN interactions
 - etc

Strategy

- **Q.** How to construct a theoretical framework?
- A. It is well known...
 - the commonly accepted and the best way is to start from QCD and try to arrive some an effective framework (unfortunately it is not completely understood yet)
 - therefore, as much as possible main peculiarities of QCD must be taken into account in arriving an effective theory or in constructing a phenomenological approach which describe the hadrons and their interactions
 - at low energies main peculiarities (which obvious in a single hadron sector) are
 - chiral symmetry and its spontaneous breakdown
 - quark confinement (the mechanism is not understood yet)
 - in addition one should take into account the structure changes of in-medium nucleons in constructing the nuclear many body systems

Content

- Topological models
- Medium modifications
- In-medium nucleons
- Nuclear matter
- Nucleon in finite nuclei
- Properties of nuclei
- Summary and Outlook

Topological models

Q. What is a nucleon? A. It seems nontrivial to answer the question...

At the fundamental level we may have

- fermions -> then bosons are trivial fermion systems
- bosons -> then fermions are <u>nontrivial topological structures</u>

Structure

From what is made a nucleon and, in particular, its core in a starting boson picture approach?

- The structure treatment depends on an energy scale
- At the limit of large number colours Nc the core still has the mesonic content



Shell is made from the meson cloud

Topological models



The free space Lagrangian (which was widely in use)

[G.S.Adkins et al. Nucl.Phys. B228 (1983)]

$$\mathcal{L} = \frac{F_{\pi}^2}{16} \operatorname{Tr} \left(\partial_{\mu} U \partial^{\mu} U^{\dagger} \right) - \frac{1}{16e^2} \operatorname{Tr} \left[U^{\dagger} \partial_{\mu} U, U^{\dagger} \partial_{\nu} U \right]^2 + \frac{F_{\pi}^2 m_{\pi}^2}{16} \operatorname{Tr} \left(U + U^{\dagger} - 2 \right)$$

 Nontrivial structure: topologically stable solitons with the corresponding conserved topological number (baryon number) A

$$U = \exp\{i\overline{\tau} \,\overline{\pi} / 2F_{\pi}\} = \exp\{i\overline{\tau} \,\overline{n}F(r)\}$$

$$B^{\mu} = \frac{1}{24\pi^{2}} \varepsilon^{\mu\nu\alpha\beta} Tr(L_{\nu}L_{\alpha}L_{\beta}) \qquad L_{\alpha} = U^{+}\partial_{\alpha}U$$

$$A = \int d^{3}rB^{0}$$

$$H = M_{cl} + \frac{\overline{S}^{2}}{2I} = M_{cl} + \frac{\overline{T}^{2}}{2I},$$

$$|S = T, s, t \ge (-1)^{t+T} \sqrt{2T + 1}D_{-t,s}^{S=T}(A)$$

 Nucleon is quantized state of the classical soliton-skyrmion which rotates in the ordinary and an internal spaces

Q. What happens in the nuclear medium?

A. The possible medium effects

- Deformations (swelling or shrinking, multipole deformations) of nucleons \bigcirc
- Characteristic changes in: effective mass, charge distributions, all possible form factors
- NN interactions may change
- etc.

One should be able to describe all those phenomena

Soliton in the nuclear medium (phenomenological way)

- Outer shell modifications (informations from pionic atoms)
- Inner core modifications, in particular, at large densities (nuclear matter properties) 0

Inner core modifications in the nuclear medium may be related to:

- vector meson properties in the nuclear medium
- nuclear matter properties at saturation density

Meson cloud modifications in the nuclear medium: Pion physics in the nuclear medium

Medium modifications

"Outer shell" modifications

- In free space three types of pions can be treated separately: isospin breaking
- In nuclear matter: there are three types of polarization operators

$$(\partial^{\mu}\partial_{\mu} + m_{\pi}^2)\vec{\pi}^{(\pm,0)} = 0$$

$$(\partial^{\mu}\partial_{\mu} + m_{\pi}^2 + \hat{\Pi}^{(\pm,0)})\vec{\pi}^{(\pm,0)} = 0$$

$$\hat{\Pi}^0 = 2\omega U_{\text{opt}} = \chi_s(\rho, b_0, c_0) + \vec{\nabla} \cdot \chi_p(\rho, b_0, c_0) \vec{\nabla}$$

 Optic potential approach: parameters from the pion- 		π -atom	$T_{\pi}=50~{\rm MeV}$
nucleon scattering	$b_0 [m_{\pi}^{-1}]$	- 0.03	- 0.04
(including the isospin	$b_1 [m_{\pi}^{-1}]$	- 0.09	- 0.09
	$c_0 [m_{\pi}^{-3}]$	0.23	0.25
	$c_1 [m_{\pi}^{-3}]$	0.15	0.16
	g'	0.47	0.47

Medium modifications

"Outer shell" modifications in the Lagrangian

[U.Meissner et al., EPJ A36 (2008)]

$$\mathcal{L}_{2}^{*} = \frac{F_{\pi}^{2}}{16} \underbrace{\alpha_{\tau}}_{16} \operatorname{Tr} \left(\partial_{0}U\partial_{0}U^{\dagger}\right) - \frac{F_{\pi}^{2}}{16} \underbrace{\alpha_{s}}_{16} \operatorname{Tr} \left(\partial_{i}U\partial_{i}U^{\dagger}\right)$$
$$\mathcal{L}_{m}^{*} = -\frac{F_{\pi}^{2}m_{\pi}^{2}}{16} \underbrace{\alpha_{m}}_{16} \operatorname{Tr} \left(2 - U - U^{\dagger}\right)$$

- Due to the non-locality of optic potential the kinetic term is also modified
- Due to energy and momentum dependence of the optic potential parameters, the following parts of the kinetic term are modified in different forms:
 - Temporal part
 - Space part

	$\pi\text{-}\mathrm{atom}$	$T_{\pi}=50~{\rm MeV}$
$b_0 [m_{\pi}^{-1}]$	- 0.03	- 0.04
$b_1 [m_\pi^{-1}]$	- 0.09	- 0.09
$c_0 [m_{\pi}^{-3}]$	0.23	0.25
$c_1 [m_{\pi}^{-3}]$	0.15	0.16
g'	0.47	0.47

 $\hat{\Pi}^0 = 2\omega U_{\text{opt}} = \chi_s(\rho, b_0, c_0) + \vec{\nabla} \cdot \chi_p(\rho, b_0, c_0) \vec{\nabla}$

"Inner core" modifications [UY & H.Ch. Kim, PRC83 (2011); UY, JKPS62 (2013); UY, PRC88 (2013)]



Medium modifications



In-medium nucleons

Structure studies 1: Energy momentum tensor

- It allows to address the questions like:
 - How are the total angular momentum and angular momentum of the nucleon shared among its constituents?
 - How are the strong forces experienced by its constituents distributed inside the nucleon?
- EMT form factors in free space studied in lattice QCD, ChPT and in different models (chiral quark soliton model, Skyrme model, etc.)
- We made further step studying EMT form factors in nuclear matter

Structure studies 1: Energy momentum tensor

Definition

$$\langle p' | \hat{T}_{\mu\nu}(0) | p \rangle = \bar{u}(p', \, s') \left[M_2(t) \, \frac{P_\mu P_\nu}{M_N} + J(t) \, \frac{i(P_\mu \sigma_{\nu\rho} + P_\nu \sigma_{\mu\rho})\Delta^\rho}{2M_N} + d_1(t) \, \frac{\Delta_\mu \Delta_\nu - g_{\mu\nu}\Delta^2}{5M_N} \right] u(p, \, s) \, ,$$

 Three form factors give an information about energy distribution, angular momentum distribution and about the stabilization of strong forces inside the nucleon

$$\begin{split} T_{00}^{*}(r) &= \frac{F_{\pi,s}^{*2}}{8} \left(\frac{2\sin^{2}F}{r^{2}} + F'^{2} \right) + \frac{\sin^{2}F}{2\,e^{*2}\,r^{2}} \left(\frac{\sin^{2}F}{r^{2}} + 2F'^{2} \right) + \frac{m_{\pi}^{*2}F_{\pi,s}^{*2}}{4} \left(1 - \cos F \right), \\ T_{0k}^{*}(r,s) &= \frac{\epsilon^{klm}r^{l}s^{m}}{(s \times r)^{2}} \, \rho_{J}^{*}(r), \\ T_{ij}^{*}(r) &= s^{*}(r) \left(\frac{r_{i}r_{j}}{r^{2}} - \frac{1}{3}\,\delta_{ij} \right) + p^{*}(r)\,\delta_{ij} \end{split} \\ M_{2}^{*}(t) - \frac{t}{5M_{N}^{*2}} \, d_{1}^{*}(t) &= \frac{1}{M_{N}^{*}} \int d^{3}r \, T_{00}^{*}(r) \, j_{0}(r\sqrt{-t}), \\ d_{1}^{*}(t) &= \frac{15M_{N}^{*}}{2} \int d^{3}r \, p^{*}(r) \, \frac{j_{0}(r\sqrt{-t})}{t}, \\ M_{2}^{*}(0) &= \frac{1}{M_{N}^{*}} \int d^{3}r \, T_{00}^{*}(r) = 1, \quad J^{*}(0) = \int d^{3}r \, \rho_{J}^{*}(r) = \frac{1}{2} \, . \end{split}$$

Structure studies1: Energy momentum tensor related quantities [H.C.Kim, P. Schweitzer, UY, Phys.Lett. B718 (2012)]

Different quantities related to the nucleon EMT densities and their form factors: $T_{00}^*(0)$ denotes the energy in the center of the nucleon; $\langle r_{00}^2 \rangle^*$ and $\langle r_J^2 \rangle^*$ depict the mean square radii for the energy and angular momentum densities, respectively; $p^*(0)$ represents the pressure in the center of the nucleon, whereas r_0^* designates the position where the pressure changes its sign; d_1^* is the value of the $d_1^*(t)$ form factor at the zero momentum transfer.

$ ho/ ho_0$	$T_{00}^*(0)$ [GeV fm ⁻³]	$\langle r_{00}^2 angle^*$ [fm ²]	$\langle r_J^2 \rangle^*$ [fm ²]	p*(0) [GeV fm ⁻³]	r ₀ [fm]	d_1^*
0	1.45	0.68	1.09	0.26	0.71	-3.54
0.5	0.96	0.83	1.23	0.18	0.82	-4.30
1.0	0.71	0.95	1.35	0.13	0.90	-4.85

In-medium nucleons

Structure studies 1: Pressure distribution inside the nucleon in free space and in symmetric matter [H.C.Kim, P. Schweitzer, UY, Phys.Lett. B718 (2012)]



FIG. 3: (Color online) The decomposition of the pressure densities $4\pi r^2 p^*(r)$ as functions of r, in free space ($\rho = 0$) and at $\rho = \rho_0$, in the left and right panels, respectively. The solid curves denote the total pressure densities, the dashed ones represent the contributions of the 2-derivative (kinetic) term, the long-dashed ones are those of the 4-derivative (stabilizing) term, and the dotted ones stand for those of the pion mass term.

Structure studies 2: Transverse EM charge densities

- Definition of EM ff's $\langle N(p', S') | J_{\mu}^{EM}(0) | N(p, S) \rangle$ = $\overline{u}_N(p', S') \Big[\gamma_{\mu} F_1^*(q^2) + i \frac{\sigma_{\mu\nu} q^{\nu}}{2m_N} F_2^*(q^2) \Big] u_N(p, S)$.
- These Pauli and Dirac ff's can be expressed by Sachs ff's

$$G_E(Q^2) = F_1(Q^2) + \frac{Q^2}{4M_N^2}F_2(Q^2)$$

$$G_M(Q^2) = F_1(Q^2) + F_2(Q^2),$$

 They give an information about transverse charge distributions inside the nucleon

$$\rho_{0}^{*}(b) = \int_{0}^{\infty} \frac{Q \, dQ}{2\pi} J_{0}(bQ) \frac{G_{E}^{*}(Q^{2}) + \tau G_{M}^{*}(Q^{2})}{1 + \tau}$$

$$\rho_{T}^{*}(\mathbf{b}) = \rho_{0}^{*}(b) - \sin(\phi_{b} - \phi_{S})$$

$$\times \int_{0}^{\infty} \frac{Q^{2} \, dQ}{4\pi \, m_{N}} J_{1}(bQ) \frac{-G_{E}^{*}(Q^{2}) + G_{M}^{*}(Q^{2})}{1 + \tau}, \qquad \mathbf{b} = b(\cos\phi_{b}\hat{\mathbf{e}}_{x} + \sin\phi_{b}\hat{\mathbf{e}}_{y})$$

Structure studies 2: Transverse EM charge densities inside an unpolarized nucleon [UY, H.C.Kim, Phys.Lett. B726 (2013)]



Fig. 3. Quark transverse charge densities inside an unpolarized proton (upper panels) and a neutron (lower panels) in free space (left panels) and at nuclear matter density $\rho_0 = 0.5m_{\pi}^3$ (right panels).

In-medium nucleons

Structure studies 2: Transverse EM charge densities inside the polarized nucleon [UY, H.C.Kim, Phys.Lett. B726 (2013)]



Fig. 4. Transverse charge densities of the proton (upper panels) and neutron (lower panels) in free space (left panels) and in nuclear matter with the density $\rho_0 = 0.5m_{\pi}^3$ (right panels).

In-medium nucleons

Masses [UY, PRC88 (2013)]

- Isoscalar effective mass
- Isovector effective mass (relevant to: universe evolution in Early stage; stability of drip line nuclei; mirror nuclei; transport in heavy-ion collisions; asymmetric nuclear matter)
- Effective masses of the nucleons

$$m_{N,s}^{*} = M_{s}^{*} + \frac{3}{8\Lambda^{*}} + \frac{\Lambda^{*}}{2} \left(a^{*2} + \frac{\Lambda_{env}^{*2}}{\Lambda^{*2}} \right)$$

$$\Delta m_{np}^* = a^* + \frac{\Lambda_{env}^*}{\Lambda^*}$$

$$m_{n,p}^* = m_{N,s}^* - \Delta m_{np}^* T_3$$

Nuclear matter

From the Bethe-Weizsacker formula

$$\varepsilon(A,Z) = -a_V + a_S \frac{(N-Z)^2}{A^2} + \mathbb{M}$$

The binding-energy-formula terms in the framework of present model can be obtained considering

We are ready to reproduce
Volume term

Symmetric infinite nuclear matter
Asymmetry term
Isospin asymmetric environment
Surface and Coulomb terms
Nucleons in a finite volume
Finite nuclei properties

Local density approximation

Nuclear matter

The volume term and Symmetry energy

 At infinite nuclear matter approximation the binding energy per nucleon takes the form

 $\varepsilon(\lambda, \delta) = \varepsilon_V(\lambda) + \varepsilon_S \delta^2 + O(\delta^4) \equiv \varepsilon_V(\lambda) + \varepsilon_A(\lambda, \delta)$

- λ is normalised nuclear matter density
- δ is asymmetry parameter
- ϵ_s is symmetry energy
- In our model
 - Symmetric matter
 - Asymmetric matter

$$\varepsilon_{V}(\lambda) = m_{N,s}^{*}(\lambda,0) - m_{N}^{\text{free}}$$

$$\varepsilon_{A}(\lambda,\delta) = \varepsilon(\lambda,\delta) - \varepsilon_{V}(\lambda)$$

$$= m_{N,s}^{*}(\lambda,\delta) - m_{N,s}^{*}(\lambda,0) + m_{N,V}^{*}(\lambda,\delta)\delta$$

Nuclear matter

Nuclear matter properties

Symmetric matter properties (pressure, compressibility and third derivative)

$$p = \rho_0 \lambda^2 \frac{\partial \varepsilon_V(\lambda)}{\partial \lambda} \bigg|_{\lambda=1}, \quad K_0 = 9\rho^2 \frac{\partial^2 \varepsilon_V(\lambda)}{\partial \rho^2} \bigg|_{\rho=\rho_0} \qquad Q = 27\lambda^3 \frac{\partial^3 \varepsilon_V(\lambda)}{\partial \lambda^3} \bigg|_{\lambda=1}$$

Symmetry energy properties (coefficient, slop and curvature)

$$\varepsilon_{s}(\lambda) = \varepsilon_{s}(1) + \frac{L_{s}}{3}(\lambda - 1) + \frac{K_{s}}{18}(\lambda - 1)^{2} + \mathbb{X}$$

Symmetric nuclear matter

Volume energy [UY, PRC88 (2013)]

- Set I solid
- Set II dashed
- Set III dotted

For comparison: Akmal-Pandharipande-Ravenhall (APR) predictions [PRC 58, 1804 (1998)] are given by stars. (From Arigonna 2 body interactions + 3 body interactions)



Symmetric nuclear matter



For comparison: Right figure from Danielewicz- Lacey-Lynch, Science 298, 1592 (2002). (Deduced from experimental flow data and simulations studies)

Asymmetric nuclear matter

Symmetry energy

• Solid $L_s = 70 \text{ MeV}$

• Dashed
$$L_s = 40 \text{ MeV}$$

For comparison: Akmal-Pandharipande-Ravenhall (APR) predictions [PRC 58, 1804 (1998)] are given by stars.

(From arigonna 2 body interactions + 3 body interactions)





For comparison: Right figure from Danielewicz- Lacey-Lynch, Science 298, 1592 (2002). (Deduced from experimental flow data and simulations studies)

Low density behaviour of symmetry energy



Neutron star properties

• TOV equations

$$-\frac{dP(r)}{dr} = \frac{G\mathcal{E}(r)\mathcal{M}(r)}{r^2} \left(1 - \frac{2G\mathcal{M}(r)}{r}\right)^{-1} \left(1 + \frac{P(r)}{\mathcal{E}(r)}\right) \left(1 + \frac{4\pi r^3 P(r)}{\mathcal{M}(r)}\right)$$

Energy-pressure relation

$$P = P(\mathcal{E}) \qquad \qquad P(\lambda) = \rho_0 \lambda^2 \frac{\partial \varepsilon(\lambda, 1)}{\partial \lambda}, \\ \mathcal{E}(\lambda) = [\varepsilon(\lambda, 1) + m_N] \lambda \rho_0$$

Neutron star's mass

$$\mathcal{M}(r) = 4\pi \int_0^r \mathrm{d}r \, r^2 \mathcal{E}(r) \, .$$

Neutron star properties [UY, PLB749 (2015)]



From Ref. [J.M. Lattimer & M. Prakash, Astrophys. J. 550 (2001)].

Asymmetric nuclear matter

Neutron star properties

[UY, PLB749 (2015)]

TABLE III: Properties of the neutron stars from the different sets of parameters (see Tables I and II for the values of parameters): n_c is central number density, ρ_c is central energy-mass density, R is radius of the neutron star, M_{max} is possible maximal mass, A is number of baryons in the star, E_b is binding energy of the star. In the left panel we represent the neutron star properties corresponding to the maximal mass M_{max} and in right panel approximately 1.4 solar mass neutron star properties. The last two lines are results from the Ref. [21].

Set	n_c	$ ho_c$	R	$M_{\rm max}$	A	E_b	n_c	$ ho_c$	R	M	A	E_b
	$[\mathrm{fm}^{-3}]$	$[10^{15}\mathrm{gr/cm^3}]$	$[\mathrm{km}]$	$[M_{\odot}]$	$[10^{57}]$	$[10^{53} \mathrm{erg}]$	$[\mathrm{fm}^{-3}]$	$[10^{15} { m gr/cm}^3]$	$[\mathrm{km}]$	$[M_{\odot}]$	$[10^{57}]$	$[10^{53} \mathrm{erg}]$
III-a	1.046	2.445	10.498	2.226	3.227	8.721	0.479	0.861	11.587	1.402	1.898	3.503
III-b	1.045	2.444	10.547	2.223	3.216	8.557	0.471	0.861	11.772	1.402	1.895	3.453
III-c	1.037	2.424	10.616	2.221	3.200	8.397	0.460	0.832	11.953	1.402	1.887	3.339
III-d	1.047	2.452	10.494	2.221	3.213	8.598	0.481	0.867	11.619	1.402	1.893	3.422
III-e	1.044	2.440	10.554	2.218	3.203	8.495	0.473	0.858	11.809	1.403	1.890	3.384
III-f	1.040	2.433	10.609	2.216	3.189	8.311	0.464	0.842	11.992	1.403	1.887	3.334
SLy230a [21]	1.15	2.69	10.25	2.10	2.99	7.07	0.508	0.925	11.8	1.4	1.85	2.60
SLy230b [21]	1.21	2.85	9.99	2.05	2.91	6.79	0.538	0.985	11.7	1.4	1.85	2.61

Nucleon in finite nuclei

The nucleon in a nucleus will include

- Local density approach for environment
- R dependence of a results
- Deformations
 - In particular, axially symmetric case allows the deformations in polar direction
- Polar deformations can be represented
 - in the isotopic vector and
 - in the profile function

$$\boldsymbol{N}(\boldsymbol{r}-\boldsymbol{R}) = \begin{pmatrix} \sin \Theta(\boldsymbol{r}-\boldsymbol{R}) \cos \varphi \\ \sin \Theta(\boldsymbol{r}-\boldsymbol{R}) \sin \varphi \\ \cos \Theta(\boldsymbol{r}-\boldsymbol{R}) \end{pmatrix}$$

$$P = P(|\boldsymbol{r} - \boldsymbol{R}|, \theta), \qquad \Theta = \Theta(|\boldsymbol{r} - \boldsymbol{R}|, \theta)$$

$$U(\boldsymbol{r}-\boldsymbol{R}) = \exp\left[i\boldsymbol{\tau}\cdot\boldsymbol{N}(\boldsymbol{r}-\boldsymbol{R})P(\boldsymbol{r}-\boldsymbol{R})\right]$$



The Equations of Motion

• The coupled partial differential equations (not an easy problem)

$$f(F_{\tilde{r}\tilde{r}}, F_{\theta\theta}, F_{\tilde{r}}, F_{\theta}, \Theta_{\theta}, F, \Theta) = 0,$$

$$g(\Theta_{\theta\theta}, \Theta_{\theta}, F_{\tilde{r}}, F_{\theta}, \Theta, F) = 0,$$

A numerical variational method can be applied

$$P(r,\theta) = 2 \arctan\left\{\frac{r_0^2}{r^2}(1+m_\pi r)(1+u(\theta))\right\} e^{-f(r)r}$$

$$\Theta(r,\theta) = \theta + \zeta(r,\theta),$$

$$F(r) = 2 \arctan\left\{\frac{r_0^2}{r^2}(1+m_\pi r)\right\} e^{-f(r)r}, \qquad u(\theta) = \sum_{n=1}^{\infty} \gamma_n \cos^n \theta$$

$$f(r) = \beta_0 + \beta_1 e^{\beta_2 r^2}.$$

$$\zeta(r,\theta) = r e^{-\delta_0^2 r^2} \sum_{n=1}^{\infty} \delta_n \sin 2n\theta,$$

$$\lim_{r \to \infty} F(r) = D(1+m_\pi r) \frac{e^{-m_\pi r}}{r^2},$$

Accuracy of the variational method

- In spherically symmetric approximation (e.g. nucleon in the centre of the spherical nucleus) one can explicitly solve Equations of Motion and compare with results of variational method
- Skyrme term is not modified in nuclear matter (table below)

Element	t	r_0	$10eta_0$	eta_1	eta_2	$m_{ m p}^{*}$	$\varDelta m^*_{ m np}$	$\Delta m_{ m np}^{ m *(EM)}$	$\mu^*_{ m p}$	$\mu^*_{ m n}$	$\langle r^2 \rangle_{\mathrm{E,S}}^{*1/2}$	$\langle r^2 \rangle_{\rm E,V}^{*1/2}$
		[fm]	$[m_{\pi}]$	$[m_{\pi}]$	$[m_{\pi}^2]$	[MeV]	[MeV]	[MeV]	[n.m.]	[n.m.]	[fm]	[fm]
	i)	_	_	_	_	938.268	1.291	-0.686	1.963	-1.236	0.481	0.739
free space	ii)	0.954	0.075	1.311	-0.009	938.809	1.313	-0.687	1.966	-1.241	0.481	0.739
	i)	_	_	_	_	593.285	1.668	-0.526	2.355	-1.276	0.656	0.850
^{14}N	ii)	1.393	0.076	0.920	0.226	598.505	1.655	-0.536	2.230	-1.209	0.648	0.810
	i)	_	_	_	_	585.487	1.697	-0.517	2.393	-1.297	0.667	0.863
$^{16}\mathrm{O}$	ii)	1.426	0.076	0.907	0.219	590.175	1.685	-0.527	2.341	-1.232	0.660	0.825
	i)	_	_	_	_	558.088	1.804	-0.480	2.584	-1.422	0.722	0.942
³⁸ K	ii)	1.493	0.076	0.841	0.153	559.957	1.802	-0.485	2.550	-1.377	0.718	0.910
	i)	_	_	_	_	557.621	1.804	-0.478	2.569	-1.428	0.724	0.947
40 Ca	ii)	1.489	0.076	0.839	0.149	559.378	1.802	-0.483	2.557	-1.383	0.720	0.914

Nucleon in finite nuclei

The Hamiltonian of the model

• Has the form as in the case of symmetric top

$$\begin{split} \hat{H} &= M_{\rm NP}^* + \mathcal{M}_-^2 \Lambda_{\rm mes} + \frac{\Lambda_{\rm env}^{*2}}{2\Lambda_{\omega\Omega,33}^*} + \frac{(\hat{T}_1^2 + \hat{T}_2^2)\Lambda_{\Omega\Omega,12}^* + (\hat{J}_1^2 + \hat{J}_2^2)\Lambda_{\omega\omega,12}^*}{2(\Lambda_{\omega\omega,12}^*\Lambda_{\Omega\Omega,12}^* - \Lambda_{\omega\Omega,12}^{*2})} \\ &+ \frac{(\hat{T}_1\hat{J}_1 + \hat{T}_2\hat{J}_2)\Lambda_{\omega\Omega,12}^*}{\Lambda_{\omega\omega,12}^*\Lambda_{\Omega\Omega,12}^* - \Lambda_{\omega\Omega,12}^{*2}} + \frac{\hat{T}_3^2}{2\Lambda_{\omega\Omega,33}^*} - \boxed{\left(a^* + \frac{\Lambda_{\rm env}^*}{\Lambda_{\omega\Omega,33}^*}\right)} \hat{T}_3 \,. \end{split}$$
Neutron-proton mass difference in finite nuclei

The densities of nuclei (left) and the isoscalar mass in nuclei (right)



On the right panel R is a distance between the geometrical centres of nucleus and nucleon

Nucleon in finite nuclei



Baryon charge distribution inside the nucleon under the consideration

In free space (left)

and

in O16 (right), R = 3fm

The densities of nuclei (left) and the isoscalar mass in nuclei (right)



On the right panel R is a distance between the geometrical centres of nucleus and nucleon

Nucleon in finite nuclei





In free space (left)

and

in O16 (right), R = 1.5fm

Nucleon in finite nuclei

Baryon charge distribution inside the nucleon under the consideration



In O16 (left), R = 3fm

and

in Ca40 (right), R = 4.5fm

The neutron-proton mass difference in finite nuclei



R is a distance between the geometrical centres of nucleus and nucleon

The Nolen-Schiffer anomaly (NSA)

• The mass difference of mirror nuclei

$$\Delta M \equiv {}^{A}_{Z+1}\mathbf{M}_{N} - {}^{A}_{Z}\mathbf{M}_{N+1} = \Delta E_{\rm EM} - \Delta m_{\rm np}^{*}$$

- EM part was calculated with high accuracy (within 1% error) in very detailed form (e.g., the exchange term, the center-of-mass motion, finite-size effects of the proton and neutron charges, magnetic interactions, vacuum effects, the dynamical effect of the neutron-proton mass difference, and short-range two-body correlations, etc.)
- If neutron-proton mass difference does not change in nuclear matter then the above formula cannot be satisfied.

$$\overline{\Delta}_{\rm NSA} = \Delta m_{\rm np} - \left(\Delta \overline{m}_{\rm np}^{*(1)} + \Delta \overline{m}_{\rm np}^{*(2)} \right)$$

The Nolen-Schiffer anomaly (NSA)

Is defined as ("bar" means averaging over the R)

$$\overline{\Delta}_{\rm NSA} = \Delta m_{\rm np} - \left(\Delta \overline{m}_{\rm np}^{*(1)} + \Delta \overline{m}_{\rm np}^{*(2)} \right)$$

where

$$\begin{split} \Delta \overline{m}_{\rm np}^* &\approx \int \left(\Delta \psi_{\rm np}^{(2)} m_{\rm p}^* + \left(\psi^{(p)} \right)^2 \Delta m_{\rm np}^* \right) {\rm d}^3 R \\ &\equiv \Delta \overline{m}_{\rm np}^{*(1)} + \Delta \overline{m}_{\rm np}^{*(2)} \,, \end{split}$$

			Present approach							
Nuclei		$\overline{m}^*_{ m p}$		$\alpha_{\rm ren} = 0 \qquad \qquad \alpha_{\rm ren} = 0.95$				$\overline{\varDelta}_{\mathrm{NSA}}$	$\overline{\varDelta}_{\mathrm{NSA}}$	
	$\alpha_{\rm ren} = 0$	$\alpha_{\rm ren} = 0.95$	$\Delta \overline{m}_{\rm np}^{*(1)}$	$\Delta \overline{m}_{ m np}^{*(2)}$	$\overline{\varDelta}_{\rm NSA}$	$\Delta \overline{m}_{\mathrm{np}}^{*(1)}$	$\Delta \overline{m}_{ m np}^{*(2)}$	$\overline{\varDelta}_{\mathrm{NSA}}$	ref. [16]	ref. [17]
$^{15}\text{O-}^{15}\text{N}$	767.45	928.30	-4.27	1.56	4.02	-0.21	1.33	0.20	_	0.16 ± 0.04
${}^{17}\text{F-}{}^{17}\text{O}$	812.35	930.54	-5.53	1.52	5.33	-0.28	1.32	0.27	0.31	0.31 ± 0.04
39 Ca- 39 K	724.78	926.16	-8.11	1.67	7.75	-0.41	1.33	0.37	_	0.22 ± 0.08
41 Sc- 41 Ca	771.71	928.51	-9.74	1.62	9.44	-0.49	1.33	0.47	0.62	0.59 ± 0.08

U.Meissner, A.Wirzba, A.Rakhimov, UY [EPJ A36(2008)]

The nucleon mass in nuclei



Present work (left) and U.Meissner, A.Wirzba, A.Rakhimov, UY [EPJ A36(2008)] (right)

The neutron-proton mass difference in finite nuclei



Present work (left) and U.Meissner, A.Wirzba, A.Rakhimov, UY [EPJ A36(2008)] (right)

The present model describes at same footing

the single nucleon properties

- in free space considering it as a structure-full system
- in nuclear matter (EM and EMT form factors)
- as well as the properties of the whole nucleonic systems
 - infinite nuclear matter properties (volume and symmetry energy properties)
 - matter under extreme conditions (e.g. neutron stars)
 - few/many nucleon systems (symmetric nuclei, mirror nuclei, rare isotopes, halo nuclei,...)
 - nucleon knock-out reactions (lepton-nucleus scattering)
 - possible changes in in-medium NN interactions

• etc

Thank you very much for your attention!