



Contribution ID: 277

Type: Oral

Preparation the map of concentration alpha-radiation radon gas in building of Warsaw University of Technology.

Radon - as an alpha-radioactive gas naturally present in atmospheric air - is the main component of the exposure of Poles to ionizing radiation, accounting for almost 40% of the annual effective dose and over 70% of the dose received by a statistical Poles.

One of the commonly used passive methods for measuring radon concentrations in the air is the CR39 type Solid State Nuclear Tracker (SSNTD), a commercial name for allyl diglycol polycarbonate (PADC). These are small plates which, when measured, are placed in appropriately sized containers of known volume and then, after appropriate thermochemical treatment, are analyzed under a microscope. Particles of alpha, protons and heavy ions passing through the detector material damage the chemical bonds, which can be visualized by thermochemical treatment. As a result of etching on the plates are formed holes.

Preparing of map radon gas concentration in buildings of Warsaw University of Technology (WUT) contain:

1. Arrangement of CR39 detectors in buildings WUT
2. Preparation of measuring system
3. Research on optimization of thermochemical process
4. Measurement execution
5. Preparing an interactive concentration map

All research was done in cooperation with the science club CAMAC from the Faculty of Physics and the science club Chemical and Process Engineering (IChiP) from the Faculty of Chemical and Process Engineering WUT as part of the Rector's grant.

Primary author: Mr BOLEK, Karol (Warsaw University of Technology)

Co-authors: Mr MILER, Michał (Warsaw University of Technology); BORZEŃCKA, Nina (Warsaw University of Technology); PODGÓRSKA, Zuzanna (Central Laboratory for Radiological Protection, Warsaw University of Technology)

Presenter: Mr BOLEK, Karol (Warsaw University of Technology)

Track Classification: Applied Research