



Contribution ID: 245

Type: Oral

## Structure of B-10 and C-11,12 nuclei in relativistic dissociation

In the framework of the BECQUEREL Project in JINR the nuclear track emulsion (NTE) technique allowed one to investigate clustering of the nuclei Li, Be, B, C and N in their relativistic dissociation. With an unsurpassed spatial resolution (about  $0.5 \mu\text{m}$ ) NTE provides a complete observation of tracks starting from fission fragments and down to relativistic particles. Fragment tracks observed in NTE is a "building blocks" the light nuclei include the lightest clusters having no excited states, namely,  $\alpha$ -particles, tritons,  $^3\text{He}$  nuclei, and deuterons. A pair and triples of protons and  $\alpha$ -particles can constitute the unstable  $^8\text{Be}$  and  $^9\text{B}$ . Analysis of NTE exposed by  $^{11,12}\text{C}$  and  $^{10}\text{B}$  and investigation the role of unstable  $^8\text{Be}$  and  $^9\text{B}$  nuclei will be presented.

### Summary

Contribution of the unstable nuclei  $^8\text{Be}$  and  $^9\text{B}$  into dissociation of relativistic nuclei  $^{10}\text{B}$  and  $^{11,12}\text{C}$  is under study on the basis of the nuclear track emulsion exposed to secondary beams of the JINR Nuclotron. In a charge state distribution of fragments the share of the channel  $^{10}\text{B} \rightarrow 2\text{He} + \text{H}$  is 77%.

On the basis of measurements of fragment emission angles it is determined that unstable nucleus  $^8\text{Be}(\text{g.s.})$  manifests itself with a probability of  $(25 \pm 5)\%$  where  $(14 \pm 3)\%$  of them occur in decays of the unstable nucleus  $^9\text{B}$ . Channel  $\text{Be} + \text{H}$  appeared subdued accounting for about 2% of "white" stars. A probability ratio of the mirror channels  $^9\text{B} + \text{n}$  and  $^9\text{Be} + \text{p}$  is estimated to be  $6 \pm 1$ .  $^8\text{Be}(\text{g.s.})$  decays are presented in  $24 \pm 7\%$  of  $2\text{He} + 2\text{H}$  and  $27 \pm 11\%$  of the  $^3\text{He}$  of the  $^{11}\text{C}$  "white" stars.  $^9\text{B}$  decays are identified in "white" stars  $^{11}\text{C} \rightarrow 2\text{He} + 2\text{H}$  constituting 14% of the  $^{11}\text{C}$  "white" stars.

**Primary author:** Mr ZAITSEV, Andrew (JINR, LHE)

**Presenter:** Mr ZAITSEV, Andrew (JINR, LHE)

**Track Classification:** Experimental Nuclear Physics