

**Adler function, Bjorken Sum Rule,  
the Crewther-Broadhurst-Kataev relation and the  
 $\{\beta\}$ -expansion with different fermion representations of  
the gauge group of order  $O(\alpha_s^4)$**

P. A. Baikov<sup>1</sup>, K. G. Chetyrkin<sup>2</sup>, S. V. Mikhailov<sup>3</sup>

based on

**PRL 104, 132004 (2010), JHEP 04(2017)169, [B.Ch.M] in preparation**

<sup>1</sup>Skobeltsyn Institute of Nuclear Physics, Moscow State University, Moscow 119991, Russia

<sup>2</sup>Institut für Theoretische Teilchenphysik, KIT, D-76128 Karlsruhe, Germany

<sup>3</sup>Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, Dubna, Russia

**Advances in Quantum Field Theory 2021, BLTP**

## What are Adler function, Bjorken Sum Rule, the Crewther relation

There are renorm-group invariant single scale  $Q^2$  quantities  $D$ ,  $C^{\text{Bjp}}$ :

Adler function

$$d_R D(\mathbf{a}_s) = D_A = -12\pi^2 Q^2 \frac{d}{dQ^2} \Pi(Q^2); \quad Q^2 = -q^2$$

Bjorken polarized Sum Rule

$$\frac{1}{6} \left| \frac{g_A}{g_V} \right| C^{\text{Bjp}}(\mathbf{a}_s) + \text{high twist} = S_{\text{NS}}^{\text{Bjp}}(Q^2) = \int_0^1 [g_1^{lp}(x, Q^2) - g_1^{ln}(x, Q^2)] dx$$

Crewther relation

–a plausible **conjecture** [Crewther 1972,1997] inspired by conformal symmetry

$$D(\mathbf{a}_s)_{\text{ns}} \cdot C^{\text{Bjp}}(\mathbf{a}_s) = 1 + \beta(\mathbf{a}_s) K(\mathbf{a}_s), \text{ where } K(\mathbf{a}_s) - \text{polynom in } a_s = \frac{\alpha_s}{4\pi}$$

[D.Broadhurst,A.Kataev,PLB1993]-Crucial 3-loop analysis in  $\overline{\text{MS}}$ -scheme

[P.Baikov,K.Chetyrkin,J.Kühn, PRL2010]- confirmation in  $O(a_s^4)$ .

# OUTLINE

1. **Intro**: What is the  $\{\beta\}$ -**expansion** for RG-invariants and what does it express?
2. How to apply the  $\{\beta\}$ -**expansion**?  
To understand and to control the corresponding PT series in each expansion order, etc.
3. How to obtain the  $\{\beta\}$ -**expansion** from RQCD (QCD with different fermion Representations of gauge group), results for **Adler**  $D_{ns}(\mathbf{a}_s)$  and **Bjorken**  $C^{Bjp}(a_s)$ .
4. **Crewther-Broadhurst-Kataev relation** and its corollaries from  $\{\beta\}$ -**expansion** point of view.
5. **Conclusions**

# Motivation for the revision of series representation

we consider 1-scale  $Q^2$  **RG-INVARIANT** quantities at  $Q^2 = \mu_R^2$ , e.g.,  $D_{ns}$


"Wild" approach:  $\forall d_n$  - numbers,  
taken **wholly**

$$D(a_s) \sim 1 + a_s d_1 + a_s^2 d_2 + a_s^3 d_3 + \dots$$



**Delicate** approach:  $\forall d_n$  has an  
intrinsic structure due to  **$a_s$ -renorm.**

$$D(a_s) \sim 1 + \hat{M}(a_s, \{\beta_i\}) \leftarrow 2D \text{ matrix}$$



$$d_2 = 31.77 - 1.84n_f;$$

$$d_3 = 1164.8 - 270.1n_f - 5.5n_f^2;$$

$$d_4 = 34765 - 8806.4n_f + 481.3n_f^2 - 2.56n_f^3.$$

$$d_2 = \beta_0 d_2[1] + d_2[0];$$

$$d_3 = \beta_0^2 d_3[2] + \beta_1 d_3[0, 1] + \beta_0 d_3[1] + d_3[0]$$



$$d_4 = \beta_0^3 d_4[3] + \beta_2 d_4[0, 0, 1] + \dots$$

$\rightarrow$  series becomes "thick"

the decomposition is named  **$\{\beta\}$ -expansion[MS2005-7]**  
it shows the **dynamic** knowledge of  $D$  exhibited as  **$a_s$ -renorm.**

# How to apply the $\{\beta\}$ -expansion? 1.(If we already have it)

Different pieces are appropriate for miscellaneous and can be cooked differently



Evident usage of the  $\{\beta\}$ -expansion is the different kinds of **optimization**: one can change the contributions of different origins of  $a_s$ -renorm playing with the choice of  $\mu_R^2$ . E.g., well-known **BLM approach** [Brodsky et al 1983]:

$$d_2 = \underline{\beta_0 d_2[1]} + d_2[0] \rightarrow d_2[0] \text{ at } \mu_R^2 \rightarrow \mu_{BLM}^2 = \exp(-d_2[1]/d_1) \mu_R^2$$

$$\beta_0 = 11/3 C_A - 4/3 T_R n_f \Big| \text{ profit at } \underline{\beta_0 d_2[1]} \gg d_2[0]$$

$$d_3 = \underline{\beta_0^2 d_3[2]} + \beta_1 d_3[0, 1] + \beta_0 d_3[1] + d_3[0] \rightarrow d_3[0]$$

$$d_n = \underline{\beta_0^{n-1} d_n[n-1]} + \dots + d_n[0] \rightarrow d_n[0]$$

**generalized BLM:**  $D = 1 + \sum_1^N a_s^n(\mu_R^2) d_n \rightarrow D_0 = 1 + \sum_1^N a_s^n(\mu_{opt}^2) d_n[0]$

But this “conformal limit” may be not an **optimized series** in any sense, for  $R(s)$  it makes the PT convergence **worse** in  $O(a_s^4)$  [A. Kataev&MS PRD2015]

## How to apply the $\{\beta\}$ -expansion 2.(If we already have it)

[MS2007, A.Kataev&MS PRD2015]  $(a_s, \mu^2) \rightarrow (a'_s, \mu'^2)$

$$\ln(\mu'^2/\mu^2) = t - t' \equiv \Delta(a') = \Delta_0 + a' \beta_0 \Delta_1 + (a' \beta_0)^2 \Delta_2 + \dots, \quad \boxed{\Delta_0 = d_2[1]/d_1}$$

↑ BLM

$$a^2 \cdot d_2 \rightarrow a'^2 \cdot [d'_2 = \beta_0 (d_2[1] - \Delta_0) + d_2[0]];$$

$$a^3 \cdot d_3 \rightarrow a'^3 \cdot [d'_3 = \beta_0^2 (d_3[2] - 2d_2[1]\Delta_0 + \Delta_0^2 - \Delta_1) + \beta_1 (d_3[0, 1] - \Delta_0) \\ + \beta_0 (d_3[1] - 2d_2[0]\Delta_0) + d_3[0]];$$

$$a^4 \cdot d_4 \rightarrow a'^4 \cdot [d'_4 = \beta_0^3 (d_4[3] - 3d_3[2]\Delta_0 \dots - \Delta_2) + \dots]$$

Fitting components  $\Delta_0, \Delta_1, \Delta_2, \dots$  of the normalization scale  $\mu'^2$  to adjust the elements  $d'_2, d'_3, d'_4, \dots$  following to any **optimization procedure**.

**The practice impact of the  $\{\beta\}$ -expansion reveals at  $\beta_0 \gg 1$ ,** different cases of optimization in  $O(a_s^3)$  [A.Kataev&MS2015].

**An optimization as numerical minimum** of all QCD corr. to  $C^{\text{Bjp}}(a_s)$ ,  $a_s c_1 + a_s^2 c_2 + a_s^3 c_3 + a_s^4 c_4$ , up to  $O(a_s^4)$  was realized in [D.Kotlorz&MS2019]. The effect is about **-20%** at  $\mu^2 \approx 3\text{GeV}^2$ .

## How to obtain the $\{\beta\}$ -expansion? 1. RQCD

We need in QCD with additional degrees of freedom - **d.o.f.**, e.g., fermion multiplets  
These **d.o.f.  $\{R\}$  interact following the universal gauge principle** entering  
only in intrinsic loops [**K.Chetyrkin PLB1997, D with MSSM gluinos  $n_{\tilde{g}}$** ].

$$T_R n_f, \frac{C_A}{2} n_{\tilde{g}}, \dots \xrightarrow{\text{general}} \{R\} \text{ [M.Zoller 2016] for } \beta(a_s, \{R\}),$$

$$D(a_s, \{R\}) \text{ or } C^{\text{Bjp}}(a_s, \{R\}) : \text{RQCD [P.A.B.\&K.G.Ch.\&S.V.M.]}$$

**d.o.f.:**

$\{R\}$ - any numbers of different quark representations [**K.Chetyrkin&M.Zoller 2017**]

$$\mathcal{L}_{\text{QCD}} = \dots + \sum_{r=1}^{N_{\text{rep}}} \sum_{q=1}^{n_{f,r}} \left\{ \frac{i}{2} \bar{\psi}_{q,r} \overleftrightarrow{\hat{D}} \psi_{q,r} - m_{q,r} \bar{\psi}_{q,r} \psi_{q,r} + g_s \bar{\psi}_{q,r} \hat{A}^a T^{a,r} \psi_{q,r} \right\},$$

$R = (q - \text{flavors}, r - \text{Representation})$

Lie Algebra:  $[T^{a,r}, T^{b,r}] = if^{abc} T^{c,r}; T_{ik}^{a,r} T_{kj}^{a,r} = \delta_{ij} C_{F,r}; T_{F,r} \delta^{ab} = \text{Tr} (T^{a,r} T^{b,r});$

$$d_R^{a_1 a_2 \dots a_n} = \frac{1}{n!} \sum_{\text{perm } \pi} \text{Tr} \left\{ T^{a_{\pi(1)},R} T^{a_{\pi(2)},R} \dots T^{a_{\pi(n)},R} \right\},$$

In the standard QCD  $D(\mathbf{a}_s, \mathbf{R})$  and  $C^{Bjp}(\mathbf{a}_s, \mathbf{R})$  at  $O(a_s^4)$  computed in [P. Baikov, K. Chetyrkin, J. Kühn, 2010] with the help of

1. reduction of FI's with  $1/D$  expansion [P. Baikov 2000...]
2. FORM [J. Vermaseren 1990 ...]
3. FORM package COLOR for color structures [T. van Ritbergen, A. Schellekens, J. Vermaseren 1999 ...]

Adding more fermions multiplets,  $\mathbf{R} \rightarrow \{\mathbf{R}\}$ ,

items 1./2. in the same way as in [2010]

item 3. extension of the package COLOR on multi-fermion-generation case [M. Zoller 2016]



## How to obtain the $\{\beta\}$ -expansion? 2.

Then one can decompose all  $\beta$ -terms explicitly following an algebraic procedure

[MS2017]: all 7 elements of  $\mathbf{d}_4$  and  $\mathbf{c}_4$  were obtained,

$$d_4 = \beta_0^3 d_4[3] + \beta_2 d_4[0, 0, 1] + \beta_1 \beta_0 d_4[1, 1] + \beta_0^2 d_4[2] + \beta_1 d_4[0, 1] + \beta_0 d_4[1] + d_4[0]$$

$$d_n = \underbrace{\beta_0^{n-1} d_n[n-1] + \dots + \beta_0 d_n[1] + d_n[0]}_{\mathbf{N}(n)}$$

$$\hat{M}(a_s, \{\beta_i\}) = \begin{pmatrix} \dots & a_s^{n-1} & & a_s^n & \dots \\ & \vdots & \mathbf{N} \begin{cases} d_n[0] \\ \beta_0 d_n[1] \\ \vdots \end{cases} & & \end{pmatrix}$$

$$\mathbf{N}(n) = \sum_{l=0}^{(n-1)} p(l) = \{1, 2, 4, 7, 12, \dots\} \sim \frac{\sqrt{6n}}{\pi} \cdot (p(n) \leftarrow \text{partition of numbers}) + \dots$$

Hardy-Ramanujan asymptotic for partition of numbers  $p(n) \sim \frac{1}{4n\sqrt{3}} \exp\left(\pi\sqrt{2n/3}\right)$

**Important!** We need new d.o.f. Only to perform the decomposition, after that we return from RQCD to the standard QCD,  $\{\mathbf{R}\} \rightarrow \mathbf{T}_R n_f$ .

The marked traces of the general gauge principle saved as  $\{\beta\}$ -expansion.

## How to obtain the $\{\beta\}$ -expansion? 3. Solving a set of equations

The key role plays the **set of zeros** of  $\beta_0(\{R\}), \beta_1(\{R\}), \beta_2(\{R\}), \dots$  and **zeros** of sets of these  $\beta_k$ .

E.g., in  $O(a_s^4)$   $\beta_0, \beta_1, \beta_2$  are defined on the axes of variables  $R_0, R_1, R_2$ :

1)  $\exists$  3D point  $\bar{R}_{0,1,2} : \beta_0(\{\bar{R}_{0,1,2}\}) = \beta_1(\{\bar{R}_{0,1,2}\}) = \beta_2(\{\bar{R}_{0,1,2}\}) = \mathbf{0}$ ,  
Then  $D(\bar{R}_{0,1,2}) = (a_s^2 d_2[0], a_s^3 d_3[0], a_s^4 d_4[0], \dots)$   
step by step

2)  $\exists$  line in 3D  $\beta_0(\{\bar{R}_{0,1}\}) = \beta_1(\{\bar{R}_{0,1}\}) = \mathbf{0}$ ,  
Then  $d_4(\bar{R}_{0,1}) = \beta_2(\{\bar{R}_{0,1}\}) \underline{d_4[0, 0, 1]} + d_4[0]$

3)  $\exists$  curve (line) in 3D  $\beta_0(\{\bar{R}_{0,2}\}) = \beta_2(\{\bar{R}_{0,2}\}) = \mathbf{0}$ ,  
Then  $d_4(\bar{R}_{0,2}) = \beta_1(\{\bar{R}_{0,2}\}) \underline{d_4[0, 1]} + d_4[0]$

...

Finally this reduces to the set of equations at non-zero determinant.

**In  $O(a_s^5)$  at 6-loop we also have single-valued solution of the similar set of equation [MS2017].**

## Crewther-Broadhurst-Kataev relation and its corollaries 1.

The elements of  $\{\beta\}$ -**expansion** provide the *appropriate “bricks”* to analyse **C-B-K relation**, which is **our main subject** here.

First time it was applied to C-B-K [A.Kataev&MS TMP2012]

$$D_{ns}(a_s) \cdot C^{\text{Bjp}}(a_s) = \mathbf{1} + \beta(a_s) \times \sum_{n=1} a_s^{n-1} K_n$$

$$K_1 = K_1[1], K_2 = K_2[1] + \beta_0 K_2[2], \quad K_3 = K_3[1] + \beta_0 K_3[2] + \beta_0^2 K_3[3] + \beta_1 K_3[1, 1]$$

1. The  $D_0 \cdot C_0^{\text{Bjp}} = \mathbf{1}$  –“conformal” part of the relation, here  $d_n \rightarrow d_n[0] \in D_0$ ,

$$c_k[0] + d_k[0] = (-)^k \det[D_0^{(k)}] \equiv (-)^k \begin{vmatrix} d_1 & 1 & 0 & \dots & 0 \\ d_2 & d_1 & 1 & \dots & 0 \\ d_3 & d_2 & d_1 & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 \\ d_{k-1} & \dots & \dots & d_1 & 1 \\ \mathbf{0} & d_{k-1} & d_{k-2} & \dots & d_2 & d_1 \end{vmatrix},$$

$$c_k[0] + d_k[0] = \text{Polynom}(d_{k-1}), \quad k = 2, 3, 4 - \text{Confirmations!}$$

$$c_5[0] + d_5[0] = \text{Polynom}(d_4), \quad k = 5 - \text{Prediction}$$

## Crewther-Broadhurst-Kataev relation and its corollaries 2.

2. The factorization of the  $\beta(\mathbf{a}_s)$ , taken wholly, sets the chain of conditions

$$\begin{aligned}
 K_1[1] &= d_2[1] + c_2[1] = d_3[0, 1] + c_3[0, 1] = \underline{c_4[0, 0, 1] + d_4[0, 0, 1]} = 3C_F \left( \frac{7}{2} - 4\zeta_3 \right) \\
 &= d_n \underbrace{[0, 0, \dots, 1]}_{n-1} + c_n \underbrace{[0, 0, \dots, 1]}_{n-1} \quad \text{Confirmations/Prediction}
 \end{aligned}$$

$$\begin{aligned}
 K_2[1] &= c_3[1] + d_3[1] + d_1(c_2[1] - d_2[1]) = \\
 &= \underline{c_4[0, 1] + d_4[0, 1]} + d_1(c_3[0, 1] - d_3[0, 1]) \quad \text{Confirmation} \\
 &= C_F^2 \left( -\frac{397}{6} - 136\zeta_3 + 240\zeta_5 \right) + C_F C_A \left( \frac{47}{3} - 16\zeta_3 \right) \\
 &= \underline{c_5[0, 0, 1] + d_5[0, 0, 1]} + d_1(c_4[0, 0, 1] - d_4[0, 0, 1]) \quad \text{Prediction} \\
 &= c_n \underbrace{[0, \dots, 1]}_{n-2} + d_n \underbrace{[0, \dots, 1]}_{n-2} + d_1(c_{n-1} \underbrace{[0, \dots, 1]}_{n-2} - d_{n-1} \underbrace{[0, \dots, 1]}_{n-2}).
 \end{aligned}$$

$$\begin{aligned}
 K_3[1] &= c_4[1] + d_4[1] + d_1(c_3[1] - d_3[1]) + d_2[0]c_2[1] + d_2[1]c_2[0] \quad \text{Prediction} \\
 &= \underline{c_5[0, 1] + d_5[0, 1]} + d_1(c_4[0, 1] - d_4[0, 1]) + d_2[0]c_3[0, 1] + c_2[0]d_3[0, 1] = \dots \\
 &= c_{n+1} \underbrace{[0, \dots, 1]}_{n-2} + d_{n+1} \underbrace{[0, \dots, 1]}_{n-2} + d_1 \left( c_n \underbrace{[0, \dots, 1]}_{n-2} - d_n \underbrace{[0, \dots, 1]}_{n-2} \right) + \\
 &\quad d_2[0] c_{n-1} \underbrace{[0, \dots, 1]}_{n-2} + c_2[0] d_{n-1} \underbrace{[0, \dots, 1]}_{n-2}
 \end{aligned}$$

## Crewther-Broadhurst-Kataev relation, the structure of $K$ – term

The universal form of the second term appears due to the cancellation of  $a_s^1$ - terms

$$K_{n \geq 3}[1] = c_{n+1}[1] + d_{n+1}[1] + d_1 (c_n[1] - d_n[1]) + \sum_{k=2}^{n-2} (d_k[0] c_{n+1-k}[1] + c_k[0] d_{n+1-k}[1]).$$

Partial results for  $K$ -term in order  $O(a_s^4)$

$$K_1[1] = d_2[1] + c_2[1] = 3C_F \left( \frac{7}{2} - 4\zeta_3 \right)$$

$$\begin{aligned} K_2[1] &= c_3[1] + d_3[1] + d_1 (c_2[1] - d_2[1]) \\ &= C_F^2 \left( -\frac{397}{6} - 136\zeta_3 + 240\zeta_5 \right) + C_F C_A \left( \frac{47}{3} - 16\zeta_3 \right) \end{aligned}$$

$$K_2[2] = c_3[2] + d_3[2] = 3C_F \left( \frac{163}{6} - \frac{76}{3}\zeta_3 \right)$$

$$K_3[1] = c_4[1] + d_4[1] + d_1 (c_3[1] - d_3[1]) + d_2[0] c_2[1] + d_2[1] c_2[0]$$

$$K_3[2] = c_4[2] + d_4[2] + d_1 (c_3[2] - d_3[2]) + d_2[1] c_2[1],$$

$$K_3[3] = c_4[3] + d_4[3],$$

$$K_3[1, 1] = c_4[1, 1] + d_4[1, 1]$$

# CONCLUSIONS

1. The  **$\{\beta\}$ -expansion** for PT series is invented and analyzed for the **Renormalization Group invariant** quantities.  
This allows to perform different optimizations of the PT series.
2. The elements of  **$\{\beta\}$ -expansion** can be determined within **RQCD** following to an algebraic procedure.
3. The **Crewther-Broadhurst-Kataev relation** is reproduced in order  $O(a_s^4)$ . The interesting relations between the elements of Adler  $D_{ns}$ , and Bjorken polarized SR  $C^{Bjp}$  are established in any orders of  $a_s$ .

## STORE, explicit form of $D$ -elements. 1

$$\begin{aligned}d_1 &= 3C_F; \quad d_2[1] = d_1 \left( \frac{11}{2} - 4\zeta_3 \right); \quad d_2[0] = d_1 \left( \frac{C_A}{3} - \frac{C_F}{2} \right); \\d_3[2] &= d_1 \left( \frac{302}{9} - \frac{76}{3}\zeta_3 \right); \quad d_3[0, 1] = d_1 \left( \frac{101}{12} - 8\zeta_3 \right); \\d_3[1] &= d_1 \left( C_A \left( -\frac{3}{4} + \frac{80}{3}\zeta_3 - \frac{40}{3}\zeta_5 \right) - C_F (18 + 52\zeta_3 - 80\zeta_5) \right); \\d_4[3] &= C_F \left( \frac{6131}{9} - 406\zeta_3 - 180\zeta_5 \right); \\d_4[1, 1] &= C_F \left( 385 - \frac{1940}{3}\zeta_3 + 144\zeta_3^2 + 220\zeta_5 \right); \\d_4[2] &= -C_F \left[ C_F \left( \frac{6733}{8} + 1920\zeta_3 - 3000\zeta_5 \right) + \right. \\&\quad \left. C_A \left( \frac{20929}{144} - \frac{12151}{6}\zeta_3 + 792\zeta_3^2 + 1050\zeta_5 \right) \right]; \\d_4[0, 0, 1] &= C_F \left( \frac{355}{6} + 136\zeta_3 - 240\zeta_5 \right); \\d_4[1] &= C_F \left[ -C_F^2 \left( \frac{447}{2} - 42\zeta_3 - 4920\zeta_5 + 5040\zeta_7 \right) + \right. \\&\quad \left. C_A C_F \left( \frac{3301}{4} - 678\zeta_3 - 2280\zeta_5 + 2520\zeta_7 \right) + \right. \\&\quad \left. C_A^2 \left( \frac{16373}{36} - \frac{17513}{3}\zeta_3 + 2592\zeta_3^2 + 3030\zeta_5 - 420\zeta_7 \right) \right], \\d_4[0, 1] &= -C_F \left[ C_A \left( \frac{139}{12} + \frac{1054}{3}\zeta_3 - 460\zeta_5 \right) + C_F \left( \frac{251}{4} + 144\zeta_3 - 240\zeta_5 \right) \right],\end{aligned}$$

## STORE, explicit form of $D_0$ , $C_0$ elements. 2

$$d_3[0] = d_1 \left( \left( \frac{523}{36} - 72\zeta_3 \right) C_A^2 + \frac{71}{3} C_A C_F - \frac{23}{2} C_F^2 \right) .$$

$$\begin{aligned} d_4[0] &= \tilde{d}_4[0] + \delta d_4 \\ &= C_F^4 \left( \frac{4157}{8} + 96\zeta_3 \right) - C_A C_F^3 \left( \frac{2409}{2} + 432\zeta_3 \right) + C_A^2 C_F^2 \left( \frac{3105}{4} + 648\zeta_3 \right) + \\ &\quad C_A^3 C_F \left( \frac{68047}{48} + \frac{8113}{2} \zeta_3 - 7110\zeta_5 \right) + \delta d_4 , \end{aligned}$$

$$\delta d_4 = -\frac{16}{dR} \left( d_F^{abcd} n_f d_F^{abcd} (13 + 16\zeta_3 - 40\zeta_5) + d_F^{abcd} d_A^{abcd} (-3 + 4\zeta_3 + 20\zeta_5) \right)$$

$$c_3[0] = c_1 \left( \left( \frac{523}{36} - 72\zeta_3 \right) C_A^2 + \frac{65}{3} C_F C_A + \frac{C_F^2}{2} \right)$$

$$\begin{aligned} c_4[0] &= \tilde{c}_4[0] - \delta d_4 \\ &= -C_F^4 \left( \frac{4823}{8} + 96\zeta_3 \right) + C_A C_F^3 \left( \frac{3201}{2} + 432\zeta_3 \right) - C_A^2 C_F^2 \left( \frac{2055}{4} + 1944\zeta_3 \right) - \\ &\quad C_A^3 C_F \left( \frac{68047}{48} + \frac{8113}{2} \zeta_3 - 7110\zeta_5 \right) - \delta d_4 ; \end{aligned}$$