

Centrality determination in MPD at NICA AYSS-2021

Pedro Antonio Nieto Marín
Universidad Autónoma de Sinaloa, México
Dr. Alexey Aparin
Joint Institute for Nuclear Research, Dubna




JOINT INSTITUTE
FOR NUCLEAR RESEARCH



MexNICA

Centrality Determination / MC-Glauber Approach



Definition

Number of ancestors parameterization:

$$N_a(f) = fN_{part} + (1 - f)N_{coll} \quad (1)$$

Negative Binomial Distribution ($\mu = M^{max} / N_a^{max}$):

$$P(n) = \frac{\Gamma(n + k)}{\Gamma(n + 1)\Gamma(k)} \frac{(\mu/k)^n}{(\mu/k + 1)^{n+k}} \quad (2)$$

Glauber-based fit function:

$$F_{fit}(f, \mu, k) = \sum_1^{N_a} N_a \times P(N_{ch}) \quad (3)$$





Figure: Relation between the parameters f and k with their corresponding χ^2 with Bi+Bi collisions at 7.7, 9 and 9.46 GeV.

Parfenov, P., Idrisov, D., Luong, V., Taranenko, A. (2021). Relating Charged Particle Multiplicity to Impact Parameter in Heavy-Ion Collisions at NICA Energies. *Particles*, 4(2), 275–287. DOI: <https://doi.org/10.3390/particles4020024>

Definition


Centrality classes based on the multiplicity distribution:

$$c[\%] = \frac{\int_{N_{max}}^{N_i} \frac{dN_{ev}}{dN_{ch}} dN_{ch}}{\int_{N_{max}}^0 \frac{dN_{ev}}{dN_{ch}} dN_{ch}} \quad (4)$$



Multiplicity selection

- $p_T > 0.15$ GeV/c
- $|\eta| < 0.5$ and $|\eta| < 1.3$
- Only charged particles
- $N_{hits} > 16$
- Primary particles.
- $\sim 600,000$ reconstructed events in MpdRoot Framework.
- Bi+Bi collisions at 7.7, 9 and 9.46 GeV using UrQMD.



Multiplicity distribution




Figure: Comparison of the multiplicity distributions at the two η ranges ($|\eta| < 0.5$ and $|\eta| < 1.3$) at $\sqrt{s_{NN}} = 7.7, 9$ and 9.46 GeV.

b , N_{coll} and N_{part} vs centrality ($|\eta| < 0.5$)




Figure: Comparison of the relation of impact parameter, N_{coll} and N_{part} with centrality of the three energies $\sqrt{S_{NN}} = 7.7, 9$ and 9.46 GeV ($|\eta| < 0.5$).

b , N_{coll} and N_{part} vs centrality ($|\eta| < 1.3$)




Figure: Comparison of the relation of impact parameter, N_{coll} and N_{part} with centrality of the three energies $\sqrt{S_{NN}} = 7.7, 9$ and 9.46 GeV ($|\eta| < 1.3$).

$|\eta| < 0.5$ and $|\eta| < 1.3$ comparison





Figure: Pseudorapidity comparison of the relation of impact parameter with centrality at the three energies $\sqrt{S_{NN}} = 7.7, 9$ and 9.46 GeV.

Centrality ranges comparison

Bi+Bi $\sqrt{s_{NN}}=9.46$ GeV, $|\eta|<0.5$, 1% ranges

- $p_T > 0.15$ GeV/c
- $|\eta| < 0.5$
- 5% centrality range vs 1% centrality range.



Centrality ranges comparison






Figure: 5% and 1% centrality ranges comparison of the impact parameter vs centrality at the three energies $\sqrt{s_{NN}} = 7.7, 9$ and 9.46 GeV.

UrQMD model

- $p_T > 0.15$ GeV/c
- $|\eta| < 0.5$
- Only charged particles
- $\sim 600,000$ events.
- Bi+Bi collisions at 7.7, 9 and 9.46 GeV using UrQMD test.f14 output files.



UrQMD model ($|\eta| < 0.5$)




Figure: UrQMD model and TPC results comparison of the impact parameter vs centrality at the three energies $\sqrt{S_{NN}} = 7.7, 9$ and 9.46 GeV ($|\eta| < 0.5$).

UrQMD model ($|\eta| < 1.3$)




Figure: UrQMD model and TPC results comparison of the impact parameter vs centrality at the three energies $\sqrt{S_{NN}} = 7.7, 9$ and 9.46 GeV ($|\eta| < 1.3$).

Parameterization comparison

- "Default"

$$N_a(f) = fN_{part} + (1 - f)N_{coll}$$

- "PSD"

$$N_a(f) = f - N_{part}$$

- "Npart"


$$N_a(f) = (N_{part})^f$$

- "Ncoll"

$$N_{coll}(f) = (N_{coll})^f$$

- "STAR"

$$N_a(f) = \frac{(1-f)}{2}N_{part} + fN_{coll}$$



- Better agreement with STAR and Default (0-40%).


- Problem with PSD parameterization.



JOINT INSTITUTE
FOR NUCLEAR RESEARCH

Different models comparison

- $p_T > 0.15$ GeV/c
- $|\eta| < 0.5$
- $N_{hits} > 16$
- **DCA cut:**
 $|DCA| < 0.5$
- **"Default"**
 $N_a(f) = fN_{part} + (1 - f)N_{coll}$



- Obtain PHSD model to compare.
- Better agreement with **PHSD, DCA correction and cut, $|\eta| < 1.3$** .



Future work

- Compare and explore Γ – *Fit* method.
- Finish analysis of new obtained reconstructed data from Bi+Bi collisions at 9.2 GeV.
- Compare results with other Monte Carlo generators (LAQGSM, PHSD and PHQMD) for Bi+Bi collisions at the four different energies (7.7, 9, 9.2 and 9.5 GeV).
- Fix PSD parameterization.

Thank you!



Backup slides



Abstract for AYSS-2021

We present a study of centrality class determination based on number of charged particles registered with Time Projection Chamber in Multi-Purpose Detector at NICA complex. Precise determination of centrality classes will allow to select the ion collisions within a certain class of initial conditions in order to study behaviour of some variables on the mean energy densities reached.

It will be important to study the QCD matter with respect to the geometric properties of the collision between nuclei, but these properties cannot be experimentally measured. Therefore, we compare in this work the centrality classes which are obtained using different sets of observables as proxies for centrality, They include the number of hits in the TPC and transverse momentum of registered particles in the TPC sub-detector for Bi+Bi collisions at $\sqrt{S_{NN}} = 7.7, 9$ and 9.46 GeV, Data generated by several Monte Carlo models are used. Results are discussed.

