

A NEW APPROACH TO THE THERMAL EVOLUTION OF NEUTRON STARS

Rodrigo Negreiros – IF/UFF

Fridolin Weber – SDSU

Stefan Schramm - FIAS





ACKNOWLEDGMENTS

- CAPES
- CNPq
- David Blaschke
- Organizes

INTRODUCTION

- Thermal evolution is driven by neutrino emissions from the core, and photon emission from the surface.
- Neutrino emissions strongly depend on the core composition.
- Depending on its mass, a neutron star may exhibit fast or slow cooling.







$$P = P(\rho, T = 0, \Omega = 0)$$

$$M = M(\rho, T = 0, \Omega = 0)$$

$$R = R(\rho, T = 0, \Omega = 0)$$



$$\varepsilon = \varepsilon(\rho, T)$$


$$K = K(\rho, T)$$

$$c_v = c_v(\rho, T)$$


TRADITIONAL PICTURE

- Structure and composition frozen in time - “Frozen In”
- Thermal properties only change due to temperature evolution
- Dynamics => TEMPERATURE (ONLY !!)

STRUCTURE/COMPOSITION MAY NOT BE STATIC









Accretion



Magnetic/Spin evolution





STUDIES

• Structure and Composition



PREVIOUS STUDIES

- Direct Urca and superfluidity



A NEW APPROACH

- Consider a dynamic structure and composition.
- Go beyond spherically symmetric stars.

$$P = P(\rho, T = 0, \Omega = 0)$$

$$M = M(\rho, T = 0, \Omega = 0)$$

$$R = R(\rho, T = 0, \Omega = 0)$$

$$\varepsilon = \varepsilon(\rho, T)$$

$$K = K(\rho, T)$$

$$c_v = c_v(\rho, T)$$



$$P = P(\rho, T = 0, \Omega(t), B(t))$$

$$M = M(\rho, T = 0, \Omega(t), B(t))$$

$$R = R(\rho, T = 0, \Omega(t), B(t))$$

$$\varepsilon = \varepsilon(\rho, T, \Omega(t), B(t))$$

$$K = K(\rho, T, \Omega(t), B(t))$$

$$c_v = c_v(\rho, T, \Omega(t), B(t))$$

PROOF OF CONCEPT: - STRUCTURE OF ROTATING NEUTRON STARS

- Metric $ds^2 = -e^{\gamma+\rho} dt^2 + e^{2\alpha} (dr^2 + r^2 d\theta^2) + e^{\gamma-\rho} r^2 \sin^2 \theta (d\phi - \omega dt)^2$
- Sources $T^{\mu\nu} = (\epsilon + p) u^\mu u^\nu + p g^{\mu\nu} + \text{magnetic terms}$
- Einstein's Equation $G^{\mu\nu} = R^{\mu\nu} - \frac{1}{2} g^{\mu\nu} R = 8\pi T^{\mu\nu}$
- Hydrostatic Equilibrium $dp - (\epsilon + p) [d \ln u^t + u^t u_\phi d\Omega] = 0 + \text{magnetic terms}$


PROOF OF CONCEPT: - STRUCTURE OF ROTATING NEUTRON

- Magnetic Braking model

$$\dot{\Omega} = -K \Omega^n$$

NOW, WE HAVE OBTAINED THE
STRUCTURE/COMPOSITION AS A
FUNCTION OF TIME!

ONTO THE COOLING....



THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS

$$P = P(\rho, T = 0, \Omega(t), B(t))$$

$$M = M(\rho, T = 0, \Omega(t), B(t))$$

$$R = R(\rho, T = 0, \Omega(t), B(t))$$

$$\varepsilon = \varepsilon(\rho, T, \Omega(t), B(t))$$

$$K = K(\rho, T, \Omega(t), B(t))$$


$$c_v = c_v(\rho, T, \Omega(t), B(t))$$




$$\begin{aligned} \partial_r \tilde{H}_{\bar{r}} + \frac{1}{r} \partial_{\theta} \tilde{H}_{\bar{\theta}} &= -r e^{\phi+2\omega} \left(\frac{1}{\Gamma} e^{2\nu} \varepsilon + \Gamma C_V \partial_t \tilde{T} \right) \\ &\quad - r \Gamma U e^{\nu+2\phi+\omega} \left(\partial_r \Omega + \frac{1}{r} \partial_{\theta} \Omega \right), \\ \partial_r \tilde{T} &= -\frac{1}{r\kappa} e^{\nu-\phi} \tilde{H}_{\bar{r}} - \Gamma^2 U e^{-\nu+\phi} \tilde{T} \partial_r \Omega, \\ \frac{1}{r} \partial_{\theta} \tilde{T} &= -\frac{1}{r\kappa} e^{-\nu-\phi} \tilde{H}_{\bar{\theta}} - \Gamma^2 U e^{-\nu+\phi} \tilde{T} \frac{1}{r} \partial_{\theta} \Omega \\ \Gamma U \partial_t \tilde{T} &= -\frac{1}{r\kappa} e^{-\omega-\phi} \tilde{H}_{\bar{\varphi}}, \end{aligned}$$

THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS

- $M_0 = 1.8 M_{\text{sun}}$, $\Omega_0 = 500$ Hz



$$\dot{\Omega} = 10^{-15} \text{ s}^{-2}$$




$$\dot{\Omega} = 10^{-13} \text{ s}^{-2}$$


THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS

• $M_0 = 1.8 M_{\text{sun}}$, $\Omega_0 = 500 \text{ Hz}$


$$\dot{\Omega} = 10^{-13} \text{ s}^{-2}$$



0.10 years



36 years




8.0E5 years


THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS

• $M_0 = 1.8 M_{\text{sun}}$, $\Omega_0 = 500 \text{ Hz}$


$$\dot{\Omega} = 10^{-15} \text{ s}^{-2}$$



0.10 years



10.05 years




22.45 years


THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS

• $M_0 = 1.8 M_{\text{sun}}$, $\Omega_0 = 500 \text{ Hz}$


$$\dot{\Omega} = 10^{-13} \text{ s}^{-2}$$



0.10 years



10.05 years




22.45 years


THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS

- $M_0 = 1.8 M_{\text{sun}}$, $\Omega_0 = 500 \text{ Hz}$

$$\dot{\Omega} = 10^{-15} \text{ s}^{-2}$$




31.67 years




5.0E5 years

THERMAL EVOLUTION OF AXIS-SYMMETRIC NEUTRON STARS


- $M_0 = 1.8 M_{\text{sun}}$, $\Omega_0 = 500 \text{ Hz}$ -> **MICROSCOPIC PROPERTIES**



Proton Singlet GAP




Neutron Triplet GAP



Neutron Singlet GAP

THERMAL-ROTATIONAL EVOLUTION



PERSPECTIVES

- We are now in a position to deal fully with axis-symmetric neutron stars thermal evolution.
- We can also consider a dynamic structure/composition.
- Investigate different neutron stars evolution:
 - Magnetic
 - Spin-Up/accreting