



Rho meson distribution amplitudes in QCD Sum Rules

Alexander Pimikov¹

in collaboration with S. V. Mikhailov¹, and N. G. Stefanis²

Bogoliubov Lab. Theor. Phys., JINR (Dubna, Russia)¹

ITP-II, Ruhr-Universität (Bochum, Germany)²

Motivation:

- B-meson decays: $B \rightarrow \rho l \nu$, $\bar{B}^0 \rightarrow \rho^0 \gamma$
- CKM matrix constrains
- ρ -meson electro-photoproduction

Leading twist ρ -meson DA definitions:

$$\langle 0 | \bar{d}(z) \gamma_\mu u(0) | \rho(P, \lambda) \rangle = f_\rho P_\mu \int_0^1 dx e^{ix(zp)} \varphi_\rho^L(x) + \text{higher twists}$$

$$\langle 0 | \bar{d}(z) \sigma_{\mu\nu} u(0) | \rho(P, \lambda) \rangle = i f_\rho^\top (\varepsilon_\mu^{(\lambda)} P_\nu - \varepsilon_\nu^{(\lambda)} P_\mu) \int_0^1 dx e^{ix(zp)} \varphi_\rho^\top(x) + \dots$$


- Typical characteristic used to describe observables:

$$\int_0^1 dx \varphi_\rho^{L,\top}(x)/x \text{ (Inverse moment) and } \int_\epsilon^1 dx \varphi_\rho^{L,\top}(x) f(x) \text{ with } \epsilon \simeq 0.1 - 0.5 .$$

- DA evolution with μ^2 , according to ERBL equation **[1979-1980]**.

Gegenbauer expansion of pion DA: $\varphi_\pi(x, \mu^2) \Leftrightarrow a_2, a_4, \dots, a_n$

$$\varphi_\pi(x, \mu^2) = 6x\bar{x}(1 + a_2(\mu^2)C_2^{3/2}(x - \bar{x}) + a_4(\mu^2)C_4^{3/2}(x - \bar{x}) + \dots)$$



QCD SR Approach

Determination of parameters from agreement of two correlator calculations [SVZ'79]:

$$\Pi(-q^2) = \int d^4x e^{-iqx} \langle 0 | J_1(0) J_2(x) | 0 \rangle .$$

- Dispersion relation: decay constants f_h and masses m_h ,

$$\Pi_{\text{had}}(Q^2) = \int_0^\infty \frac{\rho_{\text{had}}(s) ds}{s + Q^2} + \text{subtractions} .$$

- Model spectral density: $\rho_{\text{had}}(s) = f_h^2 \delta(s - m_h^2) + \rho_{\text{pert}}(s) \theta(s - s_0)$.

- Operator product expansion:

$$\Pi_{\text{OPE}}(Q^2) = \Pi_{\text{pert}}(Q^2) + \sum_n C_n \frac{\langle 0 | : O_n : | 0 \rangle}{Q^{2n}} .$$

- Condensates $\langle 0 | : O_n : | 0 \rangle \equiv \langle O_n \rangle \neq 0$ (next slides).

QCD SR reads

$$\Pi_{\text{had}}(Q^2, m_h, f_h) = \Pi_{\text{OPE}}(Q^2) .$$

Introducing condensates in QCD calculations

$$\langle 0 | T (\bar{q}_B(0) q_A(x)) | 0 \rangle = \langle 0 | : \bar{q}_B(0) q_A(x) : | 0 \rangle - i \hat{S}_{AB}(x)$$

QCD PT

$$\langle 0 | : \bar{q} q : | 0 \rangle \stackrel{\text{def}}{=} 0$$

QCD SR

$$\langle 0 | : \bar{q}_A(0) q_A(x) : | 0 \rangle = \langle \bar{q} q \rangle = \text{CONST} \neq 0$$



[SVZ'79]

Condensates

Decay constants,
masses of hadrons

NLC QCD SR

$$\langle 0 | : \bar{q}(0) q(x) : | 0 \rangle = F_S(x^2) + \hat{x} F_V(x^2)$$




M&R '86

Nonlocal condensates

Distribution Amplitudes (DA),
Form Factors

$$\langle \bar{q}_B(0) q_A(x) \rangle = \frac{\delta_{AB}}{4} \left[\langle \bar{q} q \rangle + \frac{x^2}{4} \frac{\langle \bar{q} D^2 q \rangle}{2} + \dots \right] + i \frac{\hat{x}_{AB}}{4} \frac{x^2}{4} \left[\frac{2\alpha_s \pi \langle \bar{q} q \rangle^2}{81} + \dots \right]$$


Diagrams for $\langle T (J_\nu(z) J_\mu(0)) \rangle$



- Quarks run through vacuum with nonzero momentum $k \neq 0$ with average k^2 :

$$2\langle k^2 \rangle = \frac{\langle \bar{q} D^2 q \rangle}{\langle \bar{q} q \rangle} = \lambda_q^2 = 0.40(5) \text{ GeV}^2$$

Nonlocality from Lattice data



Nonlocality of quark condensates $\lambda_q^2 = 0.42(8) \text{ GeV}^2$ from lattice data of Pisa group (Di Giacomo et.al., PRD(1999)) in comparison with **local limit**.

- Even at $|z| \simeq 0.5 \text{ fm}$ nonlocality is quite important!

QCD SR for ρ -meson DA

QCD SR technique for correlator of two vector current leads to SR for ρ -DA $\varphi_\rho^L(x)$:

$$f_\rho^2 \varphi_\rho(x) e^{-m_\rho^2/M^2} + f_{\rho'}^2 \varphi_{\rho'}(x) e^{-m_{\rho'}^2/M^2} = \int_0^{s_0} \rho_{\text{pert}}(s, x) e^{-s/M^2} ds + \Phi_\rho(x, M^2),$$

where

$$\Phi_\rho(x, M^2) = -\varphi_{4Q} + \varphi_T + \varphi_V + \varphi_G,$$


$$\Phi_\pi(x, M^2) = +\varphi_{4Q} + \varphi_T + \varphi_V + \varphi_G.$$

The largest nonperturbative term:

$$\varphi_{4Q}^{\text{NLC}} \sim x\theta(\Delta - x) \xrightarrow{\text{loc. lim}} \varphi_{4Q}^{\text{loc}} \sim \delta(x),$$

is defined by scalar quark condensate,

where $\Delta = \lambda_q^2/M^2 \in [0.1, 0.3]$.





● Nonperturbative contribution has **singularities**.

SRs for integral characteristics ($\int_0^1 dx \varphi(x) (1 - 2x)^N$, $\int_0^1 dx \varphi(x)/x$) take into account all condensates and have less model dependence.



● In end-point region only 4-quark condensate φ_{4Q} contributes without any singularities. That allows us to study slope $\varphi'(0)$ and $\int_0^1 dx \varphi(x)/x$.

ρ vs π $DA_{\mu=1}$ GeV from NLC QCD SR



ρ (here)



π (BMS, Bakulev et.al., PLB508(2001))



φ_ρ^L DA from QCD SR with NLC



- ◆ NLC QCD SR, (here)
- ◆ Asymptotic
- - -■ QCD SR, Ball&Jones 07
- ⋯□ Lattice, Arthur et.al. 11
- · - □ AdS/QCD, Ahmady&Sandapen 13
- · - ▽ Light-front quark model, Choi&Ji 07
- - -● Instanton liquid model, Dorokhov 07

$\mu = 1 \text{ GeV}$

● DA model and bunch were obtained using Gaussian condensate model with single nonlocality parameter $\lambda_q^2 = 0.4 \text{ GeV}^2$.

● Higher Gegenbauer coefficients $a_{n \geq 6} = 0$ are assumed to be equal 0 — this does not contradict QCD SR.

SR for transverse DA φ_ρ^T

$$\Pi_N^{\mu\nu;\alpha\beta}(q) \equiv \int d^4x e^{iqx} \langle 0 | T(J_{\mu\nu}^N(x) J_{\alpha\beta}^0(0)) | 0 \rangle = \sum_{i=1}^6 C_i P_i^{\mu\nu;\alpha\beta}$$

$$J_{\mu\nu}^N(x) \equiv \bar{u}(x) \sigma_{\mu\nu} (z \nabla)^N d(x)$$

Mixed SR, Ball&Braun 96:

$$\rho + b_1 \sim \Pi_N^{\mu\nu;\alpha\beta}(q) g_{\mu\alpha} n_\nu n_\beta \sim C_1 - C_2$$

$$\rho + \text{higher twists} \sim C_1$$




$$b_1 + \text{higher twists} \sim C_2$$

Pure SR, Bakulev&Mikhailov 01:




$$\rho \sim C_1 + C_4 = \frac{1}{2}(C_1 - C_2) + \Delta_{4Q}$$

$$b_1 \sim -(C_2 + C_4) = \frac{1}{2}(C_1 - C_2) - \Delta_{4Q}$$




Where Δ_{4Q} is total 4-quark condensate contribution defined by following diagrams:






4-quark condensate terms to SR for φ_ρ^T -DA



Local Condensates ($\lambda_q^2 \rightarrow 0$)



NonLocal Condensates ($\lambda_q^2 = 0.4 \text{ GeV}^2$)



For typical value of Borel parameter $M^2 = 1 \text{ GeV}^2$

Conclusions

- Using QCD SR with nonlocal condensate we recalculated the leading twist longitudinal and transverse ρ -meson DAs.
- The longitudinal ρ -meson DA was found to be close to Light-front quark model result [Choi&Ji 07] and the asymptotic form.
- Calculated 4-quark condensate contribution to transverse ρ -meson DA appeared to be non-zero in the end-point x -region.
- Our result for transverse ρ -meson DA is preliminary. Further study is still in process.

Thank you for attention!