

Applicability assessment of Fe₃O₄-SiO₂-Au nanoparticles for the cancer radiotherapy

Friday, 28 October 2022 15:05 (15 minutes)

The object of our research is a Fe₃O₄-SiO₂-Au nanoparticles. The electron microscopy study showed that the nanoparticles had a core-shell structure. The average size of the magnetite core was 227±7 nm, the thickness of the protective shell made of SiO₂ was 2 nm, the average size of gold nanoparticles covering the core was 16±0.6 nm.

To estimate the output of secondary electromagnetic radiation during irradiation of the Fe₃O₄-SiO₂-Au nanoparticles, an experimental unit was constructed.

A scintillation gamma-ray spectrometer based on a NaI(Tl) 78×78 crystal was used as a detector. The relative efficiency of registration at the ¹³⁷Cs 662 keV line was 25%, and the resolution at the ¹³⁷Cs 662 keV line was 9%. A collimated beam of gamma radiation from ¹³⁷Cs was directed to the cuvette perpendicular to the axis of the spectrometer. The activity of the ¹³⁷Cs radionuclide source was 99 MBq.

To take into account the background component of the obtained data, we measured the energy spectrum of the blank sample. A water sample was used as a blank. Measurements were carried out sequentially –replacing the test and blank samples. In the end of each measurement stage, the energy spectra of the standard gamma-ray sources (²⁴¹Am, ¹³⁷Cs) were measured.

To process the experimental data obtained, the measured energy spectra were conditionally divided into several ranges. The width of ranges was selected in accordance with the energy resolution of the spectrometer. The intensity of the analytical signal of secondary radiation of the Fe₃O₄-SiO₂-Au nanoparticles generated by gamma-ray of the ¹³⁷Cs source was determined in accordance to the equation:

where: –the intensity of the analytical signal of secondary electromagnetic radiation; –the median value of the intensity of the analytical signal in the spectrum of the sample, –the median value of the intensity of the analytical signal in the spectrum of the blank sample, –the number of the energy range.

The absolute error of the data obtained was determined by the method of calculating the errors of indirect measurements according to the equation:

where: –the absolute error of determining the intensity of the analytical signal of secondary electromagnetic radiation; –the absolute error of determining the median value of the intensity of the analytical signal in the spectrum of the sample; –the absolute error of determining the median value of the intensity of the analytical signal in the spectrum of the blank sample.

As a result, the spectrum of secondary electromagnetic radiation generated by irradiation of Fe₃O₄-SiO₂-Au nanoparticles was obtained. It was shown that the secondary radiation is caused by the photoelectric and the Compton effects.

We acknowledge the Russian Ministry of Science and Higher Education. Synthesis and structural studies of nanoparticles were carried out with financial support the Russian Ministry of Science and Higher Education, State Order No. 0657-2020-0013. Radiological studies were carried out with financial support of the Ministry of Russian Ministry of Science and Higher Education, State Order No. 0205-2021-0002

Primary authors: MARKIN, Nikita (Far Eastern Federal University; Institute of chemistry of the Far Eastern Branch of the Russian Academy of Sciences); IVANNIKOV, Sergey (Institute of chemistry of the Far Eastern Branch of the Russian Academy of Sciences); OGNEV, Alexey (Far Eastern Federal University); AFREMOV, Leonid (Far Eastern Federal University); SAMARDAK, Alexander (Far Eastern Federal University)

Presenter: MARKIN, Nikita (Far Eastern Federal University; Institute of chemistry of the Far Eastern Branch of the Russian Academy of Sciences)

Session Classification: Online poster session

Track Classification: Condensed Matter Physics