

RADIATION CORRECTIONS TO PROMPT PHOTON PRODUCTION IN COMPTON SCATTERING OF QUARK-GLUON $qg \longrightarrow q\gamma$ AND ANNIHILATION OF QUARK-ANTIQUARK PAIR $q\bar{q} \longrightarrow g\gamma$ PROCESSES

Tuesday 25 October 2022 15:30 (15 minutes)

```
\documentclass{article}
\usepackage{amssymb}
\usepackage{amsmath}
\usepackage[dvips]{graphicx}
\begin{document}
\begin{center}
\textbf{RADIATION CORRECTIONS TO PROMPT PHOTON PRODUCTION IN}
\textbf{COMPTON SCATTERING OF QUARK-GLUON } $qg \rightarrow q\gamma$ \textbf{ AND ANNIHILATION OF}
\textbf{QUARK-ANTIQUARK PAIR } $q\bar{q} \rightarrow g\gamma$ \textbf{ PROCESSES }
\textbf{}
\textbf{M.R. ALIZADA}
\textbf{}
\textit{Department of Theoretical Physics, Baku State University}
\textit{str. Z.Khalilov, 23, Az-1148 Baku, Azerbaijan E-mail: }
\textit{mohsunalizade@gmail.com}
\end{center}
\end{document}
```

Radiation corrections to processes of Compton scattering of quark-gluon:

1. $q\gamma \rightarrow q\gamma$, 2. $q\gamma \rightarrow qg$, 3. $g\gamma \rightarrow q\bar{q}$,
4. $q\gamma \rightarrow qg\gamma$, 5. $qg \rightarrow g\gamma\gamma$, 6. $qg \rightarrow qg\gamma$ and
7. $g\gamma \rightarrow q\bar{q}\gamma$ and annihilation of quark-antiquark pair:

1. $q\bar{q} \rightarrow \gamma\gamma$, 2. $q\bar{q} \rightarrow q\bar{q}\gamma$,
 3. $q\bar{q} \rightarrow g\gamma\gamma$ and 4. $q\bar{q} \rightarrow gg\gamma$ without and
- with taking into account of polarization of quark was considered.

Dependencies of differential cross section of subprocesses on energy of colliding protons \sqrt{s} , transverse momentum p_T , cosine of scattering angle $\cos(\theta)$ and y of photons, x_T were investigated.

Differential cross section of considered subprocesses decreases as the transverse momentum increases. The following results were obtained:

$$\frac{d\sigma_1}{dydp_T^2} \gg \frac{d\sigma_2}{dydp_T^2} \gg \frac{d\sigma_3}{dydp_T^2} \text{ and } \frac{d\sigma_4}{dydp_T^2} \gg \frac{d\sigma_5}{dydp_T^2} \gg \frac{d\sigma_6}{dydp_T^2} \gg \frac{d\sigma_7}{dydp_T^2}$$

Compton scattering process and $\frac{d\sigma_2}{dydp_T^2} \gg \frac{d\sigma_3}{dydp_T^2} \gg \frac{d\sigma_4}{dydp_T^2}$ for annihilation of quark-antiquark pair process.

It was been determined that, contributions of corrections to differential cross section of Compton scattering of quark-gluon process is significant than contributions of corrections to differential cross section of annihilation of quark-antiquark pair process.

The double spin asymmetry A_{LL} of subprocesses $q\bar{q} \rightarrow \gamma\gamma$, $q\bar{q} \rightarrow g\gamma\gamma$ and $q\bar{q} \rightarrow gg\gamma$ of annihilation process are independent of \sqrt{s} , p_T and $\cos(\theta)$.

The double spin asymmetry expression for these subprocesses is as follows

$A_{LL} = -\lambda_1 \lambda_2$. Doublespin asymmetry A_{LL} of subprocess $q\bar{q} \rightarrow q\bar{q}\gamma$ of annihilation process increases (decreases) with increasing transverse momentum for $\lambda_1 \lambda_2 < 0$ ($\lambda_1 \lambda_2 > 0$) and reach plateau at certain p_T .
 The value of this p_T increases with increasing of absolute value of $\lambda_1 \lambda_2$.

Author: ALIZADA, Mohsun (Baku State University)

Presenter: ALIZADA, Mohsun (Baku State University)

Session Classification: High Energy Physics

Track Classification: High Energy Physics