

Nonanalytic relativistic r -mode instability windows

Wednesday, 1 March 2023 16:10 (30 minutes)

Detectability of the r -mode gravitational-wave signal depends on the interplay between the mode amplification by the CFS instability and its damping by dissipative mechanisms, operating in the stellar matter. Those stellar parameters - usually, the angular velocity, Ω , and redshifted temperature, T^∞ , - for which the mode is unstable, define the r -mode instability window. We revisit this problem in nonbarotropic neutron stars, accounting for the previously overlooked relativistic r -mode nonanalytic behavior (in Ω) and enhanced energy dissipation due to diffusion in superconducting stellar matter. We show that, at slow rotation rates, relativistic r -modes are amplified by the CFS instability weaker than Newtonian ones, while their viscous and diffusive dissipation is, instead, significantly more efficient. At realistic rotation rates relativistic and Newtonian r -mode amplification by CFS mechanism and damping by shear viscosity become comparable, while the relativistic mode damping by diffusion and bulk viscosity remain significantly stronger than nonrelativistic ones. As a result, accounting simultaneously for diffusion and relativistic r -mode nonanalyticity drastically changes the r -mode instability window as compared to the Newtonian one. This effect is of paramount importance for the interpretation of the future gravitational-wave observations and understanding of the r -mode physics in general.

Primary authors: KANTOR, Elena (Ioffe Institute); KRAAV, Kirill (Ioffe Institute); GUSAKOV, Mikhail (Ioffe Institute)

Presenter: KRAAV, Kirill (Ioffe Institute)