

G.P. PROKHOROV ^{1,2}

O.V. TERYAEV ^{1,2}

V.I. ZAKHAROV ^{2,1}

¹ JOINT INSTITUTE FOR NUCLEAR
RESEARCH (JINR), BLTP, DUBNA

² NRC KURCHATOV INSTITUTE,
MOSCOW

BASED ON WORKS:

[1] PHYS. REV. LETT.,
129(15):151601, 2022.

[2] ARXIV: 2211.03865

INFINITE AND FINITE NUCLEAR
MATTER (INFINUM-2023),
JINR, DUBNA,
27 FEBRUARY TO 3 MARCH 2023

KINEMATICAL VORTICAL EFFECT

CONTENTS

- Introduction
- Gravitational chiral anomaly in hydrodynamics:
 - Derivation of the general relation for Kinematical Vortical Effect (KVE) (a la Son&Surowka)
- Verification: spin $\frac{1}{2}$
- Verification: **spin 3/2 fields**
 - Rarita-Schwinger-Adler (RSA) theory
 - **KVE vs anomaly**
- Conclusion

PART 1

INTRODUCTION

GRADIENT EXPANSION IN HYDRODYNAMICS

- **Hydrodynamics** is constructed as a **gradient expansion**. E.g. for an ideal fluid: (signature (-1,1,1,1))

$$T^{\mu\nu} = (\varepsilon + p)u^\mu u^\nu + p\eta^{\mu\nu}$$

- For a viscous fluid, linear gradients arise: **[Landau, Lifshitz, Fluid Mechanics, Vol. 6]**

Shear viscosity

Bulk viscosity

From string theory:

$$\eta/s \geq \hbar/(4\pi k_B)$$

$$\tau_{\mu\nu} = -\eta P_\mu^\alpha P_\nu^\beta (\partial_\alpha u_\beta + \partial_\beta u_\alpha) - \left(\zeta - \frac{2}{3}\eta\right) P_{\mu\nu} \partial_\alpha u^\alpha$$

- **New phenomena** related to **vorticity** and **magnetic fields** (see talks of A. Roenko, Yu. Ivanov, V. Nesternko, D. Sychev). Two well-known effects:

Chiral magnetic effect

$$\text{CME: } j^\mu = C \mu_5 B^\mu$$

Experimental search is in progress: in HIC and condensed matter

Chiral vortical effect

$$\text{CVE: } j_A^\mu = C \mu^2 \omega^\mu$$

$$\omega^\mu = \frac{1}{2} \varepsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha u_\beta$$

CVE AND CME – NEW ANOMALOUS TRANSPORT


- **CME** and **CVE** are related to the gauge chiral **anomaly**:

[V. I. Zakharov, Lect. Notes Phys.871,295(2013), 1210.2186]

[D.T. Son, P. Surowka, PRL, 103 (2009) 191601]

$$\text{CME: } j^\mu = C \mu_5 B^\mu$$

$$\text{CVE: } j_A^\mu = C \mu^2 \omega^\mu$$



Quantum chiral anomaly


$$\langle \partial_\mu \hat{j}_A^\mu \rangle = -\frac{C}{8} \varepsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$$

- Interplay of **infrared** (hydrodynamics) and **ultraviolet** (anomalies) physics.

What about the **gravitational chiral anomaly**?

- The gravitational chiral anomaly grows **rapidly** with **spin**:

???



$$\langle \nabla_\mu \hat{j}_A^\mu \rangle_S = \frac{(S - 2S^3)}{96\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

[M. J. Duff, in First School on Supergravity (1982) arXiv:1201.0386]

CVE AND CME – NEW ANOMALOUS TRANSPORT


- **CME** and **CVE** are related to the gauge chiral **anomaly**:

[V. I. Zakharov, Lect. Notes Phys.871,295(2013), 1210.2186]

[D.T. Son, P. Surowka, PRL, 103 (2009) 191601]

$$\text{CME: } j^\mu = C \mu_5 B^\mu$$

$$\text{CVE: } j_A^\mu = C \mu^2 \omega^\mu$$



Quantum chiral anomaly


$$\langle \partial_\mu \hat{j}_A^\mu \rangle = -\frac{C}{8} \varepsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$$

- Interplay of **infrared** (hydrodynamics) and **ultraviolet** (anomalies) physics.

What about the **gravitational chiral anomaly**?

- The gravitational chiral anomaly grows **rapidly** with **spin**:

$$j_\mu^A = \lambda_1 (\omega_\nu \omega^\nu) \omega_\mu + \lambda_2 (a_\nu a^\nu) \omega_\mu$$



$$\langle \nabla_\mu \hat{j}_A^\mu \rangle_S = \frac{(S - 2S^3)}{96\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

New phenomenon: **kinematical vortical effect (KVE)**

GRAVITY IN FLAT SPACETIME: CHESHIRE CAT GRIN

“Well! I've often seen a cat without a grin,' thought Alice 'but a grin without a cat! It's the most curious thing i ever saw in my life!”

— Lewis Carroll, Alice in Wonderland



GRAVITY IN FLAT SPACETIME: CHESHIRE CAT GRIN

“Well! I've often seen a cat without a grin,' thought Alice 'but a grin without a cat! It's the most curious thing i ever saw in my life!”

— Lewis Carroll, Alice in Wonderland



PART 2

**GRAVITATIONAL
CHIRAL ANOMALY AND
CUBIC GRADIENTS**

HYDRODYNAMICS IN CURVED SPACE-TIME

Consider an uncharged fluid of particles with an arbitrary spin in a **gravitational field**:

fluid

4-velocity of the fluid	$u_\mu(x)$
Proper temperature	$T(x)$
Inverse temperature vector	$\beta_\mu = u_\mu/T$
Thermal vorticity tensor (analogous to the acceleration tensor)	$\varpi_{\mu\nu} = -\frac{1}{2}(\nabla_\mu\beta_\nu - \nabla_\nu\beta_\mu)$

space-time

Curved space-time metric
$g_{\mu\nu}(x)$
Riemann tensor
$R_{\mu\nu\kappa\lambda}$

We consider a medium in a state of **(global) thermodynamic equilibrium**

[F. Becattini, L. Bucciardini, E. Grossi, L. Tinti,
Eur. Phys. J. C 75, 191 (2015)]

[F. Becattini, Acta Phys. Polon. B 47, 1819 (2016)]

For example, we can find the second derivative:

Killing equation

$$\nabla_\mu\beta_\nu + \nabla_\nu\beta_\mu = 0$$

i.e. *stationary* spacetime is considered

$$\nabla_\mu\nabla_\nu\beta_\alpha = R^\rho{}_{\mu\nu\alpha}\beta_\rho$$

DECOMPOSITION OF THE TENSORS

Components of the thermal vorticity tensor

6 components

[M. Buzzegoli, E. Grossi, F. Becattini, JHEP 10 (2017) 091]

$$\varpi_{\mu\nu} = \epsilon_{\mu\nu\alpha\beta} \omega^\alpha u^\beta + \alpha_\mu u_\nu - \alpha_\nu u_\mu$$

Similar to the expansion for the electromagnetic field

We also decompose the Riemann tensor into components

$$\begin{aligned} R_{\mu\nu\alpha\beta} = & u_\mu u_\alpha A_{\nu\beta} + u_\nu u_\beta A_{\mu\alpha} - u_\nu u_\alpha A_{\mu\beta} - u_\mu u_\beta A_{\nu\alpha} \\ & + \epsilon_{\mu\nu\lambda\rho} u^\rho (u_\alpha B^\lambda_\beta - u_\beta B^\lambda_\alpha) \\ & + \epsilon_{\alpha\beta\lambda\rho} u^\rho (u_\mu B^\lambda_\nu - u_\nu B^\lambda_\mu) \\ & + \epsilon_{\mu\nu\lambda\rho} \epsilon_{\alpha\beta\eta\sigma} u^\rho u^\sigma C^{\lambda\eta} \end{aligned}$$

20 components

Generalization of 3d tensors from

Coincide with 3d tensors in the fluid rest frame.

[L. D. Landau and E. M. Lifschits,
The Classical Theory of Fields, Vol. 2, 1975]

GRADIENT EXPANSION IN THE CURVED SPACETIME

The gravitational chiral anomaly has the **4th order** in gradients - it is to be related to the **3rd order** terms in gradient expansion of the axial current.

Hydrodynamic expansion of the axial current up to the **3rd order** in gradients:

$$j_{\mu}^{A(3)} = \xi_1(T)w^2w_{\mu} + \xi_2(T)\alpha^2w_{\mu} + \xi_3(T)(\alpha w)w_{\mu} \\ + \xi_4(T)A_{\mu\nu}w^{\nu} + \xi_5(T)B_{\mu\nu}\alpha^{\nu}$$

Diagram annotations:

- A grey arrow points from the text "arbitrary coefficients" to the coefficients ξ_1, ξ_2, ξ_3 .
- Red arrows point from the text "Survive in flat spacetime" to the terms $\xi_2(T)\alpha^2w_{\mu}$ and $\xi_3(T)(\alpha w)w_{\mu}$.
- A blue arrow points from the text "'gravitational' currents" to the terms $\xi_4(T)A_{\mu\nu}w^{\nu}$ and $\xi_5(T)B_{\mu\nu}\alpha^{\nu}$.

We use only:

$$\nabla_{\mu}j_{A}^{\mu} = \mathcal{N}\epsilon^{\mu\nu\alpha\beta}R_{\mu\nu\lambda\rho}R_{\alpha\beta}^{\lambda\rho}$$

ANOMALY MATCHING: SYSTEM OF EQUATIONS

The divergence of the axial current transforms into the sum of **independent** terms:

$$\begin{aligned}\nabla_{\mu} j_{A(3)}^{\mu} &= (\alpha w) w^2 (-3T\xi_1 + T^2\xi'_1 + 2T\xi_3) \\ &+ (\alpha w) \alpha^2 (-3T\xi_2 + T^2\xi'_2 - T\xi_3 + T^2\xi'_3) \\ &+ A_{\mu\nu} \alpha^{\mu} w^{\nu} (T^2\xi'_4 + 3T\xi_5 + 2T^{-1}\xi_2 + T^{-1}\xi_3) \\ &+ B_{\mu\nu} w^{\mu} w^{\nu} (-2T^{-1}\xi_1 - 3T\xi_4 - T\xi_5) \\ &+ B_{\mu\nu} \alpha^{\mu} \alpha^{\nu} (T^2\xi'_5 - T\xi_5 - T^{-1}\xi_3) \\ &+ A_{\mu\nu} B^{\mu\nu} (-T^{-1}\xi_4 + T^{-1}\xi_5) \\ &= 32\mathcal{N} A_{\mu\nu} B^{\mu\nu} .\end{aligned}$$

Principle:

In this case, the divergence is equal to the anomaly: additional terms - *macroscopic* - cannot violate the equation from the fundamental *microscopic* theory.

The coefficient in front of each pseudoscalar must be equal to zero - a **system of equations** for the unknown coefficients $\xi_n(T)$.

ANOMALY MATCHING: SYSTEM OF EQUATIONS

This **system of linear differential equations** has the form:

$$\begin{aligned} -3T\xi_1 + T^2\xi'_1 + 2T\xi_3 &= 0 \\ -3T\xi_2 + T^2\xi'_2 - T\xi_3 + T^2\xi'_3 &= 0 \\ T^2\xi'_4 + 3T\xi_5 + 2T^{-1}\xi_2 + T^{-1}\xi_3 &= 0 \\ -2T^{-1}\xi_1 - 3T\xi_4 - T\xi_5 &= 0 \\ T^2\xi'_5 - T\xi_5 - T^{-1}\xi_3 &= 0 \\ -T^{-1}\xi_4 + T^{-1}\xi_5 - 32\mathcal{N} &= 0 \end{aligned}$$

The factor from the **gravitational chiral anomaly**

ANOMALY MATCHING: SOLUTION

Since the theory does not include **dimensional** parameters other than **temperature**:

$$\xi_1 = T^3 \lambda_1 \quad \xi_2 = T^3 \lambda_2 \quad \xi_3 = T^3 \lambda_3 \quad \xi_4 = T \lambda_4 \quad \xi_5 = T \lambda_5$$

The current: $j_\mu^{A(3)} = \lambda_1 \omega^2 \omega_\mu + \lambda_2 a^2 \omega_\mu + \lambda_3 (a\omega) \omega_\mu + \lambda_4 A_{\mu\nu} \omega^\nu + \lambda_5 B_{\mu\nu} a^\nu$

The solution looks like:

$$\begin{aligned} \frac{\lambda_1 - \lambda_2}{32} &= \mathcal{N} & \lambda_4 &= -8\mathcal{N} - \frac{\lambda_1}{2} \\ \lambda_3 &= 0 & \lambda_5 &= 24\mathcal{N} - \frac{\lambda_1}{2} \end{aligned}$$

was also shown in

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, JHEP 02 (2019)]

FLAT SPACE LIMIT:

KINEMATICAL VORTICAL EFFECT (KVE)

Let's move on to the limit of **flat space-time**. Despite the absence of a gravitational field, there **remains a contribution** to the axial current induced by the gravitational chiral anomaly:

$$\left\{ \begin{array}{l} j_{\mu}^A = \lambda_1 (\omega_{\nu} \omega^{\nu}) \omega_{\mu} + \lambda_2 (a_{\nu} a^{\nu}) \omega_{\mu} \quad \leftarrow R_{\mu\nu\alpha\beta} = 0 \\ \frac{\lambda_1 - \lambda_2}{32} = \mathcal{N} \quad \leftarrow \nabla_{\mu} j_A^{\mu} = \mathcal{N} \epsilon^{\mu\nu\alpha\beta} R_{\mu\nu\lambda\rho} R_{\alpha\beta}{}^{\lambda\rho} \end{array} \right.$$

- A new type of anomalous transport - the **Kinematical Vortical Effect (KVE)**.
- Does not explicitly depend on temperature and density \rightarrow determined only by the **kinematics** of the flow.

PART 3

**VERIFICATION:
SPIN 1/2**

TRANSPORT COEFFICIENTS AND ANOMALY:

SPIN 1/2

- In [GP, O.V. Teryaev, and V.I. Zakharov, JHEP, 02:146, 2019], [V. E. Ambrus, JHEP, 08:016, 2020], [A. Palermo, et al. JHEP 10 (2021) 077] and for ω^3 in [A. Vilenkin, Phys. Rev., D20:1807-1812, 1979] the following expression was obtained:


$$j_{\mu}^A = \left(\frac{T^2}{6} + \frac{\mu^2}{2\pi^2} - \overbrace{\frac{\omega^2}{24\pi^2} - \frac{a^2}{8\pi^2}}^{\text{KVE}} \right) \omega_{\mu}$$

- Comparing it with the well-known anomaly [L. Alvarez-Gaume, E. Witten, Nucl. Phys., B234:269, 1984]:

$$\nabla_{\mu} j_A^{\mu} = \frac{1}{384\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\alpha\beta} R_{\mu\nu\lambda\rho} R_{\alpha\beta}{}^{\lambda\rho}$$

We see that the formula is **fulfilled**:

$$\frac{\lambda_1 - \lambda_2}{32} = \mathcal{N}$$



$$\left(-\frac{1}{24\pi^2} + \frac{1}{8\pi^2} \right) / 32 = \frac{1}{384\pi^2}$$

Correspondence between **gravity** and **hydrodynamics** is shown!

PART 4

VERIFICATION:

SPIN 3/2

RARITA-SCHWINGER-ADLER MODEL OF SPIN 3/2

The **Rarita-Schwinger theory** - well-known theory of spin 3/2.
But this theory has a number of **pathologies**.

Generalized Hamiltonian dynamics: **Dirac bracket** instead of **Poisson bracket**

There is **singularity** in a Dirac bracket in weak gauge field limit for RS-theory!

Doesn't allow to construct perturbation theory!

Solved in [\[Stephen L. Adler. Phys. Rev. D, 97\(4\):045014, 2018\]](#) by introducing of interaction with additional spin 1/2 field:

$$S = \int d^4x \left(-\varepsilon^{\lambda\rho\mu\nu} \bar{\psi}_\lambda \gamma_5 \gamma_\mu \partial_\nu \psi_\rho + i\bar{\lambda} \gamma^\mu \partial_\mu \lambda - im\bar{\lambda} \gamma^\mu \psi_\mu + im\bar{\psi}_\mu \gamma^\mu \lambda \right)$$

GRAVITATIONAL CHIRAL ANOMALY IN RSA THEORY

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov,
Phys. Rev. D 106 (2) (2022) 025022]

Summing 9 correlates (contributions of different fields), we will obtain:

Matches the form we want!

$$\langle T \hat{T}^{\mu\nu}(x) \hat{T}^{\sigma\rho}(y) \hat{j}_A^\omega(z) \rangle_c = -19 \left(4\pi^6 (x-y)^5 \times (x-z)^3 (y-z)^3 \right)^{-1} e_\vartheta \left(\eta^{\nu\rho} \varepsilon^{\sigma\vartheta\mu\omega} + \eta^{\mu\rho} \varepsilon^{\sigma\vartheta\nu\omega} + \eta^{\nu\sigma} \varepsilon^{\vartheta\mu\rho\omega} + \eta^{\mu\sigma} \varepsilon^{\vartheta\nu\rho\omega} - 6e^2 (e^\nu e^\rho \varepsilon^{\sigma\vartheta\mu\omega} + e^\mu e^\rho \varepsilon^{\sigma\vartheta\nu\omega} + e^\sigma e^\nu \varepsilon^{\vartheta\mu\rho\omega} + e^\sigma e^\mu \varepsilon^{\vartheta\nu\rho\omega}) \right)$$



$$\langle T \hat{T}^{\mu\nu}(x) \hat{T}^{\sigma\rho}(y) \hat{j}_A^\omega(z) \rangle = \mathcal{A} \left(4(x-y)^5 \times (x-z)^3 (y-z)^3 \right)^{-1} e_\vartheta \left(\eta^{\nu\rho} \varepsilon^{\sigma\vartheta\mu\omega} + \eta^{\mu\rho} \varepsilon^{\sigma\vartheta\nu\omega} + \eta^{\nu\sigma} \varepsilon^{\vartheta\mu\rho\omega} + \eta^{\mu\sigma} \varepsilon^{\vartheta\nu\rho\omega} - 6e^2 (e^\nu e^\rho \varepsilon^{\sigma\vartheta\mu\omega} + e^\mu e^\rho \varepsilon^{\sigma\vartheta\nu\omega} + e^\sigma e^\nu \varepsilon^{\vartheta\mu\rho\omega} + e^\sigma e^\mu \varepsilon^{\vartheta\nu\rho\omega}) \right)$$

(points on the same axis)

We can determine the factor in the anomaly:

$$\mathcal{A}_{RSA} = -19 \mathcal{A}_{s=1/2} = -\frac{19}{\pi^6}$$

$$\langle \nabla_\mu \hat{j}_A^\mu \rangle_{RSA} = \frac{-19}{384\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

-19 times different from the anomaly for spin 1/2

ZUBAREV DENSITY OPERATOR

Global Equilibrium Conditions

$$\nabla_{\mu}\beta_{\nu} + \nabla_{\nu}\beta_{\mu} = 0 \quad \nabla_{\mu}\zeta = 0$$



Form of the **density operator** for a medium with **rotation** and **acceleration**:

$$\hat{\rho} = \frac{1}{Z} \exp \left[-b_{\mu} \hat{P}^{\mu} + \frac{1}{2} \varpi_{\mu\nu} \hat{J}^{\mu\nu} + \zeta \hat{Q} \right]$$

$$\varpi_{\mu\nu} \hat{J}^{\mu\nu} = -2\alpha^{\rho} \hat{K}_{\rho} - 2\omega^{\rho} \hat{J}_{\rho}$$

\hat{K}^{μ} - boost (related to acceleration)

\hat{J}^{μ} - angular momentum (related to vorticity)

**Lorentz Transform
Generators**

$$\hat{J}^{\mu\nu} = \int_{\Sigma} d\Sigma_{\lambda} \left(x^{\mu} \hat{T}^{\lambda\nu} - x^{\nu} \hat{T}^{\lambda\mu} \right)$$

ZUBAREV DENSITY OPERATOR

Quantum statistical mean value

$$\langle \hat{O}(x) \rangle = \frac{1}{Z} \text{tr}(\hat{\rho} \hat{O}(x))_{\text{ren}}$$

statistical sum:
cancellation of
disconnected
correlators

Perturbation theory in the third order

$$\begin{aligned} \langle \hat{O}(x) \rangle &= \langle \hat{O}(x) \rangle_{\beta(x)} + \frac{\varpi_{\mu\nu}}{2|\beta|} \int_0^{|\beta|} d\tau \langle T_\tau J_{-i\tau u}^{\mu\nu} \hat{O}(0) \rangle_{\beta(x),c} \\ &+ \frac{\varpi_{\mu\nu} \varpi_{\rho\sigma}}{8|\beta|^2} \int_0^{|\beta|} d\tau_x d\tau_y \langle T_\tau J_{-i\tau_x u}^{\mu\nu} J_{-i\tau_y u}^{\rho\sigma} \hat{O}(0) \rangle_{\beta(x),c} \\ &+ \frac{\varpi_{\mu\nu} \varpi_{\rho\sigma} \varpi_{\alpha\beta}}{48|\beta|^3} \int_0^{|\beta|} d\tau_x d\tau_y d\tau_z \langle T_\tau J_{-i\tau_x u}^{\mu\nu} J_{-i\tau_y u}^{\rho\sigma} J_{-i\tau_z u}^{\alpha\beta} \hat{O}(0) \rangle_{\beta(x),c} + \dots \end{aligned}$$

Imaginary time
 $\tau = i t$
ordering

Connected correlators

$$\langle \hat{J} \hat{O} \rangle_c = \langle \hat{J} \hat{O} \rangle - \langle \hat{J} \rangle \langle \hat{O} \rangle$$

KVE IN RSA THEORY: CALCULATION

- Finally, we obtain, in particular, for $C^{02|02|02|3|111}$

$$C^{02|02|02|3|111} = \frac{T}{480\pi^2} \int \frac{dp p e^{p/T}}{(1 + e^{p/T})^5} \left\{ 126 - 291 \frac{p}{T} - 472 \frac{p^2}{T^2} + \left[126 + 873 \frac{p}{T} + 5192 \frac{p^2}{T^2} \right] e^{p/T} \right. \\ \left. + \left[-126 + 873 \frac{p}{T} - 5192 \frac{p^2}{T^2} \right] e^{2p/T} + \left[-126 - 291 \frac{p}{T} + 472 \frac{p^2}{T^2} \right] e^{3p/T} \right\} = \frac{177T^3}{80\pi^2}$$

Calculating all the diagrams we obtain:

$$\lambda_1 = -\frac{1}{6} \left(2 \cdot \frac{177}{80\pi^2} + 6 \cdot \frac{353}{240\pi^2} \right) = -\frac{53}{24\pi^2},$$

$$\lambda_2 = -\frac{1}{6} \left(\frac{33}{40\pi^2} + \frac{53}{80\pi^2} + \frac{1}{2\pi^2} + \frac{3}{4\pi^2} + \frac{47}{80\pi^2} + \frac{17}{40\pi^2} \right) = -\frac{5}{8\pi^2}$$


Thus, the KVE in the RSA theory has the form:

$$j_{A,KVE}^\mu = \left(-\frac{53}{24\pi^2} \omega^2 - \frac{5}{8\pi^2} a^2 \right) \omega^\mu$$

KVE VS GRAVITATIONAL ANOMALY

The obtained formula for **cubic gradients** (KVE):

$$j_{\mu}^{A(3)} = \left(-\frac{53}{24\pi^2}\omega^2 - \frac{5}{8\pi^2}a^2 \right) \omega_{\mu}$$



Gravitational chiral **anomaly**:

$$\nabla_{\mu} j_A^{\mu} = \frac{-19}{384\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\alpha\beta} R_{\mu\nu\lambda\rho} R_{\alpha\beta}{}^{\lambda\rho}$$

$$\frac{\lambda_1 - \lambda_2}{32} = \mathcal{N}$$



Direct **verification**:

$$\left(-\frac{53}{24\pi^2} + \frac{5}{8\pi^2} \right) / 32 = -\frac{19}{384\pi^2}$$

Coincidence of hydrodynamics and gravitational anomaly!

- For the RSA theory, the relationship between the transport coefficients in a **vortical accelerated fluid** and the **gravitational** chiral **anomaly** is shown: the factor **-19** from the anomaly is reproduced.
- Verification of the obtained formula in a very **nontrivial** case with higher spins and interaction.

PART 7

CONCLUSION

CONCLUSION

- The relationship between the hydrodynamic current in the third order of gradient expansion $\lambda_1(\omega_\nu\omega^\nu)\omega_\mu$ and $\lambda_2(a_\nu a^\nu)\omega_\mu$, the *Kinematical Vortical Effect (KVE)*, and the **gravitational chiral anomaly** has been proven:
 - The axial current in a flat space-time in a vortical and accelerated fluid turns out to be associated with a quantum violation of current conservation in a curved space-time.
- The obtained formula has been **verified** directly for **spin 1/2**.
- The obtained formula has been **verified** for **spin 3/2** using the **RSA** theory:
 - Cubic transport coefficients were derived using the statistical density operator expansion .
 - Correspondence between the KVE and the gravitational chiral anomaly is directly shown.

The background features a white area on the left with several overlapping, semi-transparent white triangles of varying sizes and orientations. On the right, a large black area is separated from the white by a jagged, diagonal boundary. The text is centered within the black area.

**ADDITIONAL
SLIDES**

EXPERIMENT: FEW WORDS

- Is it possible to observe KVE in experiment?
- Is it possible to observe a **gravitational chiral anomaly** in the hydrodynamics of the matter, produced in heavy ion collisions?
- To observe cubic terms, it is necessary that the gradients (acceleration, vorticity, magnetic field...) give a contribution at least of the order of temperature:

$$\omega, a \sim (0.1 - 2)T$$

[A. Zinchenko, A. Sorin, O. Teryaev, M. Baznat, J.Phys.Conf.Ser. 1435 (2020)]

[F. Becattini, M. Buzzegoli, G. Inghirami, I. Karpenko, A. Palermo, Phys.Rev.Lett. 127 (2021) 27, 272302]

However, the cubic terms are suppressed by the numerical factor

$$\text{KVE: } j_{A,S=1/2}^{\mu} = \left(-\frac{1}{24\pi^2}\omega^2 - \frac{1}{8\pi^2}a^2 \right) \omega^{\mu}$$

The good news: for spin 3/2 it is enhanced by cubic growth with spin:

The bad news: should be suppressed by mass $e^{-m/T}$ (omega baryon is heavy).

Idea: consider massless **quasiparticles** with spin 3/2 in semimetals?


[I. Boettcher, Phys. Rev. Lett. 124, 127602 (2020)]

DISCUSSION

- Arbitrary fields with arbitrary spin were considered:
General exact result
- Only **conservation law** for the current was used.
- Although the effect is associated with an anomaly - it exists in **flat space-time** (the *Cheshire cat grin*).
- In contrast to **CVE** and the **gauge anomaly case**, the factor from the gravitational anomaly is split into **two conductivities**:

$$\lambda_1 - \lambda_2 = 32\mathcal{N}$$

WHAT WILL BE DISCUSSED?



GRAVITATIONAL ANOMALY IN THERMAL CVE

The answer was obtained in different approaches:

- From the holography. [\[K. Landsteiner, E. Megias, F. Pena-Benitez, "Gravitational Anomaly and Transport," Phys. Rev. Lett. 107, 021601 \(2011\)\]](#)
- In [\[S.P. Robinson, F. Wilczek. Phys. Rev. Lett., 95:011303, 2005\]](#) Hawking radiation is associated with a gravitational anomaly: it is necessary to integrate the anomaly from the horizon to infinity + the condition for the conservation of the currents on the horizon.

In [\[M. Stone, J. Kim. Phys. Rev., D98\(2\):025012, 2018\]](#) the derivation was generalized to 3+1 dimensional gravitational chiral anomaly and (analogue) of rotating black hole.

- From the condition of the translational invariance of the Euclidean vacuum. [\[K. Jensen, R. Loganayagam, A. Yarom, JHEP 02, 088 \(2013\)\]](#)

$$j_A^\nu = (\sigma_T T^2 + \sigma_\mu \mu^2) \omega^\nu$$

$$\epsilon^{\mu\nu\alpha\beta} = \frac{1}{\sqrt{-g}} \varepsilon^{\mu\nu\alpha\beta}$$

$$\sigma_T = 64\pi^2 \mathcal{N}$$



$$\nabla_\mu j_A^\mu = \mathcal{N} \epsilon^{\mu\nu\alpha\beta} R_{\mu\nu\lambda\rho} R_{\alpha\beta}{}^{\lambda\rho}$$

GRAVITATIONAL ANOMALY IN THERMAL CVE

Verified for the Dirac field

$$j_A^\mu = \left(\frac{T^2}{6} + \frac{\mu^2}{2\pi^2} \right) \omega^\mu$$

[L. Alvarez-Gaume, E. Witten, Nucl. Phys. B234 (1984) 269]

$$\nabla_\mu j_A^\mu = \frac{1}{384\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

$$\sigma_T = 64\pi^2 N$$

A **problem** arose for fields with spin 3/2 (within the framework of the Rarita-Schwinger-Adler theory) [S. L. Adler, Phys. Rev. D 97 (4) (2018) 045014]

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, Phys. Rev. D 105 (4) (2022) L041701]

$$j_A^\nu = \left(\frac{5}{6} T^2 + \frac{5}{2\pi^2} \mu^2 \right) \omega^\nu$$

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, Phys. Rev. D 106 (2) (2022) 025022]

$$\nabla_\mu j_A^\mu = -\frac{19}{384\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

$$\sigma_T \neq 64\pi^2 N$$

In hydrodynamics, the **cubic** dependence on the spin from the gravitational anomaly is **not visible**?

ANOMALY MATCHING: PRINCIPLE

Following [\[D.T. Son, P. Surowka, PRL, 103 \(2009\) 191601\]](#)

- it is necessary to construct the **entropy current**.

In [\[M. Buzzegoli, Lect. Notes Phys. 987, 53-93 \(2021\)\]](#)

[\[Shi-Zheng Yang, Jian-Hua Gao, and Zuo-Tang Liang, Symmetry 14, 948 \(2022\)\]](#)

it is shown that it is possible to use the global equilibrium condition

$$\nabla_{\mu}\beta_{\nu} + \nabla_{\nu}\beta_{\mu} = 0$$

And it is enough to consider **only** the equation for the current

Simplifies analysis and allows viewing effects in **curved space**

We use only:

$$\nabla_{\mu}j_A^{\mu} = \mathcal{N}\epsilon^{\mu\nu\alpha\beta}R_{\mu\nu\lambda\rho}R_{\alpha\beta}^{\lambda\rho}$$

We substitute the gradient expansion:

$$\nabla^{\mu}\left(\xi_1(T)w^2w_{\mu} + \xi_2(T)\alpha^2w_{\mu} + \xi_3(T)(\alpha w)w_{\mu} + \xi_4(T)A_{\mu\nu}w^{\nu} + \xi_5(T)B_{\mu\nu}\alpha^{\nu}\right) = 32\mathcal{N}A_{\mu\nu}B^{\mu\nu}$$

DERIVATIVES

Using the condition of global equilibrium and relations for the gravitational field (the Bianchi identity, etc.), we obtain for the derivatives:

Luttinger relation

[J. M. Luttinger, Phys. Rev. 135, A1505-A1514 (1964)]

$$\nabla_{\mu} T = T^2 \alpha_{\mu},$$

$$\nabla_{\mu} u_{\nu} = T(\epsilon_{\mu\nu\alpha\beta} u^{\alpha} w^{\beta} + u_{\mu} \alpha_{\nu}),$$

$$\nabla_{\mu} w_{\nu} = T(-g_{\mu\nu}(w\alpha) + \alpha_{\mu} w_{\nu}) - T^{-1} B_{\nu\mu},$$

$$\begin{aligned} \nabla_{\mu} \alpha_{\nu} = T & (w^2(g_{\mu\nu} - u_{\mu} u_{\nu}) - \alpha^2 u_{\mu} u_{\nu} - w_{\mu} w_{\nu} \\ & - u_{\mu} \eta_{\nu} - u_{\nu} \eta_{\mu}) + T^{-1} A_{\mu\nu}, \end{aligned}$$

$$\nabla^{\mu}(A_{\mu\nu} w^{\nu}) = -3TB_{\mu\nu} w^{\mu} w^{\nu} - T^{-1} A_{\mu\nu} B^{\mu\nu},$$

$$\begin{aligned} \nabla^{\mu}(B_{\mu\nu} \alpha^{\nu}) = 3TA_{\mu\nu} w^{\mu} \alpha^{\nu} + T^{-1} A_{\mu\nu} B^{\mu\nu} \\ - TB_{\mu\nu} w^{\mu} w^{\nu} - TB_{\mu\nu} \alpha^{\mu} \alpha^{\nu} \end{aligned}$$

DISCUSSION

It also turns out that in the current

$$j_{\mu}^{A(3)} = \lambda_1 \omega^2 \omega_{\mu} + \lambda_2 a^2 \omega_{\mu} + \lambda_3 (a\omega) \omega_{\mu} + \lambda_4 A_{\mu\nu} \omega^{\nu} + \lambda_5 B_{\mu\nu} a^{\nu}$$

- **Difference of flat-space terms** is to be equal to **difference** of **curved-space terms**:

$$\lambda_1 - \lambda_2 = \lambda_5 - \lambda_4$$

- At the finite mass, λ begin to depend on mass and temperature:

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, JHEP 02 (2019)]

ZUBAREV DENSITY OPERATOR

- The density operator $\hat{\rho}$ plays a central role in the statistical description of a medium.
- Integration in the imaginary time plane: Quantum field theory in imaginary time.
- General covariant form of the density operator in local equilibrium.

$$\hat{\rho} = \frac{1}{Z} \exp \left[- \int_{\Sigma} d\Sigma_{\mu} \left(\hat{T}^{\mu\nu}(x) \beta_{\nu}(x) - \zeta(x) \hat{j}^{\mu}(x) \right) \right]$$

maximal

$$S = -\text{tr}(\hat{\rho} \log \hat{\rho})$$

[D.N. Zubarev, A.V. Prozorkevich and S.A. Smolyanskii, *Theor. Math. Phys.* **40** (1979) 821.]

[M. Buzzegoli, E. Grossi and F. Becattini, *JHEP* **1710** (2017).]



On condition

$$\nabla_{\mu} \beta_{\nu} + \nabla_{\nu} \beta_{\mu} = 0 \quad \nabla_{\mu} \zeta = 0$$



$$\begin{aligned} \beta_{\mu}(x) &= b_{\mu} + \varpi_{\mu\nu} x^{\nu} & \zeta &= \text{const} \\ b_{\mu} &= \text{const} & \varpi_{\mu\nu} &= \text{const} \end{aligned}$$



$\hat{\rho}$ does not depend on the choice of hypersurface $d\Sigma_{\mu}$



global thermodynamic equilibrium

RARITA-SCHWINGER-ADLER MODEL OF SPIN 3/2

- The interaction **shifts the pole** in the Dirac bracket!

$$[\Psi_i(\vec{x}), \Psi_j^\dagger(\vec{y})]_D = -i \left[(\delta_{ij} - \frac{1}{2} \sigma_i \sigma_j) \delta^3(\vec{x} - \vec{y}) - \vec{D}_{\vec{x}i} \frac{\delta^3(\vec{x} - \vec{y})}{m^2 + g\vec{\sigma} \cdot \vec{B}(\vec{x})} \overleftarrow{D}_{\vec{y}j} \right]$$

Contribution of interaction with an additional field 

- The **stress-energy tensor** can be obtained by varying with respect to the metric

$$T^{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta S}{\delta g_{\mu\nu}}$$

$$T^{\mu\nu} = \frac{1}{2} \varepsilon^{\lambda\nu\beta\rho} \bar{\psi}_\lambda \gamma_5 \gamma^\mu \partial_\beta \psi_\rho + \frac{1}{8} \partial_\eta \left(\varepsilon^{\lambda\alpha\nu\rho} \bar{\psi}_\lambda \gamma_5 \gamma_\alpha [\gamma^\eta, \gamma^\mu] \psi_\rho \right) + \frac{i}{4} \left(\bar{\lambda} \gamma^\nu \partial^\mu \lambda - \partial^\mu \bar{\lambda} \gamma^\nu \lambda \right) + \frac{i}{2} m \left(\bar{\psi}^\mu \gamma^\nu \lambda - \bar{\lambda} \gamma^\nu \psi^\mu \right) + (\mu \leftrightarrow \nu).$$

Traceless unlike the usual Rarita-Schwinger field $T^\mu_\mu = 0$

- The currents can be obtained from Noether's theorem. The **axial current** can be constructed for the $U(1)_A$ transformation:

$$j_A^\mu = -i \varepsilon^{\lambda\rho\nu\mu} \bar{\psi}_\lambda \gamma_\nu \psi_\rho + \bar{\lambda} \gamma_\mu \gamma_5 \lambda$$

KVE AND UNRUH EFFECT

- It is possible to distinguish **conserved** and **anomalous** parts of the current:

$$j_{\mu}^A = j_{\mu(\text{conserv})}^A + j_{\mu(\text{anom})}^A$$

$$\text{Thermal vorticity tensor squared } \omega^2 - a^2 = -\frac{1}{2}\varpi_{\mu\nu}\varpi^{\mu\nu}$$

$$j_{\mu(\text{anom})}^A = \left. \begin{matrix} -A_{\mu\nu}\omega^{\nu} \\ +B_{\mu\nu}a^{\nu} \end{matrix} \right\} \mathcal{N} \left\{ (\omega^2 - a^2)\omega_{\mu} - A_{\mu\nu}\omega^{\nu} + B_{\mu\nu}a^{\nu} \right\} \Rightarrow \nabla^{\mu} j_{\mu(\text{anom})}^A = \mathcal{N} \epsilon^{\mu\nu\alpha\beta} R_{\mu\nu\lambda\rho} R_{\alpha\beta}{}^{\lambda\rho}$$

$$j_{\mu(\text{conserv})}^A = \frac{\lambda_1 + \lambda_2}{2} \left\{ (\omega^2 + a^2)\omega_{\mu} - \frac{1}{2}A_{\mu\nu}\omega^{\nu} - \frac{1}{2}B_{\mu\nu}a^{\nu} \right\} \Rightarrow \nabla^{\mu} j_{\mu(\text{conserv})}^A = 0$$

- Consider the term with acceleration from the anomalous part of the current:

$$j_{\mu(\text{anom})}^A = -16\mathcal{N}a^2\omega_{\mu}$$

- Unruh effect** [W.G. Unruh, 1976] - in an accelerated frame there is a thermal bath of particles with the **Unruh temperature**:

$$T_U = |a|/(2\pi)$$

Substitute $|a| \rightarrow 2\pi T_U$:

$$j_{\mu(\text{anom})}^A = \frac{T_U^2}{6}\omega_{\mu} \quad \text{for spin } 1/2 \rightarrow \text{standard CVE}$$

$$j_{\mu(\text{anom})}^A = 64\pi^2 \mathcal{N} T_U^2 \omega_{\mu}$$

thermal CVE current is **proportional** to the **anomaly**!

- Match with [K. Landsteiner, et al. PRL, 2011] and [M. Stone, J. Kim. PRD, 2018], where **thermal CVE** $\mathbf{j}_A \sim T^2 \boldsymbol{\Omega}$ is associated with the **gravitational chiral anomaly**!

CHIRAL ANOMALY IN RSA THEORY: GAUGE PART

- Since the problem with the Dirac bracket is solved – **perturbation theory** can be constructed
- The **chiral (gauge) quantum anomaly** was obtained by the shift method:

[Stephen L. Adler, Phys. Rev. D, 97(4):045014, 2018]

see also

[S. L. Adler, P. Pais, Phys. Rev. D 99, 095037 (2019)]

$$\langle \partial_\mu \hat{j}_A^\mu \rangle = -\frac{5}{16\pi^2} \varepsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$$

Also by the method of conformal three-point functions:

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, Phys. Rev. D 106 (2) (2022) 025022]

- The factor “**5**” differs from what is expected according to the prediction “**3**” based on supergravity

[M. J. Duff, in First School on Supergravity (1982) arXiv:1201.0386]

However the correspondence is restored if we take into account that there are two additional degrees of freedom with spin $\frac{1}{2}$:
then **5=3+2**

GRAVITATIONAL CHIRAL ANOMALY IN RSA THEORY

For a conformally symmetric theory, if

$$\partial_\mu T^{\mu\nu} = 0, \quad \partial_\mu j_V^\mu = 0, \quad \partial_\mu j_A^\mu = 0, \\ T_\mu^\mu = 0, \quad T_{\mu\nu} = T_{\nu\mu}.$$

It is proven in [J. Erdmenger, Nucl. Phys. B, 562:315–329, 1999], that the three-point function $\langle T\hat{T}_{\mu\nu}(x)\hat{T}_{\sigma\rho}(y)\hat{j}_\omega^A(z)\rangle_c$ has the **universal form**:

$$\langle T\hat{T}^{\mu\nu}(x)\hat{T}^{\sigma\rho}(y)\hat{j}_A^\omega(z)\rangle_c = \frac{1}{(x-z)^8(y-z)^8} \\ \times \mathcal{I}_T^{\mu\nu,\mu'\nu'}(x-z)\mathcal{I}_T^{\sigma\rho,\sigma'\rho'}(y-z)t_{\mu'\nu'\sigma'\rho'}^{TTA\omega}(Z)$$

where the notations are introduced:

“6” - consequence of $T_\mu^\mu = 0$

$$\mathcal{I}_{\mu\nu,\sigma\rho}^T(x) = \left. \begin{matrix} \mathcal{E}_{\mu\nu,\alpha\beta}^T \\ \mathcal{E}_{\mu\nu,\alpha\beta}^T \end{matrix} \right\} (x),$$

$$t_{\mu\nu\sigma\rho\omega}^{TTA} = \frac{1}{Z^6} (\mathcal{E}_{\mu\nu,\eta}^{\sigma\perp} \mathcal{E}_{\sigma\rho,\kappa\varepsilon}^{\omega\perp} \hat{\varepsilon}_\omega^{\eta\kappa\lambda} Z_\lambda \\ - 6 \mathcal{E}_{\mu\nu,\eta\gamma}^T \mathcal{E}_{\sigma\rho,\kappa\delta}^T \varepsilon_\omega^{\eta\kappa\lambda} Z^\gamma Z^\delta Z_\lambda Z^{-2})$$

GRAVITATIONAL CHIRAL ANOMALY IN RSA THEORY

- **How to explain the factor -19?**
- How does it **relate** to **previous** calculations?

$$\langle \nabla_{\mu} \hat{j}_A^{\mu} \rangle_{RS} = \frac{-21}{384\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

[M. J. Duff, 1201.0386] [J.J.M. Carrasco, et al. JHEP, 07:029, 2013]

“ghostless” contribution [J.J.M. Carrasco, et al. JHEP, 07:029, 2013]

$$-19 = -20 + 1$$

Contribution
of spin 1/2

$$-19 = -21 + 2$$

KVE IN RSA THEORY: CALCULATION

Propagator at finite temperature

$$\langle T_\tau \Psi_{aI_1}(X) \bar{\Psi}_{bI_2}(Y) \rangle = \oint_P e^{iP(X-Y)} G_{ab(I_1 I_2)}(P)$$

contains either **2nd or 4th order poles**:

$$G(P) = G_1(P) + G_2(P), \quad G_1 \sim 1/P^2, \quad G_2 \sim 1/P^4$$

Formulas for summation over Matsubara frequencies

[M. Buzzegoli, thesis 2020. arXiv: 2004.08186]

$$\sum_{\omega_n=(2n+1)\pi T} \frac{f(\omega_n) e^{i\omega_n \tau}}{\omega_n^2 + E^2} = \frac{1}{2ET} \sum_{s=\pm 1} f(-isE) e^{\tau s E} [\theta(-s\tau) - n_F(E)]$$

$$\sum_{\omega_n=(2n+1)\pi T} \frac{f(\omega_n) e^{i\omega_n \tau}}{(\omega_n^2 + E^2)^2} = \frac{1}{T} \sum_{s=\pm 1} e^{\tau s E} \left\{ \frac{f(-isE)}{4E^2} n'_F(E) + \frac{(1 - s\tau E) f(-isE) + isE f'(-isE)}{4E^3} [\theta(-s\tau) - n_F(E)] \right\}$$

Fermi-Dirac distribution

KVE IN RSA THEORY: CALCULATION

Let us consider (for illustration) the contribution of the first of the term from the Wick's theorem and only from $G_1 \sim 1/P^2$. After **summation over the Matsubara** frequencies, we obtain:

$$C_{\text{Wick1,G1}}^{\alpha_1\alpha_2|\alpha_3\alpha_4|\alpha_5\alpha_6|\lambda|ijk} = -T^3 \int [d\tau] d^3x d^3y d^3z x^i y^j z^k \times \int \frac{d^3p_1 d^3p_2 d^3p_3 d^3p_4}{16E_1 E_2 E_3 E_4} e^{-i\mathbf{p}_1(\mathbf{y}-\mathbf{x}) - i\mathbf{p}_2\mathbf{x} - i\mathbf{p}_3(\mathbf{z}-\mathbf{y}) + i\mathbf{p}_4\mathbf{z}}$$

$$\times \sum_{s_n=\pm 1} e^{s_1 E_1(\tau_y - \tau_x) + s_2 E_2 \tau_x + s_3 E_3(\tau_z - \tau_y) - s_4 E_4 \tau_z} \left\{ \theta(-s_1[\tau_y - \tau_x]) - n_F(E_1) \right\}$$

$$\times \left\{ \theta(-s_2 \tau_x) - n_F(E_2) \right\} \left\{ \theta(-s_3[\tau_z - \tau_y]) - n_F(E_3) \right\} \left\{ \theta(-s_4 \tau_z) - n_F(E_4) \right\}$$

$$\times \text{tr}_{I,a} \left\{ \mathcal{D}^{\alpha_1\alpha_2}(-i\tilde{P}_1, i\tilde{P}_2) G_1(\tilde{P}_2) \mathcal{J}^\lambda G_1(\tilde{P}_4) \mathcal{D}^{\alpha_5\alpha_6}(-i\tilde{P}_4, i\tilde{P}_3) G_1(\tilde{P}_3) \mathcal{D}^{\alpha_3\alpha_4}(-i\tilde{P}_3, i\tilde{P}_1) G_1(\tilde{P}_1) \right\}$$

where

Contains an **explicit dependence on coordinates**.

$$\tilde{P}_\mu^n = (-is_n E_n, -\mathbf{p}_n), \quad E_n = |\mathbf{p}_n|$$

- Can be absorbed into **derivatives** by integration by parts

$$\int d^3p_1 d^3p_2 d^3p_3 d^3p_4 d^3x d^3y d^3z x^i y^j z^k e^{-i\mathbf{p}_1(\mathbf{y}-\mathbf{x}) - i\mathbf{p}_2\mathbf{x} - i\mathbf{p}_3(\mathbf{z}-\mathbf{y}) + i\mathbf{p}_4\mathbf{z}} f(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4) =$$

$$= i(2\pi)^9 \int d^3p \left(\frac{\partial^3}{\partial p_4^k \partial p_2^i \partial p_3^j} + \frac{\partial^3}{\partial p_4^k \partial p_2^i \partial p_4^j} \right) f(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}_3, \mathbf{p}_4) \Big|_{\substack{\mathbf{p}_4=\mathbf{p}_1 \\ \mathbf{p}_3=\mathbf{p}_1 \\ \mathbf{p}_2=\mathbf{p}_1}} .$$

There remains only **one** integral over the momentum.

KVE VS GRAVITATIONAL ANOMALY

- Considering also the linear terms, we obtain

$$j_{A,RSA}^{\mu} = \left(\frac{5}{6}T^2 + \frac{5}{2\pi^2}\mu^2 - \frac{53}{24\pi^2}\omega^2 - \frac{5}{8\pi^2}a^2 \right) \omega^{\mu}$$

A similar formula in the case of Dirac fields led to the expression in “all orders”:

$$j_{A,Dirac}^{\mu} = \left(\frac{T^2}{6} + \frac{\mu^2}{2\pi^2} - \frac{\omega^2}{24\pi^2} - \frac{a^2}{8\pi^2} \right) \omega^{\mu}$$

(at low temperatures $T \sim |a|, |\omega|$ instability was found)

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, *Phys. Rev. D* **100**, 125009 (2019)]

Also gives the “all-orders” formula?

- Using the remaining pair for the transport coefficient, we obtain:

$$j_{A,RSA}^{\mu} = \left(\frac{5T^2}{6} + \frac{5\mu^2}{2\pi^2} \right) \omega_{\mu} + \left(-\frac{53}{24\pi^2}\omega^2 - \frac{5}{8\pi^2}a^2 \right) \omega_{\mu} + \frac{3}{2\pi^2}A_{\mu\nu}\omega^{\nu} - \frac{1}{12\pi^2}B_{\mu\nu}a^{\nu}$$

where

$$A_{\mu\nu} = u^{\alpha}u^{\beta}R_{\alpha\mu\beta\nu}$$



$$B_{\mu\nu} = \frac{1}{2}\epsilon_{\alpha\mu\eta\rho}u^{\alpha}u^{\beta}R_{\beta\nu}{}^{\eta\rho}$$

Current (at high temperatures) in gravitational field for approximately empty space?

EXPERIMENT: FEW WORDS

In non-central collisions of heavy ions, huge magnetic fields and huge angular momentum arise. Differential rotation - different at different points: **vorticity** and **vortices**.


- Rotation 25 orders of magnitude faster, than the rotation of the earth:
- The vorticity has order 10^{22} sec^{-1}



[Phys. Rev. Lett. 117, 192301 (2016)]

EXPERIMENT: FEW WORDS

Vorticity transforms into polarization



[Nature 548 (2017) 62-65
arXiv:1701.06657 [nucl-ex]]

- Generation of **hyperon polarization**.
- Both **vorticity** and **acceleration** are essential for polarization.
- Also described based on **Chiral Vortical Effect (CVE)** [Rogachevsky, Sorin, Teryaev, Phys.Rev.C82(2010) 054910], [Baznat, Gudima, Sorin, Teryaev, Phys.Rev.C93, no.3,031902 (2016)]

$$\text{CVE: } \langle j_{\mu}^5 \rangle = \left(\frac{T^2}{6} + \frac{\mu^2}{2\pi^2} \right) \omega_{\mu}$$

- **Qualitative and quantitative correspondence!**
- Polarization from quantum anomaly \sim *spin crisis* and *gluon anomaly*: [Efremov, Soffer, Teryaev, Nucl.Phys.B 346 (1990) 97-114]

proton spin \rightarrow hyperon polarization,
gluon field \rightarrow chemical potential*4-velocity

Also described on the basis of a thermodynamic approach (Wigner function):

[I. Karpenko, F. Becattini, Nucl.Phys.A 982 (2019) 519-522]

DECOMPOSITION OF THE TENSORS: GRAVITY

Inverse formulas

$$A_{\mu\nu} = u^\alpha u^\beta R_{\alpha\mu\beta\nu}$$

Symmetric tensor

6 components

$$B_{\mu\nu} = \frac{1}{2} \epsilon_{\alpha\mu\eta\rho} u^\alpha u^\beta R_{\beta\nu}{}^{\eta\rho}$$

Nonsymmetric traceless pseudotensor dual to the Riemann tensor

8 components

$$C_{\mu\nu} = \frac{1}{4} \epsilon_{\alpha\mu\eta\rho} \epsilon_{\beta\nu\lambda\gamma} u^\alpha u^\beta R^{\eta\rho\lambda\gamma}$$

Double dual symmetric Riemann tensor

6 components

CVE AND CME – NEW ANOMALOUS TRANSPORT

However there are gradient terms that are not related to dissipation.

[V. I. Zakharov, Lect. Notes Phys.871,295(2013), 1210.2186]

Chiral magnetic effect (CME)

$$\text{CME: } j^\mu = C \mu_5 B^\mu$$

Current flows along the magnetic field

Chiral vortical effect (CVE)

$$\text{CVE: } j_A^\mu = C \mu^2 \omega^\mu$$

$$\omega^\mu = \frac{1}{2} \varepsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha u_\beta$$

Current flows along the vorticity

Consistency with quantum anomaly modifies hydrodynamic equations

[D.T. Son, P. Surowka, PRL, 103 (2009) 191601]:

- Generalization of [L. D. Landau, E. M. Lifshitz, Fluid Mechanics, Vol. 6, 1987]

Conservation equations:

$$\partial_\mu T^{\mu\nu} = F^{\nu\lambda} j_\lambda$$

+

Quantum chiral anomaly

$$\langle \partial_\mu \hat{j}_A^\mu \rangle = -\frac{C}{8} \varepsilon^{\mu\nu\alpha\beta} F_{\mu\nu} F_{\alpha\beta}$$

+

Second law of thermodynamics

$$\partial_\mu s^\mu \geq 0$$

MODERN DEVELOPMENT AND THE PROBLEM

A whole set of similar effects has been found at the intersection of quantum field theory and hydrodynamics [M. N. Chernodub et al. Phys. Rept. 977 (2022)].

Experimental search is in progress:

- CME not yet found in ion collisions at RHIC [D.E. Kharzeev et al. 2205.00120].
- Condensed matter copies of the effects are found in semimetals [Qiang Li et al. Nature Phys. 12 (2016)].

What about the **gravitational chiral anomaly**?

- The gravitational chiral anomaly grows **rapidly** with **spin**:

$$\langle \nabla_{\mu} \hat{j}_A^{\mu} \rangle_S = \frac{(S - \dots)}{96\pi^2 \sqrt{-g}} \varepsilon^{\mu\nu\rho\sigma} R_{\mu\nu\kappa\lambda} R_{\rho\sigma}{}^{\kappa\lambda}$$

[M. J. Duff, in First School on Supergravity (1982) arXiv:1201.0386]

[S. M. Christensen, M. J. Duff, Nucl. Phys. B 154, 301–342 (1979)]

How does the **gravitational** chiral anomaly manifest itself in **hydrodynamics**?

Is it possible to see the **factor** $S - 2S^3$ in hydrodynamics?

TRANSPORT COEFFICIENTS AND ANOMALY: SPIN 1/2

Using another pair of equations

$$\lambda_4 = -8\mathcal{N} - \frac{\lambda_1}{2}$$

$$\lambda_5 = 24\mathcal{N} - \frac{\lambda_1}{2}$$

we obtain a current for the Dirac field that includes “gravitational” terms (taking into account also linear terms):

$$j_\mu^{A,S=1/2} = \left(\frac{T^2}{6} + \frac{\mu^2}{2\pi^2} \right) \omega^\mu + \left(-\frac{1}{24\pi^2} \omega^2 - \frac{1}{8\pi^2} a^2 \right) \omega_\mu + \frac{1}{24\pi^2} \epsilon_{\alpha\mu\eta\rho} R^{\beta\nu\eta\rho} u^\alpha u_\beta a_\nu$$

- Exact result - shown outside of perturbation theory

(But does not work at low temperatures $T \sim |a|, |\omega|$ due to instability.

[G. Y. Prokhorov, O. V. Teryaev, V. I. Zakharov, *Phys. Rev. D* **100**, 125009 (2019)]

- In most of the cases - polynomiality in $\omega, a, \mu, T \dots$

- Describes the **current** at sufficiently high temperatures, in the **Ricci-flat space-time** $R_{\mu\nu} = 0$ e. g. in the space around the black hole?

DECOMPOSITION OF THE TENSORS: GRAVITY

Properties

$$A_{\mu\nu} = A_{\nu\mu}, \quad C_{\mu\nu} = C_{\nu\mu}, \quad B^{\mu}{}_{\mu} = 0, \\ A_{\mu\nu}u^{\nu} = C_{\mu\nu}u^{\nu} = B_{\mu\nu}u^{\nu} = B_{\nu\mu}u^{\nu} = 0.$$

The gravitational field is external. For simplicity, we impose an **additional condition**:

(for example, the field around a black hole) $R_{\mu\nu} = 0$

Additional properties appear, similar to

[L. D. Landau and E. M. Lifschits,
The Classical Theory of Fields, Vol. 2, 1975]

$$A_{\mu\nu} = -C_{\mu\nu}, \quad A^{\mu}{}_{\mu} = 0, \quad B_{\mu\nu} = B_{\nu\mu}$$

There are 10 independent components left

KVE IN RSA THEORY: CALCULATION

- Apply **point splitting** to all operators (no additional contributions arise - operators satisfy free field equations):

$$\hat{T}^{\mu\nu}(X) = \lim_{X_1, X_2 \rightarrow X} \mathcal{D}_{ab(IJ)}^{\mu\nu}(\tilde{\partial}_{X_1}, \tilde{\partial}_{X_2}) \bar{\Psi}_{aI}(X_1) \Psi_{bJ}(X_2),$$

$$\hat{j}_A^\mu(X) = \lim_{X_1, X_2 \rightarrow X} \mathcal{J}_{ab(IJ)}^\mu \bar{\Psi}_{aI}(X_1) \Psi_{bJ}(X_2),$$

where

$$X_\mu = (\tau_x, -\mathbf{x})$$

Fields are combined into one vector $\Psi_I = \{\tilde{\psi}_\mu, \lambda\}$ ($I = 0 \dots 4$)

The matrix element has the form of a product of **vertices** and **propagators**.

Vertices $\mathcal{J}_{(ij)}^\mu = i^{1-\delta_{0\mu}} \varepsilon^{ij\mu\nu} \tilde{\gamma}_\nu$

Euclidean Dirac matrices

$$\{\tilde{\gamma}_\mu, \tilde{\gamma}_\nu\} = 2\delta_{\mu\nu}$$

$$\mathcal{D}_{(ij)}^{\mu\nu} = -\frac{1}{2}(-i)^{\delta_{0\mu}+\delta_{0\nu}} \varepsilon^{ij\nu\beta} \left(\gamma_5 \tilde{\gamma}_\mu \tilde{\partial}_\beta^{X_2} - \frac{1}{4} \gamma_5 \tilde{\gamma}_\beta [\tilde{\gamma}_\nu, \tilde{\gamma}_\mu] \left(\tilde{\partial}_\nu^{X_1} + \tilde{\partial}_\nu^{X_2} \right) \right) + (\mu \leftrightarrow \nu)$$

$$0 \leq (i, j) < 4$$

Propagators

$$\langle T_\tau \tilde{\psi}_{a\mu}(X_1) \tilde{\psi}_{b\nu}(X_2) \rangle_T = \sum_P e^{iP_\alpha(X_1-X_2)^\alpha} \frac{i}{2P^2} \left(\tilde{\gamma}_\nu \not{P} \tilde{\gamma}_\mu + 2 \left[\frac{1}{m^2} - \frac{2}{P^2} \right] P_\mu P_\nu \not{P} \right)_{ab}$$

$$\langle T_\tau \tilde{\psi}_{a\mu}(X_1) \bar{\lambda}_b(X_2) \rangle_T = \sum_P e^{iP_\alpha(X_1-X_2)^\alpha} \frac{-P_\mu \not{P}_{ab}}{mP^2}$$

Mixed terms are non-zero

here $P_\mu = (p_n, -\mathbf{p})$, $p_n = (2n+1)\pi T$

$$\langle T_\tau \lambda_a(X_1) \bar{\lambda}_b(X_2) \rangle_T = 0$$

Field λ is **non-propagating**

KVE IN RSA THEORY: CALCULATION

Substituting the **split** form of the operators into the typical correlator C, we obtain:

$$C^{\alpha_1\alpha_2|\alpha_3\alpha_4|\alpha_5\alpha_6|\lambda|ijk} = -T^3 \int [d\tau] d^3x d^3y d^3z x^i y^j z^k \lim_{X,Y,Z,F} \mathcal{D}_{a_1 a_2 (I_1 I_2)}^{\alpha_1 \alpha_2}(\tilde{\partial}_{X_1}, \tilde{\partial}_{X_2}) \mathcal{D}_{a_3 a_4 (I_3 I_4)}^{\alpha_3 \alpha_4}(\tilde{\partial}_{Y_1}, \tilde{\partial}_{Y_2}) \mathcal{D}_{a_5 a_6 (I_5 I_6)}^{\alpha_5 \alpha_6}(\tilde{\partial}_{Z_1}, \tilde{\partial}_{Z_2}) \\ \times \mathcal{J}_{a_7 a_8 (I_7 I_8)}^\lambda \langle T_\tau \bar{\Psi}_{a_1 I_1}(X_1) \Psi_{a_2 I_2}(X_2) \bar{\Psi}_{a_3 I_3}(Y_1) \Psi_{a_4 I_4}(Y_2) \bar{\Psi}_{a_5 I_5}(Z_1) \Psi_{a_6 I_6}(Z_2) \bar{\Psi}_{a_7 I_7}(F_1) \Psi_{a_8 I_8}(F_2) \rangle_{T,c}$$

We transform the average of 8 fields using the **Wick theorem**

$$\langle \bar{\Psi}(X) \Psi(X) \bar{\Psi}(Y) \Psi(Y) \bar{\Psi}(Z) \Psi(Z) \bar{\Psi}(F) \Psi(F) \rangle_c = -\langle \Psi(Y) \bar{\Psi}(X) \rangle \langle \Psi(X) \bar{\Psi}(F) \rangle \langle \Psi(Z) \bar{\Psi}(Y) \rangle \langle \Psi(F) \bar{\Psi}(Z) \rangle + (5 \text{ terms})$$

- Initially there are 6×5^8 terms, but we should take into account, that:

- 1) some terms are zero because they include propagator $\langle \lambda \bar{\lambda} \rangle = 0$
- 2) some vertices are zero, e.g. $\mathcal{D}_{(ij)}^{\mu\nu} = 0$ if $i = j \neq 4$
- 3) some terms are zero in the limit $m \rightarrow \infty$ if the total number of propagators $\langle \lambda \bar{\psi} \rangle$ and $\langle \psi \bar{\lambda} \rangle$ is greater than the total number of vertices $\mathcal{D}_{(4i)}$ and $\mathcal{D}_{(i4)}$

Then there are:

94752 terms for each C correlator in λ_1

31152 terms for each C correlator in λ_2

KVE IN RSA THEORY: CALCULATION

- Our *goal* is to calculate the conductivities λ_1 and λ_2 in the KVE current:

$$j_{A,KVE}^\mu = \lambda_1(\omega_\nu \omega^\nu) \omega^\mu + \lambda_2(a_\nu a^\nu) \omega^\mu$$

- Using the described perturbation theory, we obtain:

$$\lambda_1 = -\frac{1}{6} \int_0^{|\beta|} [d\tau] \langle T_\tau \hat{J}_{-i\tau_x}^3 \hat{J}_{-i\tau_y}^3 \hat{J}_{-i\tau_z}^3 \hat{j}_A^3(0) \rangle_{T,c}$$

- Representing \hat{J}_σ , \hat{K}^μ through the stress-energy tensor, we obtain:

$$\lambda_1 = -\frac{1}{6T^3} \left(C^{02|02|02|3|111} + C^{02|01|01|3|122} + C^{01|02|01|3|212} + C^{01|01|02|3|221} \right. \\ \left. - C^{01|01|01|3|222} - C^{01|02|02|3|211} - C^{02|01|02|3|121} - C^{02|02|01|3|112} \right),$$

Typical correlator to be found: 4-point one-loop function

$$C^{\alpha_1\alpha_2|\alpha_3\alpha_4|\alpha_5\alpha_6|\lambda|ijk} = T^3 \int [d\tau] d^3x d^3y d^3z x^i y^j z^k \langle T_\tau \hat{T}^{\alpha_1\alpha_2}(-i\tau_x, \mathbf{x}) \hat{T}^{\alpha_3\alpha_4}(-i\tau_y, \mathbf{y}) \hat{T}^{\alpha_5\alpha_6}(-i\tau_z, \mathbf{z}) \hat{j}_5^\lambda(0) \rangle_{T,c}$$

When expanding the density operator \rightarrow shift along the **imaginary** axis \rightarrow field theory at **finite temperatures**.