

Polarizations in inclusive processes

M.G. Ryskin, Sov.J.Nucl.Phys. 48 (1988) 708

The model is simple
and contains the transparent physics.

Shortness –
not clear how to calculate the corrections

1. Polarization in pert. QCD is small

$$P \sim \alpha_s m_q / q_T$$

2. Non-pert. model

3. Matching with pert.QCD

4. Hyperon polarization

5. Inclusive pion asymmetry

Polarization need the shift of the phase between the 'spin-flip' and 'non-flip' amplitudes.

$$P \propto \text{Im}(M_{n.f.} M_{s.f.}^*)$$

The quark-gluon vertex γ_μ conserves the helicity.
Quark mass in the propagator needed to flip the helicity.

Next to generate the imaginary part an extra gluon exchange is needed.

Thus in pert. QCD

$$P \sim \frac{\alpha_s m_q}{\pi q_T} < 1\%$$

for $q_T = 2 \text{ GeV}$, $\alpha_s = 0.3$ and $m_s = 150 \text{ MeV}$
(we use the *current* quark mass in pert.QCD)

However $P_\Lambda \sim 0.3$ was observed experimentally !




Figure 1. Λ polarization in the pp collision at $p_T = 1\text{GeV}/c$ with the experimental data.




Figure 2. Σ polarization in the pp collision at $p_T = 1\text{GeV}/c$ with the experimental data.

Non-Pert. model

After the gluon exchange
the colour tube/string is created.
The string with the colour-electric field is unstable.
The colour-magnetic field is appeared,
like around the conductor with electric current in QED.

The strength of this colour-magnetic field is

$$H \simeq \frac{\sqrt{\alpha_s(R_c)}}{1.6R_c^2}$$


(“THE STRUCTURE OF A GLUON STRING”

A.B. Migdal S.B. Khokhlachev, JETP Lett. 41 (1985) 194.)

$$R_c \sim 1/400\text{MeV} \quad \alpha_s(R_c) \sim 0.5$$

Here it is better to use the constituent quark mass.

$$m_q = 330 \text{ MeV} \text{ and } m_s = 500 \text{ MeV}$$



The μH interaction leads to an *additional* quark transverse momentum

$$\delta q_T \simeq \mu H \simeq \frac{\alpha_s C_F}{2m_q 1.6R_c^2} \approx 100 \text{MeV}$$

Asymmetry of a polarized quark production

$$A = \frac{d\sigma/d^3q_{\uparrow} - d\sigma/d^3q_{\downarrow}}{d\sigma/d^3q_{\uparrow} + d\sigma/d^3q_{\downarrow}} = \frac{d\sigma(q_T + \delta q_T) - d\sigma(q_T - \delta q_T)}{d\sigma(q_T + \delta q_T) + d\sigma(q_T - \delta q_T)} =$$

$$= \delta q_T \frac{\partial}{\partial q_T} \left(\frac{d\sigma}{d^3q} \right) \left(\frac{d\sigma}{d^3q} \right)^{-1},$$

This can be compared with the pert.QCD result


The interference of Fig.1a and 1b diagrams gives

$$A = 0,95 \alpha_s q_T m / (q_T^2 + m^2),$$

corresponding to $\delta q_T = \alpha_s m_q / 2 \simeq 80 \text{ MeV}$

since

$$\frac{\partial}{\partial q_T} \left(\frac{d\sigma}{dq_T^2} \right) \left(\frac{d\sigma}{dq_T^2} \right)^{-1} = \frac{2q_T}{m^2 + q_T^2},$$



Single spin asymmetry for $p_{\uparrow}p \rightarrow \pi + X$ in pert.QCD.

M. Anselmino, M. Boglione, F. Murgia

Phys. Lett.B 362 (1995) 164

$$\frac{E_{\pi} d\sigma^{p_{\uparrow}p \rightarrow \pi X}}{d^3\mathbf{p}_{\pi}} \sim \frac{1}{2} \sum_{a,b,c,d} \sum_{\lambda_a, \lambda'_a; \lambda_b; \lambda_c, \lambda'_c; \lambda_d} \int d^2\mathbf{k}_{\perp a} dx_a dx_b \frac{1}{z} \rho_{\lambda_a, \lambda'_a}^{a/p_{\uparrow}} \hat{f}_{a/p_{\uparrow}}(x_a, \mathbf{k}_{\perp a}) f_{b/p}(x_b) \hat{M}_{\lambda_c, \lambda_d; \lambda_a, \lambda_b} \hat{M}_{\lambda'_c, \lambda_d; \lambda'_a, \lambda_b}^* D_{\pi/c}^{\lambda_c, \lambda'_c}(z),$$

where \hat{f} denotes the \mathbf{k}_{\perp} dependent number density.

$$\begin{aligned} A_N \propto \Delta^N f_{a/p_{\uparrow}}(x_a, \mathbf{k}_{\perp a}) &\equiv \sum_{\lambda_a} \left[\hat{f}_{a, \lambda_a/p_{\uparrow}}(x_a, \mathbf{k}_{\perp a}) - \hat{f}_{a, \lambda_a/p_{\downarrow}}(x_a, \mathbf{k}_{\perp a}) \right] \\ &= \sum_{\lambda_{\perp}} \left[\hat{f}_{a, \lambda_a/p_{\uparrow}}(x_a, \mathbf{k}_{\perp a}) - \hat{f}_{a, \lambda_a/p_{\uparrow}}(x_a, -\mathbf{k}_{\perp a}) \right] \end{aligned}$$

Λ -hyperon polarization

$$\Lambda_\lambda = s_\lambda + (ud)_{scalar} \text{ that is } P_\Lambda = P_{s-quark}$$

For $d\sigma/dq_T^2$ we use

$$x d\sigma/dx dq_T^2 \propto \exp(-4xq_T - 1,9q_T^2 + 0,66q_T^3 - 0,07q_T^4)$$

Pondrom L. G. Univ. of Wisconsin Thesis, 1984.

$$(p + Be \rightarrow \Lambda + X \quad 400 \text{ GeV}) \quad q_T > 1 \text{ GeV}$$

This gives $P_\Lambda = -0,28x - 0,27q_T + 0,14q_T^2 - 0,02q_T^3$.

$|P_\Lambda|$ increases with x and q_T (up to $q_T \simeq 1.5 \text{ GeV}$)

At small q_T the cross section $d\sigma/dq_T^2 \propto \exp(-Bq_T^2)$ leading to $P_\Lambda \propto q_T$.

Asymmetry in $p \uparrow p \rightarrow p + X$ is small

For the Pomeron exchange (PPP and PPR terms) we have no colour string near the leading proton.

In the case of RRP and RRR terms the leading valence quarks pick up the new slow quark which mainly goes to the scalar diquark. That is the polarization is lost.

Inclusive $p_{\uparrow}p \rightarrow \pi + X$ pion asymmetry

$$A_{\pi} = A_q P_q \sigma(q) / (\sigma(q) + \sigma(g))$$


A_q is the quark asymmetry

P_q is the quark polarization in p_{\uparrow} ,

factor $\sigma(q) / (\sigma(q) + \sigma(g))$ is the fraction of the pions created by this (polarized) quark.

In first approx. $P_q \simeq x$

with $\delta q_T = 100$ MeV and $d\sigma/dq_T^2 \propto \exp(-4.5q_T)$ we get



Conclusion

A simple model is suggested which enables one to evaluate the polarization effects in inclusive reactions making use of the q_T dependence of the x-section $E d\sigma / d^3q$

Other non-pert. models

A SEMICLASSICAL MODEL FOR THE POLARIZATION

Bo ANDERSSON, G. GUSTAFSON, G. INGELMAN, Ph.Lett. 85B (1979) 417

tunnelling $\bar{q}q$ pair production
local conservation of angular momentum ($L=-S$) and k_T




Fig. 5.20. A quark and an antiquark with transverse momenta \bar{k}_\perp and $-\bar{k}_\perp$ are produced at a distance $2\mu_\perp/\kappa$ from each other. They carry an orbital angular momentum \bar{L} which is compensated if the spins are polarized in the opposite direction.

$$L \propto \mu_\perp = \sqrt{k_T^2 + m_q^2}$$

Models for polarization asymmetry in inclusive hadron produc.
T.A. DeGrand, H.I. Miettinen, Ph. Rev. D24 (1981) 2419

Quark spin is rotated due to Thomas precession
when the *see* quark is accelerated.




FIG. 3. Semiclassical trajectory of a particle in an attractive potential showing the distance of the orbit from the origin \vec{r} , and the orientations of the scattering plane

A New Mechanism for Single Spin Asymmetries in Strong Interactions

JETP Lett. 72 (2000) 481

N.I. Kochelev




Figure 2: *The instanton contribution to the single spin asymmetry for pion production as a function of x_F .*





Figure 3: *The instanton contribution to the single spin asymmetry for pion production as a function of x_F and $p_t = |l_\perp|$. Solid line is for $x_F = 0.9$, dashed line is for $x_F = 0.6$ and dotted line is for $x_F = 0.3$.*

SINGLE SPIN ASYMMETRIES IN HIGH ENERGY REACTIONS AND NON-PERT. QCD EFFECTS

A. E. Dorokhov, N. I. Kochelev, W.-D. Nowak

Phys.Part.Nucl.Lett. 6 (2009) 440-445 • e-Print: 0902.3165



The diagrams giving rise to SSA in SIDIS. The symbol I denotes the instanton

QCD Instanton flips the quark spin !

Instanton is the classical solution of QCD equations.

A.A.Belavin, A.M.Polyakov, A.S.Schwartz and Yu.S.Tyupkin,

Phys.Lett.**59B**, 85 (1975)

$$A_{\mu}^a(x) = \frac{2}{g} \eta_{a\mu\nu} \frac{(x - x_0)_{\nu}}{(x - x_0)^2 + \rho^2}$$

$\alpha_s = g^2/4\pi$, $\rho =$ instanton radius,

$\eta_{a\mu\nu} = 0$ for $\mu = \nu = 4$

$\eta_{a\mu\nu} = -\delta_{a\nu}$ for $\mu = 4$

$\eta_{a\mu\nu} = \delta_{a\nu}$ for $\nu = 4$

$\eta_{a\mu\nu} = \epsilon_{a\mu\nu}$ for $\mu, \nu = 1, 2, 3$

At $x \rightarrow \infty$ instanton is the pure gauge field

$$g \frac{\tau^a}{2} A_\mu^a \rightarrow i S \partial_\mu S^+$$

with $S = i \tau_\mu^+ x_\mu / \sqrt{x^2}$

However for $x \neq \infty$ it is the real transverse gluon field which describes the transition between two different (in gauge) QCD vacuums.




Figure 1. Instanton and Sphaleron processes in the topology of a Yang-Mills vacuum; energy density of the gauge field (y-axis) vs. winding number N_{CS} (x-axis).

The statistical weight of size- ρ instanton is

$$D(\rho, \mu_R) = \frac{\kappa}{\rho^5} \left(\frac{2\pi}{\alpha_s(\mu_R)} \right)^6 (\rho\mu_R)^{b_0}$$

where $(\rho\mu_R)^{b_0} = \exp(-2S^I)$ $S^I = 2\pi/\alpha_s$

$$\kappa = 0.0025 \exp(0.292N_f) \sim 0.01$$

Instanton induced spin-spin corrⁿ.

$$q_L + q_L \implies I \implies n \cdot g + \sum_f (q'_{Rf} + \bar{q}_{Lf}) \quad n \sim 1/\alpha_s(\rho)$$

$$q_R + q_R \implies \bar{I} \implies n \cdot g + \sum_f (q'_{Lf} + \bar{q}_{Rf})$$

Instanton rearrange the Dirac basement

One extra level of light left quark appears while the level of right quark goes upstairs to continuum spectra.


In electro-weak case where the γ_5 anomaly is canceled between the quarks and the leptons this leads to the *baryon charge non-conservation*.

In QCD this is the helicity non-conservation.

In QCD this is the helicity non-conservation.

This can be checked experimentally by studying the spin-spin correlations, say in

$$p_{\uparrow} + p \rightarrow \Sigma^0 + X$$



Conclusion

A simple model is suggested which enables one to evaluate the polarization effects in inclusive reactions making use of the q_T dependence of the x-section $E d\sigma / d^3q$

THANK YOU