

Possibilities for centrality determination in the BM@N experiment

Ilya Segal, Arkadiy Taranenko, Peter Parfenov,
Mikhail Mamaev, Dim Idrisov
for the BM@N Collaboration



This work is supported by:
the NRNU program Priority 2030 and
the Special Purpose Funding Programme within the NICA Megascience Project in 2023



May 16th, 2023
10th BM@N Collaboration Meeting




Motivation for centrality determination

- Evolution of matter produced in heavy-ion collisions depends on its initial geometry
- Goal of centrality determination:
map (on average) the collision geometry parameters
to experimental observables (centrality estimators)

- Monte-Carlo sampling based on output of Glauber model
is commonly used to build such connection

- Centrality class S_1 - S_2 : group of events corresponding to
a given fraction (in %) of the total cross section:

$$C_S = \frac{1}{\sigma_{inel}^{AA}} \int_{S_1}^{S_2} \frac{d\sigma}{dS} dS$$




Why several alternative centrality estimators


Anticorrelation between charge of the spectator fragments (FW) and particle multiplicity (hits)

A number of produced protons is stronger correlated with the number of produced particles (track & RPC+TOF hits) than with the total charge of spectator fragments (FW)

HADES; Phys.Rev.C 102 (2020) 2, 024914



HADES; Phys.Rev.C 102 (2020) 2, 024914






Avoid self-correlation biases when using spectators fragments for centrality estimation

Types of centrality estimators

Produced charged particles

Spectators



(Target spectators
not measured for
fixed-target)


BM@N subsystems for centrality determination

Simulation setup

- DCM-QGSM-SMM
[M.Baznat et al. PPNL 17 \(2020\) 3, 303](#)
- Xe-Cs @ 4A GeV
- Transport: GEANT4

Subsystems

- Participants: **Tracking system**
GEM+STS, BD, SiMD
- Spectators: **FHCal**, **Hodoscope**,
ScWall, FD



Overview of centrality determination methods

Method type	MC-Glauber based	Model independent (e.g. Γ -fit method)	Based on ML
Used in	STAR, ALICE, HADES, CBM, MPD, etc.	ALICE, CMS, ATLAS <small>J. Y. Ollitrault et al. Phys.Rev. C 98 (2018) 024902</small>	Becoming popular <small>Fupeng L. et al. J.Phys.G 47 (2020) 11, 115104</small>
Advantages	Commonly used, well established procedure	Universality due to model independence	The most modern and fast methods
Disadvantages	MC-Glauber model provides non-realistic N_{part} simulations at low energies <small>M. O. Kuttan et al. e-Print: 2303.07919 [hep-ph]</small>	In strong connection with σ_{inel} which dependence on energy is not well studied at low energies (same problem for MC-Glauber based methods)	There no way to control the physicality of the methods

Centrality determination based on Monte-Carlo sampling

For **multiplicity of produced particles**
used in HADES, CBM, BM@N, NA61/SHINE

Get $(N_{\text{part}}, N_{\text{coll}})$ from MC-Glauber

Calculate $N_a = fN_{\text{part}} + (1-f)N_{\text{coll}}$

Sample multiplicity of produced particles (S_i) N_a times
from NBD (μ, k)

Result: total S_{tot}

MC-Glauber
distribution



Full Monte-Carlo (real
data) distribution

Evaluate χ^2
between $dN/dE_{\text{MC/data}}$ and dN/dE_{Gl}

Scan phase space of parameters
to find their values for minimum of χ^2

Extract relation between geometry
parameters and centrality estimator

MC-Glauber fit result Xe-Cs @ 4.0 AGeV



$\chi^2=1.31\pm 0.07$;
 $f=0.9$,
 $\mu=0.786293$,
 $k=1$;
MinFitBin=10,
MaxFitBin=250


- Fit result is good
- Impact parameter distributions in different centrality classes reproduces ones from DCM-QGSM-SMM

The Bayesian inversion method (Γ -fit): main assumptions


- Relation between multiplicity N_{ch} and impact parameter b is defined by the fluctuation kernel:

$$P(N_{ch}|c_b) = \frac{1}{\Gamma(k(c_b))\theta^k} N_{ch}^{k(c_b)-1} e^{-N_{ch}/\theta}$$

$$c_b = \int_0^b P(b') db' \simeq \frac{\pi b^2}{\sigma_{inel}} \quad \text{-- centrality based on impact parameter}$$



The results of fitting the multiplicity distribution for a fixed impact parameter



The dependence of the average value of multiplicity on centrality and the results of its fit

$$\frac{\sigma^2}{\langle N_{ch} \rangle} = \theta \simeq const$$

$$\langle N_{ch} \rangle = N_{knee} \exp\left(\sum_{j=1}^3 a_j c_b^j\right), k = \frac{\langle N_{ch} \rangle}{\theta}$$

Five fit parameters

N_{knee}, θ, a_j

Reconstruction of b

- Normalized multiplicity distribution $P(N_{ch})$

$$P(N_{ch}) = \int_0^1 P(N_{ch}|c_b)dc_b$$

- Find probability of b for fixed range of N_{ch} using Bayes' theorem:

$$P(b|n_1 < N_{ch} < n_2) = P(b) \frac{\int_{n_1}^{n_2} P(b|N_{ch})dN_{ch}}{\int_{n_1}^{n_2} P(N_{ch})dN_{ch}}$$



- The Bayesian inversion method consists of 2 steps:**

- Fit normalized multiplicity distribution with $P(N_{ch})$
- Construct $P(b|N_{ch})$ using Bayes' theorem with parameters from the fit



R. Rogly, G. Giacalone and J. Y. Ollitrault, Phys.Rev. C98 (2018) no.2, 024902

Implementation for MPD and BM@N by D. Idrisov: <https://github.com/Dim23/GammaFit>

Example of application in MPD: P. Parfenov et al., Particles 4 (2021) 2, 275-287




Γ -fit result Xe-Cs @ 4.0 AGeV







- Fit result is good
- Impact parameter distributions in different centrality classes reproduces ones from DCM-QGSM-SMM
- Γ -fit method also could be used for centrality determination based on spectators energy

Possibilities of spectators fragments as estimators




- Physical threshold of switching between estimators could be Hodoscope signal $E_{\text{Hodo}} = 0.04$ (corresponding to $b \sim 6$ fm)
- FHCal energy distribution improved and has more linear correlation with impact parameter (for range $E_{\text{Hodo}} < 0.04$)
- There is good correlation between Hodoscope charge and impact parameter (for range $E_{\text{Hodo}} > 0.04$)

Possibilities of spectators fragments as estimators



Comparison of different estimators and methods



from talk at ICPPA-2022

proceedings submitted
to Physics of Atomic Nuclei

- Impact parameter distributions in different centrality classes are similar for different centrality estimators
- These distributions for spectators energy is wider because of the width of b and energy correlation



NA61/SHINE experimental setup

Data samples:

- Pb-Pb @ $p_{\text{beam}} = 13A \text{ GeV}/c$
- data from 2016 physics run
- DCM-QGSM-SMM x Geant4
[M.Baznat et al. PPNL 17 \(2020\) 3, 303](#)

Subsystems

- Multiplicity: TPCs ($p_T > 0.05$, $\eta < 3.5$)
- Spectators energy: PSD



Centrality determination based on Monte-Carlo sampling

For **multiplicity of produced particles** used in HADES, CBM, BM@N, NA61/SHINE

Get (N_{part}, N_{coll}) from MC-Glauber

$$\text{Calculate } N_a = fN_{part} + (1-f)N_{coll}$$

Sample multiplicity of produced particles (S_i) N_a times from NBD (μ, k)

Result: total S_{tot}

For **spectators energy** from hadron calorimeters used in NA61/SHINE

Get (N_{spec}, b) from MC-Glauber

Sample hadron calorimeter response (S_i) N_{spec} times from Gauss (μ, k)

MC-Glauber distribution



Full Monte-Carlo (real data) distribution

Evaluate χ^2 between $dN/dE_{MC/data}$ and dN/dE_{GI}

Scan phase space of parameters to find their values for minimum of χ^2

Extract relation between geometry parameters and centrality estimator




Simplified MC sampling for hadron calorimeters



see for more details Segal I. Particles. 2023; 6(2):568-579.

- Gauss distribution can not reproduce energy distribution in the most central collisions
- Possible improvements are now under investigation

Comparison with standard MC-Glauber x NBD method



see for more details Segal I. Particles. 2023; 6(2):568-579.

- Centrality classes determined separately using the multiplicity of produced particles and spectators are slightly different
- This is due to the different shapes of two-dimensional distributions of impact parameters and corresponding centrality estimators
- Impact of this effect should be considered during further work

Summary

- Software implementation of MC-Glauber and Γ -fit based fitting procedures for multiplicity are used for BM@N
- Relation between impact parameter and centrality classes is extracted
- Combination of forward detectors can be used to avoid effects due to the beam hole in FHCaI
- Centrality determination procedure based on MC sampling of spectators energy is developed and tested based on NA61/SHINE data
- Results are tuned on the spectator production implemented in the DCM-QGSM-SMM model

Work in progress

- Apply developed procedures based on spectators for BM@N FHCaI
- Investigate applicability of the Glauber model for centrality determination at low energies
- Consider using of Γ -fit method for spectators energy
- Apply all procedures for BM@N run8 data

Backup

MC Glauber model

MC Glauber model provides a description of the initial state of a heavy-ion collision

- Independent straight line trajectories of the nucleons
- A-A collision is treated as a sequence of independent binary NN collisions
- Monte-Carlo sampling of nucleons position for individual collisions

Main model parameters

- Colliding nuclei

- Inelastic nucleon-nucleon cross section ($\sigma_{\text{inel}}^{\text{NN}}$)
(depends on collision energy)

- Nuclear charge densities (Wood-Saxon distribution)

$$\rho(r) = \rho_0 \cdot \frac{1 + w(r/R)^2}{1 + \exp\left(\frac{r-R}{a}\right)}$$

Geometry parameters


b – impact parameter

N_{part} – number of nucleons participating in the collision

N_{spec} – number of spectator nucleons in the collision


N_{coll} – number of binary NN collisions

Glauber Modeling in High Energy Nuclear Collisions:
ARNPS57:205-243,2007





SMM description of the ALADIN's fragmentation data



A.S. Botvina et al. NPA 584 (1995) 737



R.Ogul et al. PRC 83, 024608 (2011)




Respond of FHCaI detector



- Mean of signal has linear dependency with beam energy

Gaussian approximation for fragments energy




- Distribution of mass numbers of spectators fragments could be fitted by Gauss distribution
- Mean values equal to product of beam energy and fragment's mass
- Total spectators energy distribution is also Gauss:

$$P(E_{tot}; \mu_{tot}, k_{tot}) \approx \prod_{i=1}^{N_{frag}} P(E_{frag}^i; \mu_{frag}^i, k_{frag}^i) \approx \prod_{i=1}^{N_{spec}} P(E_{spec}^i; \mu_{spec}^i, k_{spec}^i)$$

- Measured energy distribution follows convolution of two Gauss distributions (sum of fragments energy and detector response)


Simplified MC sampling for hadron calorimeters




Segal I. Particles. 2023; 6(2):568-579.

- Shapes of energy and impact parameter distributions are similar
- Width of distribution for energy is larger than for multiplicity
- Possible decrease of width will be study




Centrality determination using inverse Bayes approaches



- Centrality determination based on spectator energy using inverse Bayes approach is being developed and tested on model (UrQMD, DCM-QGSM-SMM) and NA61/SHINE data
- Application of centrality determination based on spectator energy using MC-Glauber and inverse Bayes approaches is in progress
- Possible improvements are under investigation



Result of the fitting



NBD at different values of k

