

Inclusive π^0 production for online polarimetry in SPD



Katherin Shtejer Díaz


SPD Collaboration Meeting
27.04.2023

Inclusive π^0 for local polarimetry

$$A_N = \frac{\sigma^\uparrow - \sigma^\downarrow}{\sigma^\uparrow + \sigma^\downarrow}$$





A_N is a measure of the beam polarization









- ❑ SpdRoot version 4.1.5.1
- ❑ Two energies: $pp @ \sqrt{s} = 13 \text{ GeV}$ and $pp @ \sqrt{s} = 27 \text{ GeV}$
- ❑ Particle generator: Pythia 8 (number of events: $\sim 100\text{M}$)
- ❑ Minimum Bias: *SoftQCD:inelastic* \leftrightarrow inelastic, non diffractive events and diffractive topologies
- ❑ **MC truth info!**
- ❑ Vertex assumed at (0, 0, 0) \rightarrow Gaussian smeared: $\sigma_z = 30 \text{ cm}$ and $\sigma_{x,y} = 0.1 \text{ cm}$.
- ❑ Photon trajectory extrapolated to the ECAL endcap “planes”.
- ❑ z position fixed, assuming ECALTECMinDist = 188.6 cm
- ❑ Energy of the MC-particle, smeared by $\frac{\sigma_E}{E} = 2\% \oplus \frac{5.5\%}{\sqrt{E}}$
- ❑ $E_{min}^\gamma = 400 \text{ MeV}$
- ❑ π^0 selected from the M_{inv} of $\gamma\gamma$ pairs

General characteristics



$pp @ \sqrt{s} = 27 \text{ GeV}$

$pp @ \sqrt{s} = 13 \text{ GeV}$



$$p^\uparrow + p \rightarrow \pi^0 + X \quad \phi = 2\pi$$

The cross section of hadron production in polarized $p^\uparrow + p$ collisions, is modified in azimuth.

$$\frac{d\sigma}{d\varphi} = \frac{d\sigma}{d\varphi_0} [1 + \underbrace{P \cdot A_N \cdot \cos(\varphi + \varphi_0)}_{\text{Azimuthal cosine modulation}}]$$

Azimuthal cosine modulation


$$N_{\pi^0}(\varphi) = A[1 + P \cdot A_N \cdot \cos(\varphi + \varphi_0)]$$

$$A_N = \frac{Amp}{P}$$

$N_{\pi^0}(\varphi)$: Yield of π^0


P : Beam polarization

- $P = 0.7$ was assumed




The spin dependent π^0 yields for each bin are extracted from the invariant mass spectra in different x_F sub-ranges for each φ bin.



Azimuthal cosine modulation of π^0 yields in x_F intervals, $[p0] \cdot (1 + [p1] \cdot \cos([p2] + x))$



Azimuthal cosine modulation of π^0 yields in x_F intervals, $[p_0] \cdot (1 + [p_1] \cdot \cos([p_2] + x))$



A_N vs. x_F





$$\frac{\Delta A_N}{A_N} \text{ vs. } x_F$$

$$\frac{\Delta A_N}{A_N} \begin{matrix} \nearrow \text{SpdRoot} \\ \searrow \text{E704} \end{matrix}$$

$$\frac{\Delta A_N}{A_N} \sim \frac{\Delta P}{P}$$

$$\sigma_{tot}^{pp}(27 \text{ GeV}) = 40.0 \text{ mb} \quad \mathcal{L} = 10^{32} s^{-1} cm^{-2} \quad \text{Reaction rate (27 GeV)} = 4 \cdot 10^6 s^{-1}$$

$$\sigma_{tot}^{pp}(13 \text{ GeV}) = 38.4 \text{ mb} \quad \mathcal{L} = 10^{31} s^{-1} cm^{-2} \quad \text{Reaction rate (13 GeV)} = 3.8 \cdot 10^5 s^{-1}$$



Better precision of the polarization measurement expected at:
 $0.3 < x_F < 0.4$


Raw asymmetry:

$$P \cdot A_N \cdot \cos \phi = \epsilon(\phi) \Leftrightarrow \epsilon(\phi) = \frac{N^\uparrow(\phi) - N^\downarrow(\phi)}{N^\uparrow(\phi) + N^\downarrow(\phi)}$$

$$P \cdot A_N \sim \epsilon$$

$$\frac{\Delta A_N}{A_N} \sim \frac{\Delta P}{P}$$

$$\frac{\Delta P}{P} = \frac{1}{\sqrt{\sum_i \left(\frac{A_{Ni}}{\Delta A_{Ni}}\right)^2}}$$




Taking the last experimental 4 points ($0.3 \leq x_F < 0.7$): $\frac{\Delta P}{P} = 0.0873 \sim 9\%$ (Experiment E704)

*The error of the beam polarization in the experiment **E704** is estimated in **10%**, as reported in FERMILAB-Pub-91/15-E[E581,E704]*

Estimation of the statistical accuracy of the beam polarization measurement, with $pp \rightarrow \pi^0 X$ at $\sqrt{s} = 13$ GeV and $\sqrt{s} = 27$ GeV, in SPD ECAL endcaps.

Estimated time	$\frac{\Delta P}{P}$	
	$\sqrt{s} = 13$ GeV	$\sqrt{s} = 27$ GeV
2 min	2.44 %	0.72 %
5 min	1.54 %	0.45 %
10 min	1.09 %	0.32 %
20 min	0.77 %	0.23 %
30 min	0.63 %	0.18 %
1 h	0.45 %	0.13 %



- The accuracy of the beam polarization have been estimated at two different pp collision energies: 13 GeV and 27 GeV
- The determination of the polarization is expected to be more precise in the range $0.3 < x_F < 0.4$.
- From the asymmetry determination, based on MC truth simulations with SpdRoot, the statistical accuracy of the beam polarization, for 10 minutes, is estimated in: 1.1 % at 13 GeV and 0.32 % at 27 GeV.
- The inclusive $pp \rightarrow \pi^0 X$ reaction, detected in the ECAL Endcaps, is proposed to participate in the local polarimetry at SPD, by measuring and monitoring the transverse single spin asymmetry.
- Main difficulty: the few availability of accurate experimental data in the energy range of interest for SPD.

Many thanks to Andrei Maltsev for his contribution
on the MC info management for the ECAL!