

# Cluster particle production @ SPD experiment

D. Budkouski, I. Lapushanskii, A. Tumasyan

SPD meeting, 22 March, 2023

# Experience we have in such topic

- Hadron jets reconstruction in CMS experiment at LHC
- Jet energy scale calibration in CMS experiment using “ $\gamma+jet$ ” process and  $W \rightarrow qq$  decay
- Investigation of Jet energy resolution and treatment in CMS experiment
- Implementation of flavor tagging (e.g. b-tagging) for jet reconstruction algorithms in CMS
- Measurement of gluon jets fraction in inclusive pp-collisions using algorithms for quark-gluon discrimination
- Investigation of particles multiplicity of hadron jets in pp-collisions
- Jet energy regression using ML methods in CMS experiment
- Jet triggers development in CMS experiment


# Motivation

- Partons - products of hadron-hadron hard scattering are not accessible for direct measurement
- We can get an information about these particles from the final state products resulting from harmonization of quark-gluon shower created by the initial parton
- When the energy of parton-initiator is *high enough* in the final state a *jet* of particles will be formed, which will correspond to initial parton with high accuracy:

$$\mathbf{p}^{\text{Jet}} (E, P_x, P_y, P_z) \approx \mathbf{p}^{\text{parton}} (E, P_x, P_y, P_z)$$

- There is no closed system for *transformation* of colored initial parton to final state colorless particles
- Initial parton can exchange momentum with beam remnants and other partons
- If the energy/momentum of parton-initiator is *high enough* than the momentum of exchange, the jet can be associated with the parton.
- The goals of this study:
  - Understand the admissibility of such approximation for energies at NICA and SPD experiment
  - Study processes of parton production at energy region between non-pQCD and pQCD

# Problem statement



## Performance of reconstruction for particles clustered production

- Search for clustered production of particles (efficiency)
- Reconstruction of parton-initiator kinematics depending on reconstructed jet characteristic
- Reconstruction of parton-initiator flavour depending on reconstructed jet characteristic

## Clustering algorithms and parameters

- Cluster/Jet reconstruction algorithm (Iterative Cone, kT, Anti-kT, Cambridge-Aachen, etc.)
- Radius parameter
- Inputs of clustering algorithms as objects of reconstruction and their kinematic thresholds
- Energy/momentum of reconstructed cluster

# *Anti- $k_t$* algorithm


- Jets are clustered with *anti- $k_t$*  algorithm
- Distance between objects in *anti- $k_t$*  algorithm defined as  $d_{ij} = \min\left(\frac{1}{k_{ti}^2}, \frac{1}{k_{tj}^2}\right) \frac{\Delta_{ij}^2}{R^2}$ ,  
where  $\Delta_{ij}^2 = (y_i - y_j)^2 + (\varphi_i - \varphi_j)^2$
- The functionality of the algorithm can be understood by considering an event with a few hard particles and many soft particles
  - If hard particle 1 has no hard neighbours within a distance  $2R$  then we have one perfectly conical jet
  - If another hard particle 2 is present such that  $R < \Delta_{12} < 2R$  then we have two jets with some overlapping parts
  - If distance between particles 1 and 2  $\Delta_{12} < R$  then both formed one jet

# Objects definition




- “True” jet:
  - Initial parton during evolution gives several final state particles
  - On generator level we can find all particles originated from hard process parton
  - Totality of these particles we call “true” jet
  - In ideal case “true” jets can associated with initial parton
- Clustered jets:
  - Clustering algorithms can find many jet-like objects in single event
  - But we want to choose only objects, which could be associated with initial parton
  - We use two methods based on gen information to find such objects
    - We take all jets, which are found by clustering algorithm and select the closest one by distance  $R$  to hard scattered parton, but the distance should be less than  $R$
    - We take highest  $p_T$  jet and check, at least one jet constituent originated from hard scattered parton
  - Both methods gives similar results

# Event generation and jet reconstruction settings

- We use Pythia8 generator and FastJet package
- We generate process:  $qg \rightarrow q\gamma$
- *anti-kt* algorithm with parameter  $R = 0.4, 0.6, 0.8$  was used for jet clustering
- Minimum jet  $p_T = 0.5\text{GeV}$
- Jet was clustered from final state particles with  $p_T > 0.25\text{ GeV}$  and  $\eta < 5$
- “True jets” and clustered jets are matched to hard scattered parton (status = 23)
- Hard scattered parton cuts:  $p_{T, parton} > 0\text{ GeV}, > 3\text{ GeV}, > 5\text{ GeV}$
- Jet should have at least two particles






# “True” jet vs parton (R=0.4)








# “True” jet vs parton (R=0.6)




# “True” jet vs parton (R=0.8)




# “True” jet vs parton $\eta$ resolution

With p and n






Without p and n






- Particle identification based on generator information about particle mothers may give ambiguous results
- Some protons and neutrons could be associated with both parton and beam remnants
- It leads to some distortions in  $\eta$  resolution

# Clustered jet vs parton (R=0.4)






- There is no problem in  $\eta$  resolution for clustered jets
- Clustered jets include only particles in algorithm radius

# Clustered jet vs parton (R=0.6)





# Clustered jet vs parton (R=0.8)





# Mean values and sigma

### “True” jet vs parton




### Clustered jet vs parton



- High  $p_T$  partons produce jets, which could be better associated with them

# Process $qg \rightarrow q\gamma$ cross section



- Expected instantaneous luminosity -  $10^{32} \text{ cm}^{-2} \text{ s}^{-1}$
- For 100 days of work integral luminosity  $\sim 10^5 \text{ nb}^{-1}$
- We have enough statistics even for high  $p_T$  partons



# Conclusion and plans



- Today's situation:
  - Kinematical properties of hard scattered partons, "true" jets and clustered jets was compared
  - It seems that clustered jets could be associated with initial parton
  - With increasing of parton  $p_T$  we get better association between clustered jets and initial parton
  - We expect enough statistics to make these analysis
- Plans:
  - Choose the best clustering algorithm, i.e. *anti- $k_t$* ,  *$k_t$* , *Cambridge/Aache* and *Iterative Cone* algorithm
  - Find cuts, which select jets associated with initial parton
    - Minimum particle  $p_T$
    - Minimum jet  $p_T$  and  $\eta$
    - Size of jet cone
    - Specific channel cuts (e.g. opposite photon)

Participant	FTE
<u>D. Budkouski</u>	0.4
I. Lapushanskii	0.4
A. Tumasyan	0.2




Expert: S.Shmatov

# Mean values and sigma




## Clustered jet vs "true" jet



# Clustered jet vs "true" jet (R=0.4)



# Clustered jet vs "true" jet (R=0.6)



# Clustered jet vs "true" jet (R=0.8)

