Contribution ID: 17

Type: not specified

How to access QED at a supercritical Coulomb field with heavy ions

Tuesday 5 September 2023 12:40 (30 minutes)

It is known that in slow collisions of two bare nuclei with the total charge number larger than the critical value, $Z_1+Z_2 > Z_c = 173$, the initially neutral vacuum can spontaneously decay into the charged vacuum and two positrons. Detection of the spontaneous emission of positrons would be the direct evidence of this fundamental phenomenon. However, the spontaneous positron emission is generally masked by the dynamical positron emission, which is induced by a rapidly changing electric field created by the colliding nuclei. For many years it was believed that the vacuum decay can be observed only in collisions with nuclear sticking, when the nuclei are bound for some period of time due to nuclear forces. But to date there is no evidence for the nuclear sticking in such heavy-ion collisions. In our recent papers [I.A. Maltsev et al., PRL, 2019; R.V. Popov et al., PRD, 2020; R.V. Popov et al., PRD, 2023], it was shown that the vacuum decay can be observed without any sticking of the nuclei. This can be done via measurements of the pair-production probabilities or the positron spectra for a given set of nuclear trajectories. The results of this study will be presented in the talk.

The author acknowledges the support by RSCF grant No. 22-62-00004.

Primary author: Prof. SHABAEV, Vladimir (St. Petersburg State University/ NRC "Kurchatov Institute"-PNPI)

Presenter: Prof. SHABAEV, Vladimir (St. Petersburg State University/ NRC "Kurchatov Institute"- PNPI)

Session Classification: Plenary