


Supported in part by



# Global polarization of $\Lambda$ and $\Xi$ hyperons in Au+Au collisions in the STAR experiment

## Outline:

- Introduction
- Global hyperon polarization
- Motivation
- The STAR experiment
- Hyperon polarization measurements
- Results
- Conclusions


Egor Alpatov (for the STAR Collaboration)

National Research Nuclear University MEPhI


# Introduction



- The Quark-Gluon Plasma (QGP) formed in non-central nucleus-nucleus collisions is associated with large angular momentum, that leads to vorticity in the medium
- Spin-orbit coupling aligns spin directions of produced particles along the direction of vorticity
  - Z.-T. Liang and X.-N. Wang, PRL94, 102301 (2005)
  - S. A. Voloshin, arXiv:nucl-th/0410089
- Another possible source of particle polarization is magnetic field, created in non-central collisions in the initial stage
  - D. Kharzeev, L. McLerran, and H. Warringa, Nucl.Phys.A803, 227 (2008)
  - McLerran and Skokov, Nucl. Phys. A929, 184 (2014)



# Vorticity



- In non-central Heavy-Ion Collisions the initial collective longitudinal flow velocity depends on the velocity gradient:



$$\omega_y = \frac{1}{2} (\nabla \times v)_y \approx -\frac{1}{2} \frac{dv_z}{dx}$$

- For small polarization:

Becattini, Karpenko, Lisa, Upsal, Voloshin PRC95.054902 (2017)

$$P_\Lambda \approx \frac{1}{2} \frac{\omega}{T} + \frac{\mu_\Lambda B}{T}$$

$$P_{\bar{\Lambda}} \approx \frac{1}{2} \frac{\omega}{T} - \frac{\mu_\Lambda B}{T}$$



# How to measure global polarization?


- Hyperons are “self-analyzing” due to weak decay properties:
  - Daughter baryons are preferentially emitted along parent spin direction
- Daughter baryons of hyperons with polarization ( $\vec{P}$ ) follows the distribution:

$$\frac{dN}{d\Omega^*} = \frac{1}{4\pi} (1 + \alpha_H |\vec{P}| \cdot \widehat{\vec{p}}_b^*) = \frac{1}{4\pi} (1 + \alpha_H P \cos \theta^*)$$

- $\alpha_H$  - decay parameter, unique for each hyperon species
- $\widehat{\vec{p}}_b^*$  is the daughter baryon momentum in the parent frame
- Projection to the transverse plane can be measured:

$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\psi_1 - \varphi_p^*) \rangle}{Res(\psi_1)}$$


- $\psi_1$  is first-order event plane angle (proxy for reaction plane)
- $\psi_1$  and its resolution  $Res(\psi_1)$  can be calculated with spectator’s signal.
- $\Xi$  global polarization could also be measured via its daughter  $\Lambda$  polarization with transfer factor  $C_{\Xi\Lambda} = 0.944$




STAR *Nature* 548, 62–65 (2017)

# Motivation

- Global polarization of  $\Lambda$  hyperons was measured for  $\sqrt{s_{NN}} = 3-200$  GeV at STAR
- $P_H$  decreases with increasing collision energy
- Theoretical calculations can quantitatively explain the energy dependence of the  $\Lambda$  polarization, but many of them fail to explain differential measurements
- Nowadays there is a growing interest to measure the global polarization of other hyperons such as  $\Xi$ .
- $\Xi$  and  $\Omega$  hyperons global polarization was measured in Au+Au collisions at  $\sqrt{s_{NN}} = 200$  GeV
- $\Xi$  polarization may provide new input for global polarization and vorticity studies




PRC 108, 014910 (2023)



# The STAR experiment



iTPC      TPC      TOF      EPD



## Detectors with their $\eta$ acceptance:

### Hyperon reconstruction:

- Time Projection Chamber  
 $|\eta| \in [-1, 1]$
- Time-Of-Flight  
 $|\eta| \in [-0.9, 0.9]$

### Event plane angle measurement:

- Beam-Beam Counter  
 $|\eta| \in [3.3, 5.0]$
- Event-Plane Detector  
 $|\eta| \in [2.1, 5.1]$

# Experimental technique




- Event plane  $\Psi_1$  is determined by detectors at forward rapidity where directed flow is large

$$\Psi_1 = \tan^{-1} \left( \frac{\sum w_i \sin(\phi_i)}{\sum w_i \cos(\phi_i)} \right), \text{ where } w_i \text{ is detector's tile ADC}$$



A. M. Poskanzer, S. A. Voloshin, PRC58.1671(1998)

- $Res(\Psi_1, \text{Full } \eta) = \sqrt{2 \langle \cos(\Psi_{1, \text{Forward } \eta} - \Psi_{1, \text{Backward } \eta}) \rangle}$
- EPD was used to determine event-plane angle (BBC for systematics)



**Hyperon reconstruction** performed via decay topology with KFParticle technique


- $\Lambda$  daughters identified via TPC and TOF
- $\Xi$  were reconstructed via  $\Xi \rightarrow \Lambda + \pi$




$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\psi_1 - \varphi_p^*) \rangle}{Res(\psi_1)}$$

# $\Xi$ global polarization: $\sqrt{s_{NN}}=19.6$ GeV



Centrality dependence



$\eta$  dependence







$p_T$  dependence



- Direct polarization measurements are consistent with measurements via daughter decays
- Polarization increases with centrality
- No obvious pseudorapidity or  $p_T$  dependence




# $\Xi$ global polarization: $\sqrt{s_{NN}}=27$ GeV




- Direct polarization measurements are consistent with measurements via daughter decays
- Polarization increases with centrality
- Weak pseudorapidity or  $p_T$  dependence if any

# $\Xi$ global polarization results



## Global hyperon polarization $\sqrt{s_{NN}}$ dependence

- Observed positive  $\Xi$  global polarization
- Experimental results for  $\Xi$  global polarization consistent with each other with given large uncertainties at different collision energies
- Both  $\Xi$  global polarization measurement methods are consistent within uncertainties
- $\Xi$  and inclusive  $\Lambda$  global polarization are consistent within statistical uncertainties
- Global polarization of  $\Xi$  hyperons consistent with AMPT predictions



# Conclusions



- We presented  $\Xi + \bar{\Xi}$  global polarization measurements in Au+Au collisions at  $\sqrt{s_{NN}} = 19.6$  & 27 GeV, which helps to understand QCD spin dynamics and vorticity of QGP medium
  - This information can be used in theoretical development
- $\Xi + \bar{\Xi}$  global polarization is comparable with  $\Lambda + \bar{\Lambda}$  global polarization within uncertainties, indicating a global nature of polarization

Thank you for your attention!