




Direct search for low mass dark matter with DarkSide-50


*Sergey Chashin (SINP MSU)
on behalf of the DarkSide collaboration*

21st Lomonosov Conference, 25th August 2023, Moscow, Russia

The DarkSide-50



Dual-Phase Argon Time Projection Chamber




Liquid Scintillator Veto


Water Cherenkov Veto

Dual-Phase Argon Time Projection Chamber (TPC)

High energy event



Low energy event




- Recoil energy \rightarrow scintillation photons and ionization e^-
- Amplitude of S1+S2 \rightarrow calorimetry
- Particle identification via pulse shape discrimination (PSD)
- Drift time (between S1 and S2) \rightarrow Z coordinate

- Amplification in GAr lets us detect signals with high efficiency above photoelectronic noise \rightarrow lower energy threshold
- PSD, and drift time are not available


Detection channels: elastic scattering

Nuclear Recoil (NR)



DM high-mass range: $\sim 5 \text{ GeV}/c^2$ to $10 \text{ TeV}/c^2$

Electron Recoil (ER)




DM low-mass range: $\sim 30 \text{ MeV}/c^2$ to $5 \text{ GeV}/c^2$

Detection channels: inelastic scattering

Electron shell follows the recoiling nucleus with delay, so the atom after a DM-nucleus interaction can become polarized, which can lead to the following effects:

Migdal Effect:
keV-range electron emission
X-ray photon emission



Bremsstrahlung:
X-ray photon emission
(weak effect, low sensitivity)

DM low-mass range: $\sim 30 \text{ MeV}/c^2$ to $5 \text{ GeV}/c^2$

(Images credit: XENON1T collaboration)

DarkSide-50 dataset


The dataset consists of 653.1 live-days (12 ton-days) of underground argon data, taken from December 12, 2015, to February 24, 2018, with an average trigger rate of 1.54 Hz

Detector showed reasonable stability for the whole period of 26 months:

- $\delta T = \pm 0.02$ K, $\delta P < \pm 0.005$ psi,
- $\delta(S1) \sim 0.4\%$, $\delta(S2) < 1\%$


Multiple-pulses & pileup events
 Improperly triggered events
 & SE events (delayed electron signals)
 Surface- α events

Signal efficiency is $>95\%$ for the region of interest (low energy analysis threshold is equal to 4 Ne)



DarkSide-50 background model

- Internal background consists of ^{39}Ar and ^{85}Kr
- External background consists of impurities in PMT and cryostat materials
- Spurious electrons (SE), that follow large S2 pulses




Phys. Rev. D 107, 063001 (2023)

Standard low mass wimp search



S1+S2 analysis, search for NR with PSD

Approaches used to improve sensitivity in low mass region:

- Extended exposure
- Improved data selection criteria
- More accurate detector calibration
- Better background modeling



- DS50 2022
- PandaX-4T 2022
- LUX 2021
- DAMIC 2020
- Xenon1T 2020
- Cresst-III 2019
- Pico-60 2019
- Xenon1T Migdal 2019
- DS50 2018
- CDMSlite 2017
- PICASSO 2017
- CDMS 2013
- Cogent 2013
- DAMA/LIBRA 2008
- LAr Neutrino Floor




Phys. Rev. D 107, 063001 (2023)

Sergey Chashin, SINP MSU

Standard low mass wimp search


This approach allowed us to set the most stringent exclusion limit at $M_\chi = [1.2, 3.6] \text{ GeV}/c^2$



Phys. Rev. D 107, 063001 (2023)

S2-only low mass wimp search

- Using only S2 signal in analysis allows to increase sensitivity to low-energy interactions (which is crucial for low mass DM search), but PSD and Z-coordinate reconstruction become unavailable
- S2 signals, amplified in GAr, allow to identify even the single ionization electron




Phys. Rev. Lett. 121, 081307 (2018)

DM-electron scattering



This mechanism describes interactions between several DM candidates and bound electrons of a target atom:

- Fermion or scalar boson light DM (LDM) interact via a vector mediator \rightarrow ionization
- Pseudo-scalar DM (Axion-Like Particles) or vector boson DM (Dark Photons) are absorbed by argon shell electrons \rightarrow monoenergetic signal at the particle's rest mass
- Sterile neutrinos inelastically scatter of bound electrons \rightarrow ionization



Phys. Rev. Lett. 130, 101002 (2023)



DM-electron scattering



Phys. Rev. Lett. 130, 101002 (2023)


The most stringent exclusion limit on DM-electron interaction cross section was set in the mass region of [16, 56] MeV/c² for a heavy mediator and above 80 MeV/c² for a light mediator

DM-electron scattering




For different DM candidates the exclusion limit was calculated for model parameters:

- Axion-electron coupling strength g_{Ae}
- Dark photon-photon kinetic mixing strength κ
- Sterile neutrino mixing angle $|U_{e4}|^2$





Migdal effect and bremsstrahlung

- Migdal effect (ME): delayed movement of the electron shells after the recoiling nucleus \rightarrow polarization of the atom \rightarrow ionization and photon emission
- Bremsstrahlung: accelerated movement of the recoiling nucleus in the electric field of its electron shells \rightarrow photon emission (weak effect)



JHEP 11 (2020) 034 (with old dataset)


Migdal effect



Phys. Rev. Lett. 130, 101001 (2023)

Migdal effect

- This approach allowed us to set the most stringent exclusion limit at $M_\chi = [0.04, 3.6] \text{ GeV}/c^2$
- The limit is entirely driven by ME up to $0.5 \text{ GeV}/c^2$; also, the limit in this mass region is not affected by choice of quenching fluctuation model



Phys. Rev. Lett. 130, 101001 (2023)

Conclusions

- Detectors with dual-phase argon TPCs, such as DarkSide-50, are able to significantly increase the dark matter search capabilities in low mass region
- Advanced analysis methods and implementation of atomic effects, such as the Migdal effect, in the analysis can furthermore increase sensitivity of low mass dark matter search
- Increasing of exposure is crucial for low mass dark matter search, which will be achieved by experiments with much greater target mass, such as DarkSide-20k and DarkSide-LowMass – the next stages of the DarkSide program