

COHERENT ELASTIC NEUTRINO-NUCLEUS SCATTERING SEARCH IN THE ν GeN EXPERIMENT.

*D.V. Ponomarev^{a,b,1}, A.D. Bystryakov^{a,b}, A. Konovalov^b, A.V. Lubashevskiy^{a,b},
on behalf of the ν GeN Collaboration*

^a Dzhelapov Laboratory of Nuclear Problems, Joint Institute for Nuclear
Research, 6 Joliot-Curie, 141980, Dubna, Russia

^b Federal State Budgetary Science Institution — the Lebedev Physical Institute of
the Russian Academy of Sciences, 53 Leninskiy Prospect, 119991, Moscow, Russia

The ν GeN experiment aims to search for coherent elastic neutrino-nucleus scattering and to study the other rare processes. The experimental setup is located about 12 meters from the center of the 3.1 GWth reactor №3 of Kalinin NPP providing the antineutrino flux of $(3.6-4.4)\cdot 10^{13}$ particles \cdot cm $^{-2}\cdot$ s $^{-1}$. The intense antineutrino flux gives a possibility to detect coherent elastic scattering of reactor antineutrinos on Ge nuclei in the regime of full coherence, as well as to study other properties of neutrinos

PACS: 25.30.Pt; 29.40.Wk

Introduction

The coherent elastic neutrino-nucleus scattering (CE ν NS) is the process which was predicted theoretically 50 years ago [1]. The coherency effect takes place when $QR \ll 1$, where R is a nuclear radius and Q is a transferred momentum. CE ν NS was discovered in 2017 (but not in the region of complete coherence) by the COHERENT experiment [2]. This process is extremely difficult to detect, due to very low energy transferred by neutrino to a nucleus. Depending on the nucleus and neutrino kinetic energy, this energy can be either a few keV or even less than keV. Due to quenching, only a minor part of the transferred energy contributes to ionization. This makes detecting CE ν NS challenging, necessitating the use of low background detectors with extremely low thresholds and a potent neutrino source. An accurate measurement of CE ν NS can be a key to the physics beyond the Standard Model. The anomalous neutrino magnetic moment, higher than 10^{-15} μ_B will also indicate the existence of a new physics [3].

¹E-mail: ponom@jinr.ru

Experiment

20 The ν GeN experiment was deployed at the Kalinin nuclear power plant
 21 under reactor №3, which has a power of 3.1 GW_{th} [4]. The distance to the
 22 center of the reactor core is about 11 meters from the floor of the experi-
 23 mental hall. In this hall it is possible to operate with antineutrino flux up
 24 to $4.4 \cdot 10^{13} \text{ particles} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$. The reactor and its building provide about 50
 25 meters of water equivalent shielding against cosmic radiation [5].

26 To detect $\text{CE}\nu\text{NS}$, a 1.4 kg HPGe low-threshold, low-background detec-
 27 tor is used in the ν GeN experiment. The detector is cooled to a cryogenic
 28 temperature by the cryocooler CP5+ [6]. The detector shielding against ra-
 29 dioactive background consist of 3D printed nylon as inner layer (antiradon
 30 shielding), 10 cm of oxygen-free high-purity copper, 8 cm of 3 % borated
 31 polyethylene, 10 cm of lead, another 8 cm of 3 % borated polyethylene and
 32 5 cm thick plastic muon veto. The nitrogen flushing of the internal volume
 33 of the shield allows to suppress potential radon-related backgrounds. The
 34 cryocooler is placed on antivibration platform TS-C30 [7] to reduce mechan-
 35 ical noises of the KNPP equipment. The experimental hall is equipped with
 36 air conditioners to provide a stable temperature. Scheme of the shielding is
 37 presented in figure 1.




Fig. 1. Scheme of the shielding of the ν GeN experiment

38 To suppress electronic noises the several amplifiers with different shap-
 39 ing times connected to the real time ADC are used. The energy scale is
 40 calibrated using a welding tungsten rod, cosmogenic isotopes of germanium,
 41 and a pulse generator. The energy resolution of the pulse generator signals is
 42 $101.6(5) \text{ eV}$ (FWHM). The detection efficiency for signals from events with
 43 energy higher than 250 eV is more than 80 %. The efficiency of the applied
 44 cuts, as determined by 10.37 keV cosmogenic line, is 85.3(19) %.

45 The whole setup is mounted on the special lifting mechanism (figure 2),
 46 which allows to change the distance between spectrometer and the center of

47 the reactor core from 11.1 to 12.2 meters. Thus, we can change the antineu-
 48 trino flux passing through the detector $(3.6\text{--}4.4)\cdot 10^{13}$ particles $\cdot\text{cm}^{-2}\cdot\text{s}^{-1}$.




Fig. 2. Scheme of the lifting mechanism.

Results

49
 50 Up until 2022, measurements were conducted at a distance of 11.84 meters
 51 from the reactor core. At this location, data corresponding to 94 kg \cdot days
 52 days with the reactor operational and 47 kg \cdot days days with the reactor non-
 53 operational were collected after implementing all cuts. The analysis focused
 54 on the energy range of 0.32–0.36 keV. No $\text{CE}\nu\text{NS}$ signal was detected, leading
 55 to the establishment of an upper limit on the quenching parameter $k < 0.26$ [4].

56 In the fall of 2022, upgrades were implemented that enhanced noise and
 57 background levels. Following these improvements, the spectrometer was re-
 58 located to a higher position (11.09 meters). In this new position, data from
 59 217 kg \cdot days with the reactor operational and 55 kg \cdot days with the reactor non-
 60 operational were analyzed. The sensitivity to $\text{CE}\nu\text{NS}$ was assessed using the
 61 reactor OFF spectrum (55 kg \cdot days) and Azimov data sets, which include
 62 generated $\text{CE}\nu\text{NS}$ spectra combined with the OFF spectra. The statistical
 63 analysis was based on the same dataset size as the current one (217 kg \cdot days).

64 In new data the region of interest 0.29–0.36 was chosen, thanks to a lower
 65 noise with respect to the previous measurements. The excess in the count
 66 rate during reactor ON was not observed (figure 3).




Fig. 3. The residual spectrum of the ν GeN experiment data collected in the upper position. Black dots with errors—experimental data. Red solid line—expected CE ν NS spectra, in case of the quenching parameter $k=0.23$.

67

Conclusion

68 Measurement with the ν GeN spectrometer is ongoing. More than 1200
 69 kg·days of data has been accumulated so far. The spectrometer demon-
 70 strates good performance. The background level during reactor ON and
 71 OFF regimes is stable. The combined analysis of data taken in the lower
 72 and upper positions will be presented soon, as well as results of the study
 73 of neutrino electromagnetic properties. It is planned to install new digitizer
 74 based DAQ and inner veto. This work has been partly supported by the
 75 Ministry of science and higher education of the Russian Federation.

76

REFERENCES

- 77 1. *Freedman D.Z.* Coherent effects of a weak neutral current // Phys. Rev.
 78 D. — 1974. — V. 9, no. 5. — P. 1389–1392.
- 79 2. *Akimov D. et al.* Observation of Coherent Elastic Neutrino-Nucleus Scat-
 80 tering // Science. — 2017. — V. 357, no. 6356. — P. 1123–1126.
- 81 3. *Abdullah M. et al.* Coherent elastic neutrino-nucleus scattering: Terrestrial
 82 and astrophysical applications [Электронный ресурс] // ArXiv e-prints. —
 83 2022.
- 84 4. *Alekseev I. et al.* First results of the ν GeN experiment on coherent elastic
 85 neutrino-nucleus scattering // Phys. Rev. D. — 2022. — V. 106, no. 5. —
 86 P. L051101.

- 87 5. *Alekseev I. et al.* DANSS: Detector of the reactor AntiNeutrino based on
88 Solid Scintillator // Journal of Instrumentation. — 2016. — V. 11, no. 11. —
89 P. 11011.
- 90 6. Cryo-Pulse 5 Plus Electrically Refrigerated Cryostat:
91 [https://www.mirion.com/products/technologies/spectroscopy-scientific-](https://www.mirion.com/products/technologies/spectroscopy-scientific-analysis/gamma-spectroscopy/detectors/hpge-cryostats-coolers-options/cryo-pulse-5-plus-electrically-refrigerated-cryostat)
92 [analysis/gamma-spectroscopy/detectors/hpge-cryostats-coolers-](https://www.mirion.com/products/technologies/spectroscopy-scientific-analysis/gamma-spectroscopy/detectors/hpge-cryostats-coolers-options/cryo-pulse-5-plus-electrically-refrigerated-cryostat)
93 [options/cryo-pulse-5-plus-electrically-refrigerated-cryostat.](https://www.mirion.com/products/technologies/spectroscopy-scientific-analysis/gamma-spectroscopy/detectors/hpge-cryostats-coolers-options/cryo-pulse-5-plus-electrically-refrigerated-cryostat)
- 94 7. Antivibration platform TS-C30:
95 [http://tablestable.com/en/products/view/45/.](http://tablestable.com/en/products/view/45/)