

# Effect of magnetic field on Dual QCD Quark-hadron phase transition

*Garima Punetha*<sup>1</sup>

LSM Campus Pithoragarh, SSJ University Almora Uttarakhand, India

In order to better understand the dynamics of the quark-hadron phase transition in the presence of a magnetic field, the thermal behaviour of the non-perturbative QCD vacuum has been examined. The dynamical configuration of the resulting dual QCD vacuum and its flux tube configuration have been explored in order to  
1 investigate the non-perturbative properties of QCD. Within the context of dual QCD-based hadronic bag, which ensures the critical parameters and the accompanying critical sites for quark-hadron phase transition, related thermodynamic quantities and equation of state (EoS) to define quark matter have also been stud-  
2 ied in presence of magnetic field.

3 PACS: 12.38.-t, 14.80.Hv, 64.60.av, 12.38.Mh

## 4 Introduction

5 High-energy collisions have been effective in producing quark-gluon plasma  
6 (QGP), a state of matter thought to have existed shortly after the big  
7 bang [1]. During off-center collisions, the charged ions can generate sub-  
8 stantial magnetic fields up to  $eB(1 - 15)m_\pi^2$  [2]. This method can produce  
9 magnetic fields that last just a brief duration yet are stationary during that  
10 time [3]. There has been a flurry of research in this area [4–7], and the  
11 theoretical methods used to explore QGP need to be modified to reflect the  
12 effects of external magnetic fields. Insights can be gained from measurements  
13 at the Large Hadron Collider and relativistic heavy ion collider energies [8, 9],  
14 which can be used to constrain theoretical modelling. The evolution of QGP  
15 is significantly influenced by the equation of state [10]. Therefore, it is cru-  
16 cial to investigate the behaviour of magnetised QGP [11]. Different methods  
17 have been used to investigate how magnetic fields affect QGP [12–14]. In  
18 the present paper with the help of Grand canonical partition function within  
19 the domain of Bag Model, we calculated normalised pressure for magnetized  
20 QGP at finite temperature.

## 21 Field Decomposition Formulation and Magnetic Symmetry

22 The mathematical foundation for the dual gauge theory comes from the  
23 observation that the non-Abelian gauge symmetry allow an extra internal

---

<sup>1</sup>E-mail: garimapunetha@gmail.com

24 symmetry called magnetic symmetry which restricts and reduces the dynam-  
25 ical degrees of the theory [15–18],

$$D_\mu \hat{m} = 0, \text{ i.e. } (\partial_\mu + g \mathbf{W}_\mu \times) \hat{m} = 0, \quad (1)$$

26 where  $\hat{m}$  is a multiplet defined by the adjoint representation of the group. The  
27 most general gauge potential which satisfies the above constraint is written  
28 as,

$$\mathbf{W}_\mu = A_\mu \hat{m} + A'_\mu \hat{m}' - g^{-1} (\hat{m} \times \partial_\mu \hat{m}) - g^{-1} (\hat{m}' \times \partial_\mu \hat{m}'), \quad (2)$$

29 where,  $A_\mu$  and  $A'_\mu$  are the Abelian component of  $\mathbf{W}_\mu$  along  $\hat{m}$  and  $\hat{m}'$  respec-  
30 tively and are unrestricted by the constraint. The topological structure may  
31 be brought into dynamics in a dual symmetric way by imposing magnetic  
32 symmetry and for  $SU(3)$  the relevant homotopy group is  $\pi_2[SU(3)/U(1) \otimes$   
33  $U'(1)]$  so that  $\pi_2[SU(3)/U(1) \otimes U'(1)] = \pi_1[U(1) \otimes U'(1)]$ , where  $U(1) \otimes U'(1)$   
34 is the two Abelian subgroups generated by  $\lambda_3$  and  $\lambda_8$ . The  $\hat{m}$  consist of  $n$   
35 winding of the i-spin subgroup followed by  $n'$  winding of u-spin subgroup of  
36  $SU(3)$  and is given as,

$$\hat{m} = \begin{pmatrix} \sin \alpha \cos \frac{1}{2} \alpha \cos(\beta - \beta') \\ \sin \alpha \cos \frac{1}{2} \alpha \sin(\beta - \beta') \\ \frac{1}{4} \cos \alpha (3 + \cos \alpha) \\ \sin \alpha \sin \frac{1}{2} \alpha \cos \beta \\ \sin \alpha \sin \frac{1}{2} \alpha \sin \beta \\ \frac{-1}{2} \sin \alpha \cos \alpha \cos \beta' \\ \frac{-1}{2} \sin \alpha \cos \alpha \sin \beta' \\ \frac{1}{4} \sqrt{3} \cos \alpha (1 - \cos \alpha) \end{pmatrix}, \quad (3)$$

37 where  $\alpha$  and  $\beta$  are coset variables. Rotating the magnetic vector  $\hat{m}$  to a fix  
38 time independent direction by a gauge transformation leads to the value of  
39 gauge potential as,

$$\mathbf{W}_\mu \xrightarrow{U} g^{-1} \left[ \left( (\partial_\mu \beta - \frac{1}{2} \partial_\mu \beta') \cos \alpha \right) \hat{\xi}_3 + \frac{1}{2} \sqrt{3} (\partial_\mu \beta' \cos \alpha) \hat{\xi}_8 \right], \quad (4)$$

40 where  $\hat{\xi}_3$  and  $\hat{\xi}_8$  are the space-time independent unit vectors pointing to  $\lambda_3$   
41 and  $\lambda_8$  directions of the internal space respectively and the associated field  
42 strength takes the form as,

$$\mathbf{G}_{\mu\nu} \xrightarrow{U} (F_{\mu\nu} + B_{\mu\nu}^{(d)}) \hat{\xi}_3 + (F'_{\mu\nu} + B_{\mu\nu}'^{(d)}) \hat{\xi}_8, \quad (5)$$

43 where  $F_{\mu\nu}$  and  $B_{\mu\nu}^{(d)}$  are the electric and magnetic field strength respectively.  
44 The confinement mechanism of the QCD vacuum can be understood in ab-  
45 sence of color electric sources (quarks) and the Lagrangian may be reduced  
46 in the following form,

$$\mathcal{L}_d^{(m)} = -\frac{1}{4} B_{\mu\nu}^2 - \frac{1}{4} B_{\mu\nu}'^2 + |(\partial_\mu + i \frac{4\pi}{g} B_\mu^{(d)}) \phi|^2 + |(\partial_\mu + i \frac{4\pi \sqrt{3}}{g} B_\mu'^{(d)}) \phi'|^2 - V, \quad (6)$$

$\gamma$	$\alpha_s$	$\bar{m}_\phi(GeV)$	$\bar{m}_B(GeV)$	$\lambda_{QCD}^{(d)}(GeV^{-1})$	$\xi_{QCD}^{(d)}(GeV^{-1})$	$\kappa_{QCD}^{(d)}$
5.617	0.25	1.21	1.75	0.57	0.83	0.69
6.828	0.24	1.69	1.62	0.61	0.59	0.99
8.093	0.23	2.17	1.52	0.65	0.46	1.42
9.833	0.22	2.90	1.41	0.70	0.34	2.05

Table 1. The masses of vector and scalar glueball as functions of  $\alpha_s$ , where  $\lambda_{QCD}^{(d)}$ ,  $\xi_{QCD}^{(d)}$  and  $\kappa_{QCD}^{(d)}$  are the penetration depth, coherence length and Ginzburg-Landau parameter respectively.

where,  $V$  is the effective potential responsible for the dynamical breaking of magnetic symmetry and  $\phi$  is a complex scalar field for monopoles is given below,

$$V = \frac{48\pi^2}{g^4} \lambda(\phi^* \phi - \phi_0^2)^2 + \frac{432\pi^2}{g^4} \lambda'(\phi^* \phi' - \phi_0'^2)^2. \quad (7)$$

Utilizing the asymptotic solutions, the energy per unit length of the resulting flux tube configuration is obtained as,

$$k_{(n,n')} = 2\pi \int_0^\infty \rho d\rho \left[ \frac{n^2 g^2}{32\pi^2 \rho^2} \left( \frac{dF}{d\rho} \right)^2 + \frac{n^2}{\rho^2} F^2(\rho) \chi^2(\rho) + \left( \frac{d\chi}{d\rho} \right)^2 + 3\lambda \alpha_s^{-2} (\chi^2 - \phi_0^2)^2 \right] + 2\pi \int_0^\infty \rho d\rho \left[ \frac{n'^2 g^2}{96\pi^2 \rho^2} \left( \frac{dG}{d\rho} \right)^2 + \frac{n'^2}{\rho^2} G^2(\rho) \chi'^2(\rho) + \left( \frac{d\chi'}{d\rho} \right)^2 + 27\lambda' \alpha_s^{-2} (\chi'^2 - \phi_0'^2)^2 \right], \quad (8)$$

where  $n, n'$  are color quantum numbers and  $F(\rho) \xrightarrow{\rho \rightarrow \infty} C\sqrt{\rho} \exp(-m_B \rho)$ ,  $G(\rho) \xrightarrow{\rho \rightarrow \infty} C'\sqrt{\rho} \exp(-m'_B \rho)$  and  $\phi(x) = \exp(in\varphi)\chi(\rho)$ ,  $\phi'(x) = \exp(in'\varphi)\chi'(\rho)$ . Incorporating color reflection invariance the masses of the magnetic glueballs is estimated by evaluating the string tension  $k_{(n,n')}$  of the resulting flux tube written as,  $k_{(n,n')} = \frac{1}{2\pi\alpha'} = \gamma_{(n,n')} \phi_0^2$ . As a result, using the numerical computation of equation (8), for  $\gamma_{(n,n')}$ , one may obtain the vector ( $\bar{m}_B$ ) and scalar ( $\bar{m}_\phi$ ) magnetic glueball masses as a function of  $\alpha_s$  and these results are obtained in table (1).

### Equation of State in presence of Magnetic Field

The transition from a confined hadronic phase to a QGP phase, is foreseen at high temperatures and/or baryonic chemical potentials and for some decades, the bag model equation of state (EoS) has been accustomed to characterize QGP. Employing the dual QCD formalism, the bag energy ( $B_{su3}$ )

may be identified and manifest to perform an significant role in determining the phase structure of the QCD vacuum. The dominant part of the energy related with the confinement regime is explicated from equation (8) and may be recognized as bag energy expressed as,

$$B_{su3}^{1/4} = \left( \frac{\lambda}{\pi^2} \right)^{1/4} \frac{\bar{m}_B}{4}, \quad (9)$$

where  $\bar{m}_B$  is the non-thermal contribution to the vector glueball mass. To contemplate the QGP phase, we utilize a Bag model EoS with bag energy. We have chosen the limit of large  $B$  in order to make the numerical evaluation of thermal integrals less complicated. The pressure for the QGP phase in presence of magnetic field may be expressed in the following form [19],

$$P_{QGP} = \frac{37}{90}\pi^2 T^4 + T \sum_f \frac{q_f B}{2\pi^2} \int dk_z \ln \left[ 1 + e^{-\frac{\sqrt{k_z^2 + m_f^2}}{T}} \right] - B_{su3}, \quad (10)$$

where  $B$  is the external magnetic field in the z-direction,  $f$  is the flavor index and  $q_f$  is the electric charge. The pressure for hadronic phase may be expressed as [18],

$$P_h = \frac{7}{180}\pi^2 T^4. \quad (11)$$

Utilizing the grand canonical ensemble formalism, the EoS for quark-hadron system at non-vanishing external magnetic field has been constructed within the framework of  $SU(3)$  dual QCD hadronic bag in which the bag pressure has been shown to depend on the vector glueball mass mode of the magnetically condensed QCD vacuum. In fig. 1 3-D plot of normalised pressure for hadronic and QGP phase given by equations (10) and (11) respectively and has been plotted in the  $T - eB$  plane for different couplings.

## Conclusion

The effect of magnetic field ( $B$ ) on the equation of state of hot quark matter is investigated. A magnetic field induces phase-space Landau-level quantization and for lowest Landau-level ( $l = 0$ ) normalised pressure for QGP and hadronic phase have been obtained within the framework of dual QCD hadronic bag at non-zero external magnetic field. With increasing magnetic field strength the critical temperature reduces which leads to inverse magnetic catalysis. The normalised pressure increases with increasing magnetic field especially at low temperatures. At high temperature the normalised pressure is not affected by magnetic field and is apparently limited to the Stefan Boltzmann limits.

## ACKNOWLEDGEMENT

The author is grateful to the Uttarakhand State Council for Science and Technology (UCOST), Dehradun (India) for the financial assistance in terms of the research project for the study.




Fig. 1. The 3-D plot of normalized pressure (hadron phase and QGP phase) in  $T - eB$  plane for different couplings.

- 99 1. B. B. Back et al., “The PHOBOS Perspective on Discoveries at RHIC,”  
 100 Nucl.Phys. A **757**, 28 (2005) doi.org/10.1016/j.nuclphysa.2005.03.084  
 101 [arXiv:nucl-ex/0410022]; J. Adams et al., “Experimental and Theoretical  
 102 Challenges in the Search for the Quark Gluon Plasma: The STAR  
 103 Collaboration’s Critical Assessment of the Evidence from RHIC Col-  
 104 lisions,” *ibid.* **757**, 102 (2005) doi.org/10.1016/j.nuclphysa.2005.03.085  
 105 [arXiv:nucl-ex/0501009]; K. Adcox et al., “Formation of dense partonic  
 106 matter in relativistic nucleus-nucleus collisions at RHIC: Experimental  
 107 evaluation by the PHENIX collaboration,” *ibid.* **757**, 184 (2005)  
 108 doi.org/10.1016/j.nuclphysa.2005.03.086 [arXiv:nucl-ex/0410003]; I. Arse-  
 109 neet et al., “Quark Gluon Plasma and Color Glass Condensate at RHIC?  
 110 The perspective from the BRAHMS experiment,” *ibid.* **757**, 1 (2005)  
 111 doi.org/10.1016/j.nuclphysa.2005.02.130 [arXiv:nucl-ex/0410020].
- 112 2. V.V.Skokov, A. Y. Illarionov and V. D. Toneev, “Estimate of the mag-  
 113 netic field strength in heavy-ion collisions,” IJMPA **24**, 5925 (2009)  
 114 doi.org/10.1142/S0217751X09047570
- 115 3. K. Tuchin, “Electromagnetic radiation by quark-gluon plasma in magnetic  
 116 field,” Phys. Rev. C **83**, 017901 (2011) doi: 10.1103/PhysRevC.87.024912  
 117 [ arXiv:1206.0485 [hep-ph]]
- 118 4. B. Karmakar, R. Ghosh, A. Bandyopadhyay, N. Haque, and M. G.  
 119 Mustafa, “Anisotropic pressure of deconfined QCD matter in presence of  
 120 strong magnetic field within one-loop approximation,” Phys. Rev. D **99**,

- 094002 (2019) doi.org/10.1103/PhysRevD.99.094002 [ arXiv:1902.02607 [hep-ph]]
5. A. Bandyopadhyay, C. A. Islam, and M. G. Mustafa, “Electromagnetic spectral properties and Debye screening of a strongly magnetized hot medium,” Phys. Rev. D **94**, 114034 (2016) doi.org/10.1103/PhysRevD.94.114034 [arXiv:1602.06769 [hep-ph]]
6. L. Levkova and C. DeTar, “Quark-gluon plasma in an external magnetic field,” Phys. Rev. Lett. **112**, 012002 (2014) doi.org/10.1103/PhysRevLett.112.012002 [arXiv:1309.1142 [hep-lat]]
7. A. Bandyopadhyay, R. L. S. Farias, B. S. Lopes, and R. O. Ramos, “Quantum chromodynamics axion in a hot and magnetized medium,” Phys. Rev. D **100**, 076021 (2019) doi.org/10.1103/PhysRevD.100.076021 [arXiv:1906.09250 [hep-ph]]
8. S. Acharya et al., “Probing the Effects of Strong Electromagnetic Fields with Charge-Dependent Directed Flow in Pb-Pb Collisions at the LHC,” Phys. Rev. Lett. **125**, 022301 (2020) doi.org/10.1103/PhysRevLett.125.022301
9. J. Adam et al., “First Observation of the Directed Flow of  $D^0$  and  $\bar{D}^0$  in Au + Au collision at  $\sqrt{S_{NN}} = 200$  GeV,” Phys. Rev. Lett. **123**, 162301 (2019) <https://doi.org/10.1103/PhysRevLett.123.162301> [ arXiv:1905.02052 [nucl-ex]]
10. S. Bhadury, M. Kurian, V. Chandra, and A. Jaiswal, “First order dissipative hydrodynamics and viscous corrections to the entropy four-current from an effective covariant kinetic theory,” J. Phys. G: Nucl. Part. Phys. **47**, 085108 (2020) doi 10.1088/1361-6471/ab907b [arXiv:1902.05285 [hep-ph]]
11. S. S. Avancini, V. Dexheimer, R. L. S. Farias, and V. S. Timteo, “Anisotropy in the equation of state of strongly magnetized quark matter within the Nambu–Jona-Lasinio model,” Phys. Rev. C **97**, 035207 (2018) doi:10.1103/PhysRevC.97.035207
12. M. Kurian and V. Chandra, “Effective description of hot QCD medium in strong magnetic field and longitudinal conductivity,” Phys. Rev. D **96**, 114026 (2017) doi.org/10.1103/PhysRevD.96.114026 [arXiv:1709.08320 [nucl-th]]
13. S. Koothottil and V. M. Bannur, “Thermodynamic Behaviour of Magnetized QGP within the Self-Consistent Quasiparticle Model,” Phys. Rev. C **99**, 035210 (2019) doi: 10.1103/PhysRevC.99.035210 [ arXiv:1811.05377 [nucl-th]]

- 159 14. R. L. S. Farias, V. S. Timteo, S. S. Avancini, M. B. Pinto and G.  
160 Krein, “Thermo-magnetic effects in quark matter: Nambu-Jona-Lasinio  
161 model constrained by lattice QCD” Eur. Phys. J. A **53**, 101 (2017) doi:  
162 10.1140/epja/i2017-12320-8 [arXiv:1603.03847 [hep-ph]]
- 163 15. Y. M. Cho, “Restricted gauge theory,” Phys. Rev. D **21** (1980) 1080  
164 doi.org/10.1103/PhysRevD.21.1080
- 165 16. Y. M. Cho, “Extended gauge theory and its mass spectrum” Phys. Rev.  
166 D **23** (1981) 2415 doi.org/10.1103/PhysRevD.23.2415
- 167 17. Y. M. Cho, “Glueball Spectrum in Extended Quantum Chromodynam-  
168 ics,” Phys. Rev. Lett. **46** (1981) 302 doi.org/10.1103/PhysRevLett.46.302
- 169 18. Garima Punetha, Aritra Bandhyopadhyay and Shuchi Bisht, “Dual QCD  
170 thermodynamics at finite temperature and chemical potential, ” IJMPA  
171 **37** (2022) doi.org/10.1142/S0217751X22501755 2250175.
- 172 19. Eduardo S. Fraga, Leticia F. Palhares, “Deconfinement in the presence of  
173 a strong magnetic background: an exercise within the MIT bag model,”  
174 Phys. Rev. D **86**, 016008 (2012) doi.org/10.1103/PhysRevD.86.016008  
175 [arXiv:1201.5881 [hep-ph]].