

Open charm production at low energies

Short review and some ideas worth trying at SPD conditions

Evgeny Leshchenko, Aleksandr Berezhnoy

October 9, 2023

The review on charm production in hadronic interaction at low and middle energies

By now, we have a draft of the review in preparation to be published in the Bulletin of Moscow University. This draft contains dozens of pages and more than 100 references. In this study we itemize the experiments on hadronic charm production, as well as the theoretical approaches to the description of the charm production. In this report the content of the review will not be described in details, but the main theoretical approaches to the problem will be briefly highlighted and also some proposals for studying open charm at SPD will be put forward.

Open charm production at high energies

Main approaches:

- Gluon fusion

$$gg \rightarrow c\bar{c}$$

- Quark-antiquark annihilation

$$q\bar{q} \rightarrow c\bar{c}$$

- Charm excitation

$$gc \rightarrow gc$$

GM-VFNS **General-mass variable-flavor-number scheme**

(B. A. Kniehl, G. Kramer, I. Schienbein and H. Spiesberger)

The heavy quark is treated as any other massless parton, the mass is taken into account as large logarithms $\ln(p_T/m)$ in parton distribution and fragmentation functions, where they are resummed by imposing the DGLAP evolution [1].

FONLL **Fixed-order Next-to-Leading-Logarithm**

(M. Cacciari, M. Greco and P. Nason)

NLO (massive quark) + resummation of large logarithms.

At $p_T < 5m_c$ NLO works without logarithm resummation (see eq. 6.1 in [2]).

k_T fact. LO (massive quark) + virtual initial gluons

(It seems that sea c quark is not needed) [3]

Widely applied by S. P. Baranov, A. V. Lipatov, N. P. Zotov and V. A. Saleev [3, 4, 5].

Interactive cross section estimation within FONLL:

<http://www.lpthe.jussieu.fr/~cacciari/fonll/fonllform.html>

Fragmentation mechanism

The isolated quark cannot be observed and therefore must be hadronized:

$$c(\mathbf{p}) \xrightarrow{D(z)dz} H_c(z\mathbf{p})$$

$$\frac{d\sigma_{H_c}}{dz} = D(z) \cdot \sigma_{c\bar{c}}$$

$D(z)dz$ — the process-independent probability for the charm quark to transform into the open charm hadron H_c .

$$\frac{d\sigma_{H_c}}{dp_T} = \int_{2p_T/\sqrt{s}}^1 \frac{d\sigma_{c\bar{c}}}{dk_T} \left(\frac{p_T}{z} \right) \frac{D(z)}{z} dz$$
$$0 < z < 1$$

Works fairly good at $|\vec{p}| \rightarrow \infty$. Difficulties at $|\vec{p}| \sim m_c$:





- $m_Q \neq m_H$ (The invariant mass can not be conserved).
- The predictions depend essentially on the fragmentation variable choose ($|\vec{p}|, E, p_+$).
- The predictions depend on the coordinate system.

pQCD motivated FF

FF for B_c can be calculated.

$$\frac{M_{B_c}^2}{s_{e^+e^-}} \rightarrow 0$$

$$\frac{d\sigma_{B_c}}{dz} = D_{\bar{b} \rightarrow B_c}(z) \cdot \sigma_{b\bar{b}}$$



Feynman diagrams for $e^+e^- \rightarrow B_c + X$.

$$D_{\bar{b} \rightarrow B_c^*}(z) = \frac{2\alpha^2 |R_S(0)|^2}{81\pi m_c^3} \frac{rz(1-z)^2}{(1-(1-r)z)^6} \times \\ \times (6 - 18(1-2r)z + (21 - 74r + 68r^2)z^2 - 2(1-r)(6 - 19r + 18r^2)z^3 + 3(1-r)^2(1-2r+2r^2)z^4)$$

$$D_{\bar{b} \rightarrow B_c}(z) = \frac{2\alpha^2 |R_S(0)|^2}{27\pi m_c^3} \frac{rz(1-z)^2}{(1-(1-r)z)^6} \times \\ \times (2 - 2(3-2r)z + 3(3-2r+4r^2)z^2 - 2(1-r)(4-r+2r^2)z^3 + (1-r)^2(3-2r+2r^2)z^4)$$

$$r = \frac{m_c}{m_c + m_b}$$

The pQCD motivated FFs are used within FONLL to describe $c \rightarrow D$ (r — free parameter).

Peterson FF

To describe heavy quark hadronization into heavy-light meson the Peterson FF is usually used:

$$D_{Q \rightarrow (Q\bar{q})}(z) \sim \frac{1}{z \left(1 - \frac{1}{z} - \frac{\epsilon}{1-z}\right)^2}.$$


The dependence of "nonperturbative" Peterson FF is partially determined by denominator of perturbative propagator for $Q^* \rightarrow (Q\bar{q}) + jet$ process:

$$m_Q^2 - (P_{(Q\bar{q})} + P_{jet})^2.$$

$$m_Q^2 - (P_{(Q\bar{q})} + P_{jet})^2 \approx m_Q^2 - \frac{m_{(Q\bar{q})}^2}{z} - \frac{m_{jet}^2}{1-z}.$$

For $m_Q \approx m_{(Q\bar{q})}$:

$$m_Q^2 - (P_{(Q\bar{q})} + P_{jet})^2 \sim 1 - \frac{1}{z} - \frac{m_{jet}^2}{m_Q^2} \frac{1}{1-z}.$$



FF for $b \rightarrow B_c^{(*)}$ obtained within pQCD vs. Peterson parametrization.

Kartvelishvili-Likhoded-Petrov FF [6]

$$D_{KLP}(z) = (\alpha + 1)(\alpha + 2)z^\alpha(1 - z)$$

FF	the main z dependence
pQCD motivated	propagator (+WF)
Peterson	propagator
KLP	WF


- Shape is close to shapes of other FFs.
- Formula looks like the formula for the hadronic structure function.
- Based on Gribov-Lipatov reciprocity.

$$|\mathbf{hadron}\rangle = \sum_i |\mathbf{parton}\rangle_i \iff$$

$$|\mathbf{parton}\rangle = \sum_j |\mathbf{hadron}\rangle_j$$



- Any approach for heavy quark production can be used with any FF parameterizations.
- But the FF parameter values essentially depend on the choice of heavy quark production model.
- Only the cross section of heavy hadrons can be measured experimentally.

Open beauty production in e^+e^- collisions



OPAL [7], ALEPH [8], SLD [9]



Open charm production at LHCb



Differential cross-sections for Λ_c^+ baryon production compared to the theoretical prediction from the GMVFNS scheme.



Differential cross-sections for (a) D^0 , (b) D^+ , (c) D^{*+} , and (d) D_s^+ meson production compared to theoretical predictions.

Hadronization effects in LHCb data [12] (I)




The considered models do not account precisely for all the features observed in the LHCb data, but theoretical predictions including 1% intrinsic charm and 10% recombination contributions (MS [10]) better describe the data than the other models considered (Pythia, Vogt [11]).

Hadronization effects in LHCb data [12] (II)



Good news: the cross section values can be fairly described within the mentioned approaches.

Recombination mechanism




$$\frac{d\hat{\sigma}[\bar{q}g \rightarrow (Q\bar{q}) + \bar{Q}]}{d\hat{\sigma}[gg \rightarrow Q\bar{Q}]} \Big|_{\theta=\pi/2} \approx \frac{256\pi}{189} \alpha_s \frac{m_Q^2}{m_Q^2 + p_T^2}$$

$$\frac{d\hat{\sigma}[\bar{q}g \rightarrow (Q\bar{q}) + \bar{Q}]}{dp_T} \sim \frac{1}{p_T^6}$$

$$\frac{d\hat{\sigma}[\bar{q}g \rightarrow (Q\bar{q}) + \bar{Q}]}{d\hat{\sigma}[gg \rightarrow Q\bar{Q}]} \Big|_{\theta=0} \approx \frac{256\pi}{81} \alpha_s \sim 1$$

The result above can be obtained within HQET [13] or within NRQCD motivated model [14]. The recombination is not suppressed in the forward (backward) region!


$\Lambda_c \rightarrow pK\pi$ at SPD



Seems like Λ_c can be investigated at SPD conditions using the simplest mode $\Lambda_c \rightarrow pK\pi$. The signal significance after 1 year data taking was preliminary estimated as

$$\frac{N_{sig}}{\sqrt{N_{bg}}} \sim 36$$

Many thanks to Atrem Smirnov for his study!



$$\Lambda_c \rightarrow K_s p \pi^- \pi^+ \text{ and } \Lambda_c \rightarrow \Lambda^0 \pi^- \pi^+ \pi^+$$

Experiment: BIS-2 [15]

Serpukhov accelerator

Beam: neutrons;


$\sqrt{s} \approx 10$ GeV;

Target: C.

The mode was chosen due to a reliable detection of the decays of subsequent long-living states:

$K_s \rightarrow \pi^+ \pi^-$ and $\Lambda^0 \rightarrow p \pi^-$.

Kinematic region: ($X_F > 0.5$)



$$\Lambda_c \rightarrow K_s p \text{ and } \Lambda_c \rightarrow \Lambda^0 \pi^+$$

Experiment: FOCUS [16] [15]

FermiLab

Beam: bremsstrahlung photons from electrons;


$\sqrt{s_e} \approx 300 \text{ GeV}$;

Target: Be.

A broad structure at around 2.2 GeV/c coming from the decay mode

$\Lambda_c \rightarrow \Sigma^0(\Lambda\gamma)\pi^+$ where the photon from the Σ^0 decay is not reconstructed.

(See also [17, 18])



$$\bar{D}^0 \rightarrow K^+ \pi^- \pi^+ \pi^-$$

Experiment: BIS-2 [19]

Serpukhov accelerator

Beam: neutrons;

$\sqrt{s} \approx 10$ GeV;



Target: H, C and Al.

At least four charged particles with a common vertex in the target region were selected. Cerenkov threshold hodoscopes were used for kaon and pion identification.

Kinematic region:

$30 < P_L < 60$ GeV/c ($X_F \gtrsim 0.5$)
and $p_T < 1$ GeV/c.

As $\sim 78\%$ of the decay goes through the intermediate ρ^0 state, the signal quality can be improved with cut on $\pi^+ \pi^-$ invariant mass.



At least one of the $\pi^+ \pi^-$ pairs is in the ρ^0 mass region:

$$660 \leq M_{\pi^+ \pi^-} \leq 860 \text{ GeV}/c^2$$

$$D^{+*} \rightarrow D^0 \pi^+$$

Experiment: BEATRICE [20]

CERN

Beam: π^- ;

$\sqrt{s} \approx 350$ GeV;

Target: Si, W and Cu.


It is possible to observe a narrow peak in the mass difference distribution:

$$m_{D^{*+}(K^- \pi^+ \pi^+)} - m_{D^0(K^- \pi^+)}$$

The momentum of soft pion from $D^{+*} \rightarrow D^0 \pi^+$ can be estimated as

$$P_{\pi_{soft}^+} \sim \frac{m_\pi}{m_D} P_D \sim 0.1 P_D.$$

Thus an essential amount of D^0 with $P_{D^0} \gtrsim 1.5$ GeV is needed to observe $m_{D^{*+}}$ at SPD.



Plans, suggestions and conclusions

Λ_c -baryon:

- Try to improve selection quality for $\Lambda_c \rightarrow pK\pi$ decay;
- Try modes $\Lambda_c \rightarrow \Lambda^0\pi^-\pi^+\pi^+$ and $\Lambda_c \rightarrow K_s p\pi^-\pi^+$;

D -meson:

- Search for D^* meson in mass difference $m(K\pi\pi) - m(K\pi)$ to obtain narrower peak;
- $D^+ \rightarrow (K^+(890) \rightarrow K_s^0\pi^+)\pi^+\pi^-$;
- $D^0 \rightarrow K^-\pi^+\pi^+\pi^-$.

This study was carried out within the project "Production of open charm and tau leptons at the NICA SPD: phenomenology and simulation" supported by JINR.

Thank you for your attention!

- [1] B. A. Kniehl et al. “Inclusive Charmed-Meson Production at the CERN LHC”. In: *Eur. Phys. J. C* 72 (2012), p. 2082. DOI: [10.1140/epjc/s10052-012-2082-2](https://doi.org/10.1140/epjc/s10052-012-2082-2). arXiv: [1202.0439 \[hep-ph\]](https://arxiv.org/abs/1202.0439).
- [2] Matteo Cacciari, Mario Greco, and Paolo Nason. “The P(T) spectrum in heavy flavor hadroproduction”. In: *JHEP* 05 (1998), p. 007. DOI: [10.1088/1126-6708/1998/05/007](https://doi.org/10.1088/1126-6708/1998/05/007). arXiv: [hep-ph/9803400](https://arxiv.org/abs/hep-ph/9803400).
- [3] S. P. Baranov, A. V. Lipatov, and N. P. Zotov. “Heavy quark production at LHC as a test of the k(T)-factorization approach”. In: *Nucl. Phys. B Proc. Suppl.* 146 (2005). Ed. by R. Fiore et al., pp. 228–230. DOI: [10.1016/j.nuclphysbps.2005.02.102](https://doi.org/10.1016/j.nuclphysbps.2005.02.102).
- [4] S. P. Baranov, A. V. Lipatov, and N. P. Zotov. “Prompt J/Ψ production at LHC: new evidence for the k_t -factorization”. In: *Phys. Rev. D* 85 (2012), p. 014034. DOI: [10.1103/PhysRevD.85.014034](https://doi.org/10.1103/PhysRevD.85.014034). arXiv: [1108.2856 \[hep-ph\]](https://arxiv.org/abs/1108.2856).
- [5] A. V. Lipatov, V. A. Saleev, and N. P. Zotov. “Heavy quark production at the Tevatron in the semihard QCD approach and the unintegrated gluon distribution”. In: (Dec. 2001). arXiv: [hep-ph/0112114](https://arxiv.org/abs/hep-ph/0112114).

- [6] V. G. Kartvelishvili, A. K. Likhoded, and V. A. Petrov. “On the Fragmentation Functions of Heavy Quarks Into Hadrons”. In: *Phys. Lett. B* 78 (1978), pp. 615–617. DOI: [10.1016/0370-2693\(78\)90653-6](https://doi.org/10.1016/0370-2693(78)90653-6).
- [7] G. Abbiendi et al. “Inclusive analysis of the b quark fragmentation function in Z decays at LEP”. In: *Eur. Phys. J. C* 29 (2003), pp. 463–478. DOI: [10.1140/epjc/s2003-01229-x](https://doi.org/10.1140/epjc/s2003-01229-x). arXiv: [hep-ex/0210031](https://arxiv.org/abs/hep-ex/0210031).
- [8] A. Heister et al. “Study of the fragmentation of b quarks into B mesons at the Z peak”. In: *Phys. Lett. B* 512 (2001), pp. 30–48. DOI: [10.1016/S0370-2693\(01\)00690-6](https://doi.org/10.1016/S0370-2693(01)00690-6). arXiv: [hep-ex/0106051](https://arxiv.org/abs/hep-ex/0106051).
- [9] Kenji Abe et al. “Precise measurement of the b quark fragmentation function in Z0 boson decays”. In: *Phys. Rev. Lett.* 84 (2000), pp. 4300–4304. DOI: [10.1103/PhysRevLett.84.4300](https://doi.org/10.1103/PhysRevLett.84.4300). arXiv: [hep-ex/9912058](https://arxiv.org/abs/hep-ex/9912058).
- [10] Rafal Maciula and Antoni Szczurek. “Recombination mechanism for D0-meson production and D0–antiD0 production asymmetry in the LHCb p+20Ne fixed-target experiment”. In: *Phys. Lett. B* 835 (2022),

p. 137530. DOI: 10.1016/j.physletb.2022.137530. arXiv: 2206.02750 [hep-ph].

- [11] R. Vogt. “Limits on Intrinsic Charm Production from the SeaQuest Experiment”. In: *Phys. Rev. C* 103.3 (2021), p. 035204. DOI: 10.1103/PhysRevC.103.035204. arXiv: 2101.02858 [hep-ph].
- [12] R. Aaij et al. “Open charm production and asymmetry in pNe collisions at $\sqrt{s_{NN}} = 68.5$ GeV”. In: *Eur. Phys. J. C* 83.6 (2023), p. 541. DOI: 10.1140/epjc/s10052-023-11641-5. arXiv: 2211.11633 [hep-ex].
- [13] Eric Braaten, Yu Jia, and Thomas Mehen. “Charm anti-charm asymmetries in photoproduction from heavy quark recombination”. In: *Phys. Rev. D* 66 (2002), p. 014003. DOI: 10.1103/PhysRevD.66.014003. arXiv: hep-ph/0111296.
- [14] A. V. Berezhnoy and A. K. Likhoded. “Charge asymmetry in the photonic production of charmed mesons”. In: *Phys. Atom. Nucl.* 69 (2006), pp. 103–112. DOI: 10.1134/S1063778806010145. arXiv: hep-ph/0506124.

- [15] A. N. Alev et al. "The $\Lambda(c)+$ Baryon Production in Interactions of 40-GeV to 70-GeV Neutrons With Carbon Nuclei". In: *Z. Phys. C* 23 (1984), p. 333. DOI: [10.1007/BF01572656](https://doi.org/10.1007/BF01572656).
- [16] J. M. Link et al. "Study of Lambda+(c) Cabibbo favored decays containing a Lambda baryon in the final state". In: *Phys. Lett. B* 624 (2005), pp. 22–30. DOI: [10.1016/j.physletb.2005.08.014](https://doi.org/10.1016/j.physletb.2005.08.014). arXiv: [hep-ex/0505077](https://arxiv.org/abs/hep-ex/0505077).
- [17] P. Avery et al. "Inclusive production of the charmed baryon Lambda(c) from e+ e- annihilations at $s^{*(1/2)} = 10.55\text{-GeV}$ ". In: *Phys. Rev. D* 43 (1991), pp. 3599–3610. DOI: [10.1103/PhysRevD.43.3599](https://doi.org/10.1103/PhysRevD.43.3599).
- [18] H. Albrecht et al. "A Measurement of asymmetry in the decay $\lambda(c)+ \rightarrow \lambda \pi^+$ ". In: *Phys. Lett. B* 274 (1992), pp. 239–245. DOI: [10.1016/0370-2693\(92\)90529-D](https://doi.org/10.1016/0370-2693(92)90529-D).
- [19] A. N. Alev et al. "The Study of anti-D0 $\rightarrow K^+ \pi^- \pi^+ \pi^-$ in neutron interactions with hydrogen, carbon and aluminium at $s^{*(1/2)}$ approximately 10-GeV". In: *Czech. J. Phys.* 42 (1992), pp. 953–1048. DOI: [10.1007/BF01589533](https://doi.org/10.1007/BF01589533).

- [20] M. Adinolfi et al. “ D^{*+} production in 350-GeV/c π -N interactions”.
In: *Nucl. Phys. B* 547 (1999), pp. 3–18. DOI:
[10.1016/S0550-3213\(99\)00173-X](https://doi.org/10.1016/S0550-3213(99)00173-X).