

ELECTRIC DIPOLE POLARIZABILITY OF MAGIC NUCLEI

Arsenyev Nikolay

Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna, Russia

In collaboration with

A. P. Severyukhin — BLTP, JINR, Dubna

Session-Conference
of the Section of Nuclear Physics
Russian Academy of Sciences
Dubna, Russia
1 April 2024

Outline

Introduction

Realization of RPA with the Skyrme EDF

Results and discussion


Conclusions

N. Arsenyev



Introduction

$E1$ strength in (spherical) atomic nuclei





Courtesy: N. Pietralla

N. Arsenyev



Introduction

The electric dipole polarizability



External Field



Balanced



Displacement
Dipole Polarizability

Restoring Force

Courtesy: A. Tamii

$$\alpha_D = \frac{8\pi}{9} \sum_{\nu} \frac{B(E1; 0_{g.st.}^+ \rightarrow 1_{\nu}^-)}{E_{1_{\nu}^-}}$$

N. Arsenyev



REALIZATION OF RPA

N. Arsenyev



Realization of RPA

The starting point of the method is the HF-BCS calculations of the ground state, where spherical symmetry is assumed for the ground states. The continuous part of the single-particle spectrum is discretized by diagonalizing the HF Hamiltonian on a harmonic oscillator basis.

J. P. Blaizot and D. Gogny, Nucl. Phys. A284, 429 (1977).

We employ the effective Skyrme interaction in the particle-hole channel

$$V(\vec{r}_1, \vec{r}_2) = t_0 \left(1 + x_0 \hat{P}_\sigma\right) \delta(\vec{r}_1 - \vec{r}_2) + \frac{t_1}{2} \left(1 + x_1 \hat{P}_\sigma\right) \left[\delta(\vec{r}_1 - \vec{r}_2) \vec{k}^2 + \vec{k}'^2 \delta(\vec{r}_1 - \vec{r}_2)\right] \\ + t_2 \left(1 + x_2 \hat{P}_\sigma\right) \vec{k}' \cdot \delta(\vec{r}_1 - \vec{r}_2) \vec{k} + \frac{t_3}{6} \left(1 + x_3 \hat{P}_\sigma\right) \delta(\vec{r}_1 - \vec{r}_2) \rho^\alpha \left(\frac{\vec{r}_1 + \vec{r}_2}{2}\right) \\ + iW_0 (\vec{\sigma}_1 + \vec{\sigma}_2) \cdot \left[\vec{k}' \times \delta(\vec{r}_1 - \vec{r}_2) \vec{k}\right].$$

T. H. R. Skyrme, Nucl. Phys. 9, 615 (1959).

D. Vautherin and D. M. Brink, Phys. Rev. C5, 626 (1972).

The residual interaction in the particle-hole channel V_{res}^{ph} can be obtained as the second derivative of the energy density functional \mathcal{H} with respect to the particle density ρ .

G. T. Bertsch and S. F. Tsai, Phys. Rep. 18, 125 (1975).

N. Arsenyev



Realization of RPA

We introduce the phonon creation operators

$$Q_{\lambda\mu i}^+ = \frac{1}{2} \sum_{jj'} \left[X_{jj'}^{\lambda i} A^+(jj'; \lambda\mu) - (-1)^{\lambda-\mu} Y_{jj'}^{\lambda i} A(jj'; \lambda - \mu) \right],$$
$$A^+(jj'; \lambda\mu) = \sum_{mm'} C_{jmj'm'}^{\lambda\mu} \alpha_{jm}^+ \alpha_{j'm'}^+.$$

The index λ denotes total angular momentum and μ is its z -projection in the laboratory system. One assumes that the ground state is the RPA phonon vacuum $|0\rangle$ and one-phonon excited states are $Q_{\lambda\mu i}^+|0\rangle$ with the normalization condition

$$\langle 0|[Q_{\lambda\mu i}, Q_{\lambda\mu i'}^+]|0\rangle = \delta_{ii'}.$$

Making use of the linearized equation-of-motion approach one can get the RPA equations

$$\begin{pmatrix} A & B \\ -B & -A \end{pmatrix} \begin{pmatrix} X \\ Y \end{pmatrix} = \omega \begin{pmatrix} X \\ Y \end{pmatrix}.$$

Solutions of this set of linear equations yield the one-phonon energies ω and the amplitudes X, Y of the excited states.

RESULTS AND DISCUSSION

Details of calculations

The dipole transition probabilities can be expressed as

$$B(E1; 0_{gs}^+ \rightarrow 1_i^-) = \left| e_{eff}^{(n)} \langle i | \hat{M}^{(n)} | 0 \rangle + e_{eff}^{(p)} \langle i | \hat{M}^{(p)} | 0 \rangle \right|^2,$$

where $\hat{M}^{(p)} = \sum_i^Z r_i Y_{1\mu}(\hat{r}_i)$ and $\hat{M}^{(n)} = \sum_i^N r_i Y_{1\mu}(\hat{r}_i)$. The spurious 1^- state is excluded from the excitation spectra by introduction of the effective neutron $e_{eff}^{(n)} = -Z/A e$ and proton $e_{eff}^{(p)} = N/A e$ charges.

A. Bohr and B. Mottelson, Nuclear Structure Vol. II (Benjamin, New York 1975).


For our systematic analysis of the electric dipole polarizability, we employ 21 Skyrme interactions: LNS, SAMi, SGII, SIII, SK255, SkI2, SkI3, SkI5, SkM*, SkP, SkT5, SkT6, SkT7, SkX, SLy4, SLy5, SVbas, SVmas07, SVmas08, SVmas10, and SVmin. The choice of these parameterizations is due to the large range of values for the effective nucleon mass $m^* = 0.58-1.00$ and the symmetry energy at saturation density $J = 26.8-37.4$ MeV.

M. Dutra et al., Phys. Rev. C85, 035201 (2012).

N. Arsenyev



The photoabsorption cross section for ^{40}Ca




N. N. Arsenyev, A. P. Severyukhin, in preparation.

R. W. Fearick et al., Phys. Rev. Res. 5, L022044 (2023).

N. Arsenyev



The photoabsorption cross section for ^{48}Ca




N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

J. Birkhan et al., PRL 118, 252501 (2017).

N. Arsenyev



The electric dipole polarizability α_D in $^{40,48}\text{Ca}$



N. N. Arsenyev, A. P. Severyukhin, in preparation.

J. Birkhan et al., PRL 118, 252501 (2017).


R. W. Fearick et al., Phys. Rev. Res. 5, L022044 (2023).

G. Hagen et al., Nature Phys. 12, 186 (2016).

N. Arsenyev



The electric dipole polarizability α_D : Expt. vs Theory




N. N. Arsenyev, A. P. Severyukhin, in preparation.

N. Arsenyev



The neutron skin thickness ΔR_{np} in $^{40,48}\text{Ca}$



N. N. Arsenyev, A. P. Severyukhin, in preparation.

J. Birkhan et al., PRL 118, 252501 (2017).


R. W. Fearick et al., Phys. Rev. Res. 5, L022044 (2023).

G. Hagen et al., Nature Phys. 12, 186 (2016).

N. Arsenyev



The neutron skin thickness ΔR_{np} in ^{132}Sn and ^{208}Pb



N. N. Arsenyev, A. P. Severyukhin, in preparation.

A. Tamii et al., PRL 107, 062502 (2011).

A. Klimkiewicz et al., Phys. Rev. C76, 051603 (2007).

K. Hebeler et al., PRL 105, 161102 (2010).

N. Arsenyev



Pearson correlation coefficient

In statistics, the Pearson correlation coefficient (r) is a measure of linear correlation between two sets of data: $(a_1, b_1), \dots, (a_N, b_N)$. The Pearson coefficient calculate as

$$r = \frac{\sum_{i=1}^N (a_i - \bar{a})(b_i - \bar{b})}{\sigma_a \sigma_b},$$

where \bar{a} , \bar{b} represent the mean values and σ_a , σ_b the root mean square of the sample. The coefficient r always has a value between -1 and $+1$. Correlations equal to -1 or $+1$ correspond to data points lying exactly on a line. In fact, we may transform a_k to \tilde{a}_k , where

$$\tilde{a}_k = r \frac{\sigma_a}{\sigma_b} (b_k - \bar{b}) + \bar{a}.$$





For any value of the coefficient r , the value of \tilde{a}_k is the best linear approximation for a_k .

N. Draper and H. Smith, Applied Regression Analysis (Wiley, New York, 1998).

N. Arsenyev



Correlations: α_D vs ΔR_{np}



N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

N. Arsenyev



The nuclear symmetry energy J

One of the key properties of nuclear matter is the symmetry energy (J), which is particularly important in modeling nuclear matter and finite nuclei because it probes the isospin part of the Skyrme interaction. It is defined as

$$J(\rho) = \frac{\hbar^2}{6m} \left(\frac{3\pi^2}{2} \right)^{2/3} \rho^{2/3} - \frac{1}{8} t_0 (2x_0 + 1) \rho - \frac{1}{24} \left(\frac{3\pi^2}{2} \right)^{2/3} \rho^{2/3} [3t_1 x_1 - t_2 (4 + 5x_2)] - \frac{1}{48} t_3 (2x_3 + 1) \rho^{\alpha+1}.$$

Saturation properties of a few Skyrme parametrizations used in this work.





	SLy4	SAMi	SIII	SGII	SkM*	SkP	SkI2	SkT5	SK255	LNS
J	32.00	28.00	28.16	26.83	30.03	30.00	33.37	37.00	37.40	33.43
ρ_{sat}	0.160	0.159	0.145	0.158	0.160	0.163	0.158	0.164	0.157	0.175

M. Dutra et al., Phys. Rev. C85, 035201 (2012).

N. Arsenyev



Correlations: α_D vs ΔR_{np} and J




N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

N. Arsenyev



Estimation of the symmetry energy J

We carried out a theoretical analysis of the recently measured α_D and ΔR_{np} in ^{48}Ca , ^{68}Ni , and ^{208}Pb to extract information about the symmetry energy, by using a strong correlation between $\alpha_D J$ and ΔR_{np} . Combining the experimental data and the RPA theory constraints yields the interval of $J = 30 - 37 \text{ MeV}^\dagger$.



N. E. Solonovich, N. N. Arsenyev, A. P. Severyukhin, Phys. Part. Nucl. Letters. 19, 473 (2022)

A. Tamii et al., PRL 107, 062502 (2011).

D. M. Rossi et al., PRL 111, 242503 (2013).

J. Birkhan et al., Phys. Rev. Lett. 118, 252501 (2017).

[†] The current global average $J = 31.60 \pm 0.92 \pm 2.66 \text{ MeV}$ obtained using the 28 analyses of terrestrial nuclear experiments and astrophysical observations [B. A. Li and X. Han, Phys. Lett. B727, 276 (2013)]

N. Arsenyev



Conclusions

The Skyrme RPA calculations have been performed for the electric dipole response in the magic nuclei ^{16}O , $^{40,48,60}\text{Ca}$, $^{56,68,78}\text{Ni}$, ^{90}Zr , $^{100,132}\text{Sn}$ and ^{208}Pb . We have computed the nuclear dipole polarizability (α_{D}) and neutron skin thickness (ΔR_{np}) of these nuclei using a broad set of Skyrme functionals. It is shown that the neutron skin thickness is correlated the product of the electric dipole polarizability and the symmetry energy (J) at saturation density.

We carried out a theoretical analysis of the recently measured α_{D} and ΔR_{np} in ^{48}Ca , ^{68}Ni , and ^{208}Pb to extract information about the symmetry energy, by using a strong correlation between α_{D} and ΔR_{np} . Combining the experimental data and the RPA theory (using a broad set of Skyrme functionals) constraints yields the interval of $J=30-37$ MeV.

This work was supported within the framework of the scientific program of the National Center for Physics and Mathematics, topic No. 6 “Nuclear and Radiation Physics” (stage 2023–2025).

N. Arsenyev



THE END

N. Arsenyev

