

Evaluating 'elliptic' master integrals at special kinematic values: using differential equations and their solutions via expansions near singular points

Vladimir A. Smirnov

Skobeltsyn Institute of Nuclear Physics of Moscow State University

calc-2018, Dubna, July 24, 2018

Based on [R. Lee, A. Smirnov & V.S., arXiv:1805.00227]

Based on [R. Lee, A. Smirnov & V.S., arXiv:1805.00227]

It is a sequel of [R. Lee, A. Smirnov & V.S.'17]:

Based on [R. Lee, A. Smirnov & V.S., arXiv:1805.00227]

It is a sequel of [R. Lee, A. Smirnov & V.S.'17]:

an algorithm to find a solution of differential equations for master integrals in the form of an ϵ -expansion series with numerical coefficients.

Based on [R. Lee, A. Smirnov & V.S., arXiv:1805.00227]

It is a sequel of [R. Lee, A. Smirnov & V.S.'17]:

an algorithm to find a solution of differential equations for master integrals in the form of an ϵ -expansion series with numerical coefficients.

The algorithm is based on using generalized power series expansions near singular points of the differential system, solving difference equations for the corresponding coefficients in these expansions and using matching to connect series expansions at two neighbouring points.

Typically, analytical results for Feynman integrals can be expressed in terms of harmonic polylogarithms or multiple polylogarithms which are very well mathematically studied special functions introduced by physicists.

Typically, analytical results for Feynman integrals can be expressed in terms of harmonic polylogarithms or multiple polylogarithms which are very well mathematically studied special functions introduced by physicists.

HPL, GiNaC

Typically, analytical results for Feynman integrals can be expressed in terms of harmonic polylogarithms or multiple polylogarithms which are very well mathematically studied special functions introduced by physicists.

HPL, GiNaC

The possibility to arrive at a result written in terms of these functions exists if one succeeds to turn to a so-called *canonical* basis [J.M. Henn'13] using rational transformations.

Typically, analytical results for Feynman integrals can be expressed in terms of harmonic polylogarithms or multiple polylogarithms which are very well mathematically studied special functions introduced by physicists.

HPL, GiNaC

The possibility to arrive at a result written in terms of these functions exists if one succeeds to turn to a so-called *canonical* basis [J.M. Henn'13] using rational transformations.

The ε -form is not always possible. The simplest counter example is the two-loop sunset diagram with three equal non-zero masses. Elliptic functions and their generalizations appear.

Our approach [R. Lee, A. Smirnov & V.S.'17] is oriented at situations where the ε -form is not possible.

Our approach [R. Lee, A. Smirnov & V.S.'17] is oriented at situations where the ε -form is not possible.

It is very natural to try to introduce new functions.

Our approach [R. Lee, A. Smirnov & V.S.'17] is oriented at situations where the ε -form is not possible.

It is very natural to try to introduce new functions.

Elliptic generalization of multiple polylogarithms motivated by two-loop examples, where the ε -form is impossible

[L. Adams, C. Bogner, A. Schweitzer & S. Weinzierl'16;
E. Remiddi & L. Tancredi'17; M. Hidding & F. Moriello'17;
J. Broedel, C. Duhr, F. Dulat & L. Tancredi'17, J. Ablinger et al.'17, J. Broedel, C. Duhr, F. Dulat, B. Penante & L. Tancredi'18]

Our approach [R. Lee, A. Smirnov & V.S.'17] is oriented at situations where the ε -form is not possible.

It is very natural to try to introduce new functions.

Elliptic generalization of multiple polylogarithms motivated by two-loop examples, where the ε -form is impossible

[L. Adams, C. Bogner, A. Schweitzer & S. Weinzierl'16;
E. Remiddi & L. Tancredi'17; M. Hidding & F. Moriello'17;
J. Broedel, C. Duhr, F. Dulat & L. Tancredi'17, J. Ablinger et al.'17, J. Broedel, C. Duhr, F. Dulat, B. Penante & L. Tancredi'18]

Still we are far, even in lower loops orders, from answering the following question:

'What is the class of functions which can appear in results for Feynman integrals in situations where ε -form is impossible'?

- Our setup

- Our setup
- Our example: four-loop generalized sunset diagrams with three massive and two massless propagators

- Our setup
- Our example: four-loop generalized sunset diagrams with three massive and two massless propagators
- Our goal: to use our algorithm and the corresponding code for our example in order to obtain new analytical results.

- Our setup
- Our example: four-loop generalized sunset diagrams with three massive and two massless propagators
- Our goal: to use our algorithm and the corresponding code for our example in order to obtain new analytical results.

We analytically evaluate the master integrals at threshold, $p^2 = 9m^2$, in an expansion in ε up to ε^1 .

- Our setup
- Our example: four-loop generalized sunset diagrams with three massive and two massless propagators
- Our goal: to use our algorithm and the corresponding code for our example in order to obtain new analytical results.
We analytically evaluate the master integrals at threshold, $p^2 = 9m^2$, in an expansion in ε up to ε^1 .
- Perspectives

Let us consider Feynman integrals with two scales and let x be the ratio of these scales.

Let us consider Feynman integrals with two scales and let x be the ratio of these scales.

DE

$$\partial_x \mathbf{J} = M(x, \varepsilon) \mathbf{J},$$

where $\mathbf{J} = (J_1, \dots, J_N)$ are N master integrals.

Let us consider Feynman integrals with two scales and let x be the ratio of these scales.

DE

$$\partial_x \mathbf{J} = M(x, \varepsilon) \mathbf{J},$$

where $\mathbf{J} = (J_1, \dots, J_N)$ are N master integrals.

We imply that all the singular points of DE are regular, i.e. we can reduce the DE to a local Fuchsian form at any singular point, i.e. if x_j is a singular point then

$$M(x) = \frac{A_j(x)}{x - x_j}$$

where $A_j(x)$ is regular at $x = x_j$ and $A_j(x_j) \neq 0$.

General solution

$$\mathbf{J}(x) = U(x) \mathbf{C},$$

General solution

$$\mathbf{J}(x) = U(x) \mathbf{C},$$

where \mathbf{C} is a column of constants, and U is an evolution operator

$$U(x) = P \exp \left[\int M(x) dx \right].$$

Expanding in a vicinity of each singular point.

Expanding in a vicinity of each singular point.

Take $x = 0$.

Expanding in a vicinity of each singular point.

Take $x = 0$.

The expansion is

$$U(x) = \sum_{\lambda \in S} x^\lambda \sum_{n=0}^{\infty} \sum_{k=0}^{K_\lambda} \frac{1}{k!} C(n + \lambda, k) x^n \ln^k x,$$

where S is a finite set of powers of the form $\lambda = r\epsilon$ with integer r , $K_\lambda \geq 0$ is an integer number corresponding to the maximal power of the logarithm.

Expanding in a vicinity of each singular point.

Take $x = 0$.

The expansion is

$$U(x) = \sum_{\lambda \in S} x^\lambda \sum_{n=0}^{\infty} \sum_{k=0}^{K_\lambda} \frac{1}{k!} C(n + \lambda, k) x^n \ln^k x,$$

where S is a finite set of powers of the form $\lambda = r\epsilon$ with integer r , $K_\lambda \geq 0$ is an integer number corresponding to the maximal power of the logarithm.

The goal is to determine S , K_λ , and the matrix coefficients $C(n + \lambda, k)$.

Suppose that DE are in a global normalized Fuchsian form

$$M(x) = \frac{A_0}{x} + \sum_{k=1}^s \frac{A_k}{x - x_k}$$

Suppose that DE are in a global normalized Fuchsian form

$$M(x) = \frac{A_0}{x} + \sum_{k=1}^s \frac{A_k}{x - x_k}$$

and for any $k = 0, \dots, s$ the matrix A_k is free of resonances, i.e. the difference of any two of its distinct eigenvalues is not integer.

Suppose that DE are in a global normalized Fuchsian form

$$M(x) = \frac{A_0}{x} + \sum_{k=1}^s \frac{A_k}{x - x_k}$$

and for any $k = 0, \dots, s$ the matrix A_k is free of resonances, i.e. the difference of any two of its distinct eigenvalues is not integer.

In particular, the 'elliptic' cases, as a rule, can algorithmically be reduced to a global normalized Fuchsian form using, e.g., the algorithm of Lee [R.N. Lee'14].

DE \rightarrow difference equations for the matrix coefficients
 $C(n + \lambda, k)$.

DE \rightarrow difference equations for the matrix coefficients
 $C(n + \lambda, k)$.

Our algorithm provides solutions with no more than a linear growth of computational complexity with respect to a required number of terms.

DE \rightarrow difference equations for the matrix coefficients
 $C(n + \lambda, k)$.

Our algorithm provides solutions with no more than a linear growth of computational complexity with respect to a required number of terms.

This is very important for the subsequent step: the matching procedure which enables one to connect series expansions at two neighbouring points and thereby to obtain the possibility to evaluate Feynman integrals at any given point.

DE \rightarrow difference equations for the matrix coefficients
 $C(n + \lambda, k)$.

Our algorithm provides solutions with no more than a linear growth of computational complexity with respect to a required number of terms.

This is very important for the subsequent step: the matching procedure which enables one to connect series expansions at two neighbouring points and thereby to obtain the possibility to evaluate Feynman integrals at any given point.

Boundary conditions are included at one of the singular points and then series expansions at other points can be obtained by matching, step by step, pairs of expansions at neighboring points.

Using series expansions at singular points and solving difference equations:

Using series expansions at singular points and solving difference equations:

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

(evaluating three-loop massive vacuum diagrams)

Using series expansions at singular points and solving difference equations:

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

(evaluating three-loop massive vacuum diagrams)

[R. Mueller & D. G. Öztürk'16; J. M. Henn, A. V. Smirnov & V. A. Smirnov'16]

(applying general theory of DE for evaluating expansion of two-scale integrals at a given singular point)

Using series expansions at singular points and solving difference equations:

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

(evaluating three-loop massive vacuum diagrams)

[R. Mueller & D. G. Öztürk'16; J. M. Henn, A. V. Smirnov & V. A. Smirnov'16]

(applying general theory of DE for evaluating expansion of two-scale integrals at a given singular point)

[K. Melnikov, L. Tancredi and C. Wever'16]

(evaluating expansions of solutions of DE at a given singular point by difference equations)

Using series expansions at singular points and solving difference equations:

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

(evaluating three-loop massive vacuum diagrams)

[R. Mueller & D. G. Öztürk'16; J. M. Henn, A. V. Smirnov & V. A. Smirnov'16]

(applying general theory of DE for evaluating expansion of two-scale integrals at a given singular point)


[K. Melnikov, L. Tancredi and C. Wever'16]

(evaluating expansions of solutions of DE at a given singular point by difference equations)


[X. Liu, Y.Q. Ma & C.Y. Wang'17]

(solving DE wrt η in propagators $1/(k^2 + i0) \rightarrow 1/(k^2 + i\eta)$)

Feynman integrals corresponding to the generalized sunset graph with two massless and three massive lines




Feynman integrals corresponding to the generalized sunset graph with two massless and three massive lines



$$F_{a_1, \dots, a_{14}} = \int \cdots \int \frac{d^D k_1 \dots d^D k_4 (k_1 \cdot p)^{a_6} (k_2 \cdot p)^{a_7} (k_3 \cdot p)^{a_8} (k_4 \cdot p)^{a_9}}{(-k_1^2)^{a_1} (-k_2^2)^{a_2} (m^2 - k_3^2)^{a_3} (m^2 - k_4^2)^{a_4} (m^2 - (\sum k_i + p)^2)^{a_5} \times (k_1 \cdot k_2)^{a_{10}} (k_1 \cdot k_3)^{a_{11}} (k_1 \cdot k_4)^{a_{12}} (k_2 \cdot k_3)^{a_{13}} (k_2 \cdot k_4)^{a_{14}}},$$

with $x = p^2/m^2$.

Feynman integrals corresponding to the generalized sunset graph with two massless and three massive lines




$$F_{a_1, \dots, a_{14}} = \int \cdots \int \frac{d^D k_1 \dots d^D k_4 (k_1 \cdot p)^{a_6} (k_2 \cdot p)^{a_7} (k_3 \cdot p)^{a_8} (k_4 \cdot p)^{a_9}}{(-k_1^2)^{a_1} (-k_2^2)^{a_2} (m^2 - k_3^2)^{a_3} (m^2 - k_4^2)^{a_4} (m^2 - (\sum k_i + p)^2)^{a_5} \times (k_1 \cdot k_2)^{a_{10}} (k_1 \cdot k_3)^{a_{11}} (k_1 \cdot k_4)^{a_{12}} (k_2 \cdot k_3)^{a_{13}} (k_2 \cdot k_4)^{a_{14}}},$$

with $x = p^2/m^2$.

There are four master integrals in this family.

Feynman integrals corresponding to the generalized sunset graph with two massless and three massive lines



$$F_{a_1, \dots, a_{14}} = \int \cdots \int \frac{d^D k_1 \dots d^D k_4 (k_1 \cdot p)^{a_6} (k_2 \cdot p)^{a_7} (k_3 \cdot p)^{a_8} (k_4 \cdot p)^{a_9}}{(-k_1^2)^{a_1} (-k_2^2)^{a_2} (m^2 - k_3^2)^{a_3} (m^2 - k_4^2)^{a_4} (m^2 - (\sum k_i + p)^2)^{a_5} \times (k_1 \cdot k_2)^{a_{10}} (k_1 \cdot k_3)^{a_{11}} (k_1 \cdot k_4)^{a_{12}} (k_2 \cdot k_3)^{a_{13}} (k_2 \cdot k_4)^{a_{14}}},$$

with $x = p^2/m^2$.

There are four master integrals in this family. We choose

$$\{F_{1,1,1,1,1,0,\dots,0}, F_{1,1,2,1,1,0,\dots,0}, F_{1,2,1,1,1,0,\dots,0}, F_{1,2,1,1,2,0,\dots,0}\}.$$

The singular points are

$$x_0 = 0, x_1 = 1, x_2 = 9, x_3 = x_{-1} = \infty$$

The singular points are

$$x_0 = 0, x_1 = 1, x_2 = 9, x_3 = x_{-1} = \infty$$

The code DESS.m

<https://bitbucket.org/feynmanintegrals/dess>

The singular points are

$$x_0 = 0, x_1 = 1, x_2 = 9, x_3 = x_{-1} = \infty$$

The code DESS.m

<https://bitbucket.org/feynmanintegrals/dess>

Using this code it is possible to evaluate master integrals at a given point as well as expansions at singular points with a required precision in an ϵ -expansion with a required number of terms.

The singular points are

$$x_0 = 0, x_1 = 1, x_2 = 9, x_3 = x_{-1} = \infty$$

The code DESS.m

<https://bitbucket.org/feynmanintegrals/dess>

Using this code it is possible to evaluate master integrals at a given point as well as expansions at singular points with a required precision in an ϵ -expansion with a required number of terms.

The goal: to evaluate master integrals considered at threshold, $p^2 = 9m^2$,

$$\{J_1 = F_{1,1,1,1,1,0,\dots,0}, J_2 = F_{1,1,2,1,1,0,\dots,0}, J_3 = F_{1,2,1,1,1,0,\dots,0}\}.$$

Choose a point to fix boundary conditions.

Choose a point to fix boundary conditions.

$x_0 = 0$: simple (leads to vacuum integrals with less indices)

Choose a point to fix boundary conditions.

$x_0 = 0$: simple (leads to vacuum integrals with less indices)

It is far from $x_2 = 9$.

Choose a point to fix boundary conditions.

$x_0 = 0$: simple (leads to vacuum integrals with less indices)

It is far from $x_2 = 9$.

$x_1 = 1$: it is closer to $x_2 = 9$.

Choose a point to fix boundary conditions.

$x_0 = 0$: simple (leads to vacuum integrals with less indices)

It is far from $x_2 = 9$.

$x_1 = 1$: it is closer to $x_2 = 9$.

analytical results up to weight 7 can be taken from

[P. Marquard, A. V. Smirnov, V. A. Smirnov &
M. Steinhauser'15]

Choose a point to fix boundary conditions.

$x_0 = 0$: simple (leads to vacuum integrals with less indices)

It is far from $x_2 = 9$.

$x_1 = 1$: it is closer to $x_2 = 9$.

analytical results up to weight 7 can be taken from

[P. Marquard, A. V. Smirnov, V. A. Smirnov &
M. Steinhauser'15]

$x_3 = \infty$: simple constants; it is also a neighbour of $x_2 = 9$.

Choose a point to fix boundary conditions.

$x_0 = 0$: simple (leads to vacuum integrals with less indices)

It is far from $x_2 = 9$.

$x_1 = 1$: it is closer to $x_2 = 9$.

analytical results up to weight 7 can be taken from

[P. Marquard, A. V. Smirnov, V. A. Smirnov & M. Steinhauser'15]

$x_3 = \infty$: simple constants; it is also a neighbour of $x_2 = 9$.

The corresponding expansion is a large-momentum expansion [K.G. Chetyrkin'88, V.S.'90] where every term is a product of one-loop tadpoles and massless propagator integrals. It provides any required accuracy and any required number of terms in ε -expansions in the boundary conditions.

`DESS[rdatas, x, f(x), oe, np, nt, ns]`
where `ns` means the number of a singular point and this number is 1 for x_0 , 2 for x_1 , and 4 for x_3 .

`DESS[rdatas, x, f(x), oe, np, nt, ns]`
where ns means the number of a singular point and this number is 1 for x_0 , 2 for x_1 , and 4 for x_3 .

We choose $ns=4$.

`DESS[rdatas, x, f(x), oe, np, nt, ns]`

where ns means the number of a singular point and this number is 1 for x_0 , 2 for x_1 , and 4 for x_3 .

We choose $ns=4$.

Using DESS we obtain numerical results for the threshold master integrals in an ε -expansion up to ε^2 with the accuracy of 20000 digits for the corresponding coefficients.

`DESS[rdatas, x, f(x), oe, np, nt, ns]`
where ns means the number of a singular point and this number is 1 for x_0 , 2 for x_1 , and 4 for x_3 .

We choose $ns=4$.

Using DESS we obtain numerical results for the threshold master integrals in an ε -expansion up to ε^2 with the accuracy of 20000 digits for the corresponding coefficients.

Apply the PSLQ algorithm

[H.R.P. Ferguson, D.H. Bailey & S. Arno'99]

`DESS[rdatas, x, f(x), oe, np, nt, ns]`
where ns means the number of a singular point and this number is 1 for x_0 , 2 for x_1 , and 4 for x_3 .

We choose $ns=4$.

Using DESS we obtain numerical results for the threshold master integrals in an ε -expansion up to ε^2 with the accuracy of 20000 digits for the corresponding coefficients.

Apply the PSLQ algorithm

[H.R.P. Ferguson, D.H. Bailey & S. Arno'99]

`FindIntegerNullVector` in Mathematica

The choice of a basis of constants?

The choice of a basis of constants?

Results for the two-loop sunset diagram at threshold

[F.A. Berends & A.I. Davydychev '97, A.I. Davydychev & V.S. '99]:

multiple polylogarithm values at sixth roots of unity up to weight 3 [D.J. Broadhurst '98]

and $\frac{\pi}{\sqrt{3}}$.

The choice of a basis of constants?

Results for the two-loop sunset diagram at threshold

[F.A. Berends & A.I. Davydychev'97, A.I. Davydychev & V.S.'99]:

multiple polylogarithm values at sixth roots of unity up to weight 3 [D.J. Broadhurst'98]

and $\frac{\pi}{\sqrt{3}}$.

[J. Fleischer & M.Y. Kalmykov'99, A.I. Davydychev & M.Y. Kalmykov'00, M.Y. Kalmykov & B.A. Kniehl'10]:

include $\sqrt{3}$ separately.

The choice of a basis of constants?

Results for the two-loop sunset diagram at threshold

[F.A. Berends & A.I. Davydychev'97, A.I. Davydychev & V.S.'99]:

multiple polylogarithm values at sixth roots of unity up to weight 3 [D.J. Broadhurst'98]

and $\frac{\pi}{\sqrt{3}}$.

[J. Fleischer & M.Y. Kalmykov'99, A.I. Davydychev & M.Y. Kalmykov'00, M.Y. Kalmykov & B.A. Kniehl'10]:

include $\sqrt{3}$ separately.

Let us use multiple polylogarithm values at sixth roots of unity constructed up to weight 6

[J.M. Henn, A.V. Smirnov & V.S.'17] and $\sqrt{3}$.

$$G(a_1, \dots, a_w; 1),$$

where the indices a_i are equal to zero or a sixth root of unity, i.e. taken from the alphabet $\{0, r_1, r_3, -1, r_4, r_2, 1\}$ with

$$r_{1,2} = \frac{1}{2} \left(1 \pm \sqrt{3}i \right) = \lambda^{\pm 1}, \quad r_{3,4} = \frac{1}{2} \left(-1 \pm \sqrt{3}i \right) = \lambda^{\pm 2},$$

$$\lambda = e^{\pi i/3} = r_1 \text{ and } a_1 \neq 1.$$

$$G(a_1, \dots, a_w; 1),$$

where the indices a_i are equal to zero or a sixth root of unity, i.e. taken from the alphabet $\{0, r_1, r_3, -1, r_4, r_2, 1\}$ with

$$r_{1,2} = \frac{1}{2} \left(1 \pm \sqrt{3}i \right) = \lambda^{\pm 1}, \quad r_{3,4} = \frac{1}{2} \left(-1 \pm \sqrt{3}i \right) = \lambda^{\pm 2},$$

$$\lambda = e^{\pi i/3} = r_1 \text{ and } a_1 \neq 1.$$

$$G(a_1, \dots, a_w; z) = \int_0^z \frac{1}{t - a_1} G(a_2, \dots, a_w; t) dt$$

with $a_i, z \in \mathbb{C}$ and $G(z) = 1$.

$$G(0, \dots, 0; z) = \frac{1}{n!} \log^n z.$$

$$G(a_1, \dots, a_w; 1) = G_R(a_1, \dots, a_w) + i G_I(a_1, \dots, a_w)$$

$$G(a_1, \dots, a_w; 1) = G_R(a_1, \dots, a_w) + i G_I(a_1, \dots, a_w)$$

Let us denote by $B_R(w)$ ($B_I(w)$) the bases generated by $G_R(a_1, \dots, a_w)$ ($G_I(a_1, \dots, a_w)$).

$$G(a_1, \dots, a_w; 1) = G_R(a_1, \dots, a_w) + i G_I(a_1, \dots, a_w)$$

Let us denote by $B_R(w)$ ($B_I(w)$) the bases generated by $G_R(a_1, \dots, a_w)$ ($G_I(a_1, \dots, a_w)$).

[J.M. Henn, A.V. Smirnov & V.S.'17]:

$$B_R(1) = \left\{ G_R(-1) = \log(2), \quad G_R(r_4) = \frac{1}{2} \log(3) \right\},$$

$$B_I(1) = \left\{ G_I(r_2) = -\frac{\pi}{3} \right\}.$$

$$B_R(2) = \{GR[r2, -1], GR[-1]^2, GI[r2]^2, GR[-1] GR[r4], GR[r4]^2\}$$

$$B_I(2) = \{GI[0, r2], GI[r2] GR[-1], GI[r2] GR[r4]\}$$

$$B_R(3) = \{GR[0, 0, 1], GR[r2, 1, -1], GR[r2, 1, r3], GR[-1]^3, GI[r2]^2 GR[-1], GR[-1]^2 GR[r4], GI[r2]^2 GR[r4], GR[-1] GR[r4]^2, GR[r4]^3, GI[r2] GI[0, r2], GR[-1] GR[r2, -1], GR[r4] GR[r2, -1]\}$$

$$B_I(3) = \{GI[0, 1, r4], GI[0, r2, -1], GI[r2] GR[-1]^2, GI[r2]^3, GI[r2] GR[-1] GR[r4], GI[r2] GR[r4]^2, GI[0, r2] GR[-1], GI[0, r2] GR[r4], GI[r2] GR[r2, -1]\}$$

$$B_R(4) =$$

```
{GR[0, 0, r2, -1], GR[0, 0, r4, 1], GR[r2, 1, 1, -1],
GR[r2, 1, 1, r3], GR[r2, 1, r2, -1]}
```

and

```
{GR[-1]^4, GI[r2]^2 GR[-1]^2, GI[r2]^4, GR[-1]^3 GR[r4],
GI[r2]^2 GR[-1] GR[r4], GR[-1]^2 GR[r4]^2, GI[r2]^2 GR[r4]^2,
GR[-1] GR[r4]^3, GR[r4]^4, GI[r2] GI[0, r2] GR[-1],
GI[r2] GI[0, r2] GR[r4], GI[0, r2]^2, GR[-1]^2 GR[r2, -1],
GI[r2]^2 GR[r2, -1], GR[-1] GR[r4] GR[r2, -1], GR[r4]^2 GR[r2, -1],
GR[r2, -1]^2, GR[-1] GR[0, 0, 1], GR[r4] GR[0, 0, 1],
GI[r2] GI[0, 1, r4], GI[r2] GI[0, r2, -1], GR[-1] GR[r2, 1, -1],
GR[r4] GR[r2, 1, -1], GR[-1] GR[r2, 1, r3], GR[r4] GR[r2, 1, r3]}
```

$$B_l(4) =$$

```
{GI[0, 0, 0, r2], GI[0, 1, 1, r4], GI[0, 1, r2, -1], GI[0, 1, r2, r3],
  GI[0, r2, 1, -1]}
```

and

```
{GI[r2] GR[-1]^3, GI[r2]^3 GR[-1], GI[r2] GR[-1]^2 GR[r4],
  GI[r2]^3 GR[r4], GI[r2] GR[-1] GR[r4]^2, GI[r2] GR[r4]^3,
  GI[0, r2] GR[-1]^2, GI[r2]^2 GI[0, r2], GI[0, r2] GR[-1] GR[r4],
  GI[0, r2] GR[r4]^2, GI[r2] GR[-1] GR[r2, -1],
  GI[r2] GR[r4] GR[r2, -1], GI[0, r2] GR[r2, -1], GI[r2] GR[0, 0, 1],
  GI[0, 1, r4] GR[-1], GI[0, 1, r4] GR[r4], GI[0, r2, -1] GR[-1],
  GI[0, r2, -1] GR[r4], GI[r2] GR[r2, 1, -1], GI[r2] GR[r2, 1, r3]}
```

$$B_R(5) =$$

```
{GR[0, 0, 0, 0, 1], GR[0, 0, 1, 1, -1], GR[0, 0, 1, 1, r4],
GR[0, 0, 1, r2, -1], GR[0, 0, 1, r2, r3], GR[0, 0, 1, r2, r4],
GR[0, 0, r2, 1, -1], GR[r2, 1, 1, -1, r2], GR[r2, 1, 1, 1, -1],
GR[r2, 1, 1, 1, r3], GR[r2, 1, 1, r2, -1], GR[r2, 1, 1, r2, r3],
GR[r2, 1, 1, r4, -1]}
```

and

```
{GR[-1]^5, GI[r2]^2 GR[-1]^3, GI[r2]^4 GR[-1], GR[-1]^4 GR[r4],
GI[r2]^2 GR[-1]^2 GR[r4], GI[r2]^4 GR[r4], GR[-1]^3 GR[r4]^2,
GI[r2]^2 GR[-1] GR[r4]^2, GR[-1]^2 GR[r4]^3, GI[r2]^2 GR[r4]^3,
GR[-1] GR[r4]^4, GR[r4]^5, GI[r2] GI[0, r2] GR[-1]^2,
GI[r2]^3 GI[0, r2], GI[r2] GI[0, r2] GR[-1] GR[r4],
GI[r2] GI[0, r2] GR[r4]^2, GI[0, r2]^2 GR[-1], GI[0, r2]^2 GR[r4],
GR[-1]^3 GR[r2, -1], GI[r2]^2 GR[-1] GR[r2, -1],
GR[-1]^2 GR[r4] GR[r2, -1], GI[r2]^2 GR[r4] GR[r2, -1],
GR[-1] GR[r4]^2 GR[r2, -1], GR[r4]^3 GR[r2, -1],
GI[r2] GI[0, r2] GR[r2, -1], GR[-1] GR[r2, -1]^2,
GR[r4] GR[r2, -1]^2, GR[-1]^2 GR[0, 0, 1], GI[r2]^2 GR[0, 0, 1],
GR[-1] GR[r4] GR[0, 0, 1], GR[r4]^2 GR[0, 0, 1],
GR[r2, -1] GR[0, 0, 1], GI[r2] GI[0, 1, r4] GR[-1],
GI[r2] GI[0, 1, r4] GR[r4], GI[0, r2] GI[0, 1, r4],
GI[r2] GI[0, r2, -1] GR[-1], GI[r2] GI[0, r2, -1] GR[r4],
GI[0, r2] GI[0, r2, -1], GR[-1]^2 GR[r2, 1, -1],
GI[r2]^2 GR[r2, 1, -1], GR[-1] GR[r4] GR[r2, 1, -1],
GR[r4]^2 GR[r2, 1, -1], GR[r2, -1] GR[r2, 1, -1],
GR[-1]^2 GR[r2, 1, r3], GI[r2]^2 GR[r2, 1, r3],
GR[-1] GR[r4] GR[r2, 1, r3], GR[r4]^2 GR[r2, 1, r3],
GR[r2, -1] GR[r2, 1, r3], GI[r2] GI[0, 0, 0, r2],
GR[-1] GR[0, 0, r2, -1], GR[r4] GR[0, 0, r2, -1],
GR[-1] GR[0, 0, r4, 1], GR[r4] GR[0, 0, r4, 1],
GI[r2] GI[0, 1, 1, r4], GI[r2] GI[0, 1, r2, -1],
GI[r2] GI[0, 1, r2, r3], GI[r2] GI[0, r2, 1, -1],
GR[-1] GR[r2, 1, 1, -1], GR[r4] GR[r2, 1, 1, -1],
GR[-1] GR[r2, 1, 1, r3], GR[r4] GR[r2, 1, 1, r3],
GR[-1] GR[r2, 1, r2, -1], GR[r4] GR[r2, 1, r2, -1]}
```


$$B_I(5) =$$

```
{GI[0, 0, 0, 1, r2], GI[0, 0, 0, 1, r4], GI[0, 0, 0, r2, -1],
GI[0, 1, 1, -1, r2], GI[0, 1, 1, -1, r4], GI[0, 1, 1, 1, r4],
GI[0, 1, 1, r2, r3], GI[0, 1, 1, r4, -1], GI[0, 1, 1, r4, r1],
GI[0, 1, r2, r3, r2], GI[0, r2, 1, 1, -1]}
```

and

```
{GI[r2] GR[-1]^4, GI[r2]^3 GR[-1]^2, GI[r2]^5, GI[r2] GR[-1]^3 GR[r4],
GI[r2]^3 GR[-1] GR[r4], GI[r2] GR[-1]^2 GR[r4]^2, GI[r2]^3 GR[r4]^2,
GI[r2] GR[-1] GR[r4]^3, GI[r2] GR[r4]^4, GI[0, r2] GR[-1]^3,
GI[r2]^2 GI[0, r2] GR[-1], GI[0, r2] GR[-1]^2 GR[r4],
GI[r2]^2 GI[0, r2] GR[r4], GI[0, r2] GR[-1] GR[r4]^2,
GI[0, r2] GR[r4]^3, GI[r2] GI[0, r2]^2, GI[r2] GR[-1]^2 GR[r2, -1],
GI[r2]^3 GR[r2, -1], GI[r2] GR[-1] GR[r4] GR[r2, -1],
GI[r2] GR[r4]^2 GR[r2, -1], GI[0, r2] GR[-1] GR[r2, -1],
GI[0, r2] GR[r4] GR[r2, -1], GI[r2] GR[r2, -1]^2,
GI[r2] GR[-1] GR[0, 0, 1], GI[r2] GR[r4] GR[0, 0, 1],
GI[0, r2] GR[0, 0, 1], GI[0, 1, r4] GR[-1]^2, GI[r2]^2 GI[0, 1, r4],
GI[0, 1, r4] GR[-1] GR[r4], GI[0, 1, r4] GR[r4]^2,
GI[0, 1, r4] GR[r2, -1], GI[0, r2, -1] GR[-1]^2,
GI[r2]^2 GI[0, r2, -1], GI[0, r2, -1] GR[-1] GR[r4],
GI[0, r2, -1] GR[r4]^2, GI[0, r2, -1] GR[r2, -1],
GI[r2] GR[-1] GR[r2, 1, -1], GI[r2] GR[r4] GR[r2, 1, -1],
GI[0, r2] GR[r2, 1, -1], GI[r2] GR[-1] GR[r2, 1, r3],
GI[r2] GR[r4] GR[r2, 1, r3], GI[0, r2] GR[r2, 1, r3],
GI[0, 0, 0, r2] GR[-1], GI[0, 0, 0, r2] GR[r4],
GI[r2] GR[0, 0, r2, -1], GI[r2] GR[0, 0, r4, 1],
GI[0, 1, 1, r4] GR[-1], GI[0, 1, 1, r4] GR[r4],
GI[0, 1, r2, -1] GR[-1], GI[0, 1, r2, -1] GR[r4],
GI[0, 1, r2, r3] GR[-1], GI[0, 1, r2, r3] GR[r4],
GI[0, r2, 1, -1] GR[-1], GI[0, r2, 1, -1] GR[r4],
GI[r2] GR[r2, 1, 1, -1], GI[r2] GR[r2, 1, 1, r3],
GI[r2] GR[r2, 1, r2, -1]}
```

In our case, with additional $\sqrt{3}$, we use the bases
 $B(w) = \{B_R(w), \sqrt{3}B_I(w)\}$ of weights $w = 1, 2, \dots$

In our case, with additional $\sqrt{3}$, we use the bases
 $B(w) = \{B_R(w), \sqrt{3}B_I(w)\}$ of weights $w = 1, 2, \dots$

The element $\sqrt{3}$ does not contribute to the weight and it is 'imaginary' in its character, so that elements from $\sqrt{3}B_I(w)$ are 'real'.

The numbers of elements are 3, 8, 21, 55, 144 for weights $w = 1, 2, 3, 4, 5$, correspondingly.

In our case, with additional $\sqrt{3}$, we use the bases $B(w) = \{B_R(w), \sqrt{3}B_I(w)\}$ of weights $w = 1, 2, \dots$

The element $\sqrt{3}$ does not contribute to the weight and it is 'imaginary' in its character, so that elements from $\sqrt{3}B_I(w)$ are 'real'.

The numbers of elements are 3, 8, 21, 55, 144 for weights $w = 1, 2, 3, 4, 5$, correspondingly.

If a constant is expected to be uniformly transcendental one can use these bases. Otherwise, one uses

$$\bar{B}(w) = \bigcup_{i=1}^w B(i) .$$

In our case, with additional $\sqrt{3}$, we use the bases $B(w) = \{B_R(w), \sqrt{3}B_I(w)\}$ of weights $w = 1, 2, \dots$

The element $\sqrt{3}$ does not contribute to the weight and it is 'imaginary' in its character, so that elements from $\sqrt{3}B_I(w)$ are 'real'.

The numbers of elements are 3, 8, 21, 55, 144 for weights $w = 1, 2, 3, 4, 5$, correspondingly.

If a constant is expected to be uniformly transcendental one can use these bases. Otherwise, one uses

$$\bar{B}(w) = \bigcup_{i=1}^w B(i) .$$

The numbers of elements in these bases are 4, 12, 33, 88, 232 for weights $w = 1, 2, 3, 4, 5$, correspondingly.

The accuracy of 2000 digits was quite enough to obtain results with PSLQ in an ε -expansion up to the finite part in ε , or, in other words, up to weight 4, in a straightforward way.

The accuracy of 2000 digits was quite enough to obtain results with PSLQ in an ε -expansion up to the finite part in ε , or, in other words, up to weight 4, in a straightforward way.

Let us look for uniformly transcendental threshold integrals. At $p^2 = m^2$, the integrals

$$\{J_4 = F_{1,2,2,2,2,0,\dots,0}, J_5 = F_{2,2,2,2,1,0,\dots,0}\}.$$

are uniformly transcendental. Let us assume that these integrals at $p^2 = 9m^2$ also have this property. PSLQ with $B(w)$ confirms it and gives

$$\begin{aligned} J_4 = & \frac{1}{\epsilon} \left(-\frac{20}{3} G_I(r_2) G_I(0, r_2) - \frac{26}{9} G_R(0, 0, 1) \right) \\ & - 16 G_I(r_2) G_R(r_4) G_I(0, r_2) + 124 G_I(r_2) G_I(0, 1, r_4) \\ & + 72 G_I(r_2) G_I(0, r_2, -1) \\ & - \frac{100}{3} G_I(0, r_2)^2 + 8 G_R(0, 0, r_4, 1) + \frac{1153 G_I(r_2)^4}{15} + O(\epsilon), \end{aligned}$$

$$\begin{aligned}
J_5 &= \frac{\sqrt{3}G_I(r_2)}{18\epsilon^3} + \frac{1}{\epsilon^2} \left(\frac{5}{9}\sqrt{3}G_I(0, r_2) - \frac{5}{9}\sqrt{3}G_I(r_2)G_R(r_4) - \sqrt{3}G_R(-1)G_I(r_2) \right) \\
&+ \frac{1}{\epsilon} \left(-\frac{52}{9}\sqrt{3}G_R(r_4)G_I(0, r_2) - 10\sqrt{3}G_R(-1)G_I(0, r_2) + \frac{40}{9}G_I(r_2)G_I(0, r_2) + 6\sqrt{3}G_I(0, r_2, -1) \right. \\
&+ \frac{26}{3}\sqrt{3}G_I(0, 1, r_4) + \frac{52}{27}G_R(0, 0, 1) + \frac{25}{9}\sqrt{3}G_I(r_2)G_R(r_4)^2 + 10\sqrt{3}G_R(-1)G_I(r_2)G_R(r_4) \\
&+ \left. 9\sqrt{3}G_R(-1)^2G_I(r_2) + \frac{253}{36}\sqrt{3}G_I(r_2)^3 \right) \\
&+ \frac{1060}{27}\sqrt{3}G_R(r_4)^2G_I(0, r_2) + \frac{32}{3}G_I(r_2)G_R(r_4)G_I(0, r_2) - 60\sqrt{3}G_R(r_4)G_I(0, r_2, -1) \\
&+ 104\sqrt{3}G_R(-1)G_R(r_4)G_I(0, r_2) + \frac{5101}{324}\sqrt{3}G_R(0, 0, 1)G_I(r_2) + 90\sqrt{3}G_R(-1)^2G_I(0, r_2) \\
&- 54\sqrt{3}G_R(-1)G_I(0, r_2, -1) + 14\sqrt{3}G_I(0, r_2)G_R(r_2, -1) - \frac{530}{9}\sqrt{3}G_R(r_4)G_I(0, 1, r_4) \\
&- 96\sqrt{3}G_R(-1)G_I(0, 1, r_4) - 60\sqrt{3}G_I(0, 1, r_2, r_3) - \frac{248}{3}G_I(r_2)G_I(0, 1, r_4) + \frac{5695}{36}\sqrt{3}G_I(r_2)^2G_I(0, r_2) \\
&- \frac{7438}{81}\sqrt{3}G_I(0, 0, 0, r_2) - 48G_I(r_2)G_I(0, r_2, -1) + \frac{200}{9}G_I(0, r_2)^2 - 74\sqrt{3}G_I(0, 1, r_2, -1) \\
&+ 54\sqrt{3}G_I(0, r_2, 1, -1) + \frac{250}{9}\sqrt{3}G_I(0, 1, 1, r_4) - \frac{16}{3}G_R(0, 0, r_4, 1) - \frac{1021}{9}\sqrt{3}G_I(r_2)^3G_R(r_4) \\
&- \frac{250}{27}\sqrt{3}G_I(r_2)G_R(r_4)^3 - 50\sqrt{3}G_R(-1)G_I(r_2)G_R(r_4)^2 - 90\sqrt{3}G_R(-1)^2G_I(r_2)G_R(r_4) \\
&- \frac{287}{2}\sqrt{3}G_R(-1)G_I(r_2)^3 - 54\sqrt{3}G_R(-1)^3G_I(r_2) - \frac{2306}{45}G_I(r_2)^4 + O(\epsilon).
\end{aligned}$$

To evaluate the ϵ -term of J_1 let us construct the following linear combination:

$$J_6 = \left(1 + \frac{1}{2}\epsilon + \frac{95}{12}\epsilon^2 + \frac{2615}{144}\epsilon^3 + \frac{1154333}{1728}\epsilon^4 \right) J_1 \\ + 48\epsilon J_4 - 3024\epsilon^3 J_5 .$$

To evaluate the ε -term of J_1 let us construct the following linear combination:

$$J_6 = \left(1 + \frac{1}{2}\varepsilon + \frac{95}{12}\varepsilon^2 + \frac{2615}{144}\varepsilon^3 + \frac{1154333}{1728}\varepsilon^4 \right) J_1 \\ + 48\varepsilon J_4 - 3024\varepsilon^3 J_5 .$$

The coefficients here are adjusted in such a way that the available result up to the finite part in ε is uniformly transcendental.

Moreover, analytical result for its ε -term can be revealed with the help of the basis

$$\tilde{B}(5) = B(5) \cup \left\{ 1, \sqrt{3}G_I(r_2), -\frac{20}{3}G_I(r_2)G_I(0, r_2) - \frac{26}{9}G_R(0, 0, 1) \right\}$$

which differs from the uniformly transcendental basis of weight 5 adding three elements proportional to the leading terms of J_1, J_5, J_4 in their ε -expansions.

$$\begin{aligned}
J_1 = & -\frac{1}{4\epsilon^4} + \frac{1}{8\epsilon^3} + \frac{1}{\epsilon^2} \left(\frac{23}{12} - \frac{3G_I(r_2)^2}{4} \right) + \frac{1}{\epsilon} \left(-\frac{1}{3} G_R(0, 0, 1) + \frac{3G_I(r_2)^2}{8} + \frac{1493}{576} \right) \\
& - 120G_I(r_2)G_R(r_4)G_I(0, r_2) + \frac{1941G_I(r_2)^4}{20} + \frac{23G_I(r_2)^2}{4} + 180G_I(r_2)G_I(0, 1, r_4) + 320G_I(r_2) \\
& G_I(0, r_2) + 72G_R(0, 0, r_4, 1) + \frac{833}{6} G_R(0, 0, 1) - 56\sqrt{3}\pi + \frac{1024805}{6912} \\
& + \epsilon \left(-1056G_I(r_2)G_R(r_4)^2G_I(0, r_2) - 2592G_R(-1)G_I(r_2)G_I(0, 1, r_4) + 828G_I(r_2)G_R(r_4)G_I(0, r_2) \right. \\
& + 1584G_I(r_2)G_R(r_4)G_I(0, 1, r_4) + 2592G_I(r_2)G_R(r_4)G_I(0, r_2, -1) - \frac{15563}{9} G_R(0, 0, 1)G_I(r_2)^2 \\
& + 1728G_I(r_2)G_I(0, r_2)G_R(r_2, -1) + 2592G_I(r_2)G_I(0, 1, r_2, r_3) - 6042G_I(r_2)G_I(0, 1, r_4) \\
& - 2880G_I(r_2)G_I(0, 1, 1, r_4) + 1704G_I(0, r_2)G_I(0, 1, r_4) - \frac{72172}{9} G_I(r_2)^3G_I(0, r_2) + \frac{320}{9} G_I(r_2)G_I(0, r_2) \\
& - 3456G_I(r_2)G_I(0, r_2, -1) + \frac{14816}{3} G_I(r_2)G_I(0, 0, 0, r_2) + 864G_I(r_2)G_I(0, 1, r_2, -1) + 1600G_I(0, r_2)^2 \\
& + 1680\sqrt{3}G_I(0, r_2) + 1136G_R(0, 0, 1, r_2, r_4) + 288G_R(r_4)G_R(0, 0, r_4, 1) - 420G_R(0, 0, r_4, 1) \\
& - 288G_R(0, 0, 1, 1, r_4) + \frac{485}{27} G_R(0, 0, 1) - \frac{397811}{405} G_R(0, 0, 0, 0, 1) + \frac{15396}{5} G_I(r_2)^4G_R(r_4) \\
& - 1680\sqrt{3}G_I(r_2)G_R(r_4) + 1512G_R(-1)G_I(r_2)^4 - 3024\sqrt{3}G_R(-1)G_I(r_2) + \frac{28000}{9}\sqrt{3}G_I(r_2) \\
& \left. - \frac{29905G_I(r_2)^4}{8} + \frac{1493G_I(r_2)^2}{192} + 28\sqrt{3}\pi + \frac{232538063}{82944} \right) + O(\epsilon^2).
\end{aligned}$$

A similar procedure is applied to J_2 and J_3 .
Two linear combinations

$$\begin{aligned} J_7 &= \left(1 + \frac{1}{3}\epsilon + \frac{37}{9}\epsilon^2 + \frac{571}{108}\epsilon^3 + \frac{139585}{324}\epsilon^4 \right) J_2 \\ &\quad - 37\epsilon J_4 + 2112\epsilon^3 J_5, \\ J_8 &= \left(1 + 8\epsilon^2 - \frac{277}{2}\epsilon^3 - \frac{29551}{12}\epsilon^4 \right) J_3 \\ &\quad + 8(6\epsilon - 1)J_4 + 16(743\epsilon + 48)\epsilon^2 J_5. \end{aligned}$$

One can also use smaller (by 20-25 percents) bases defined in terms of values of harmonic polylogarithms at sixth roots of unity

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

One can also use smaller (by 20-25 percents) bases defined in terms of values of harmonic polylogarithms at sixth roots of unity

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

ε^2 terms of the ε -expansion (weight 6) using 20000 digits?

One can also use smaller (by 20-25 percents) bases defined in terms of values of harmonic polylogarithms at sixth roots of unity

[B. A. Kniehl, A. F. Pikelner O. L. Veretin'17]

ε^2 terms of the ε -expansion (weight 6) using 20000 digits?

At least one more irreducible constant is missing?

- Using an 'elliptic' four-loop example of Feynman integrals, we have demonstrated that although we don't know analytical results for the integrals we can obtain analytical results for these integrals at singular points.

- Using an 'elliptic' four-loop example of Feynman integrals, we have demonstrated that although we don't know analytical results for the integrals we can obtain analytical results for these integrals at singular points.
- Transporting simple information about the master integrals at infinity to other singular points.

- Using an 'elliptic' four-loop example of Feynman integrals, we have demonstrated that although we don't know analytical results for the integrals we can obtain analytical results for these integrals at singular points.
- Transporting simple information about the master integrals at infinity to other singular points.
- Our algorithm works very effectively and provides high-precision numerical results, with a subsequent successful application of the PSLQ algorithm.

- Using an 'elliptic' four-loop example of Feynman integrals, we have demonstrated that although we don't know analytical results for the integrals we can obtain analytical results for these integrals at singular points.
- Transporting simple information about the master integrals at infinity to other singular points.
- Our algorithm works very effectively and provides high-precision numerical results, with a subsequent successful application of the PSLQ algorithm.
- Other applications of our algorithm are in progress.