



At The Frontiers of Particle Physics

Sergei Shmatov (MLIT JINR, Dubna)
shmatov@jinr.ru shmatov@cern.ch

MLIT, JINR, Dubna

16-20 October, 2023



Lecture 1. At The Frontiers of Particle Physics

- What does Particle Physics do?
- How does Particle Physics do?
 - ✓ Physics Tools
 - ✓ Why do we need accelerator facilities?
 - ✓ Do we need more and more new accelerator facilities?
- Where Particle Physics Frontiers are (mainly LHC examples)
 - ✓ Selected hot points of particle physics
 - ✓ Is new physics really needed?

Lecture 2. Data Analysis in High Energy Physics (18 October)

- How do we achieve results?
 - ✓ Monte Carlo tools
 - ✓ Reconstruction of physics objects
 - ✓ Reconstruction of physics processes
 - ✓ Physics Analysis and Statistics
- Something else?

What Do Particle Physicists Do?



Some eternal questions

People have long asked,

- "What is the world made of?"
- "What holds it together?"

Physicists hope to fill in their answers to these questions through the analysis of data from High Energy Physics experiments


Particle physics or high energy physics is the study of fundamental particles and forces that constitute matter (c) Wiki

- Where can I get elementary particles?
 - ✓ in Nature (cosmic sources, earth sources, i.e. natural radioactivity)
 - ✓ man-made sources (reactors, accelerators)

- How can you catch particles \Rightarrow detector facilities


- What is needed for data processing?
 - ✓ algorithms and software for reconstruction of physics objects and processes

- What is needed for data analysis?
 - ✓ Theory
 - ✓ Monte Carlo Tools
 - ✓ Statistics Tools



Along Three Paths


Information Technologies



Accelerators : powerful machines that accelerate particles to extremely high energies and bring them into collision with other particles

Detectors : gigantic instruments that record the resulting particles as they “stream” out from the point of collision.

Computing : to collect, store, distribute and analyse the vast amount of data produced by these detectors




It's been a global effort, a global success. It has only been possible because of the extraordinary achievements of the experiments, infrastructure and the grid computing“
(c) Rolf Heuer, the Director General of CERN, when the discovery of the Higgs

Collaborative Science on Worldwide scale : thousands of scientists, engineers, technicians and support staff to design, build and operate these complex “machines”.





- The Energy Frontier**, using high-energy colliders to discover new particles and directly probe the architecture of the fundamental forces.
- The Intensity Frontier**, using intense particle beams to uncover properties of neutrinos and observe rare processes that will tell us about new physics beyond the Standard Model.
- The Cosmic Frontier**, using underground experiments and telescopes, both ground and space based, to reveal the natures of dark matter and dark energy and using high-energy particles from space to probe new phenomena.



JINR LONG-TERM DEVELOPMENT STRATEGIC PLAN UP TO 2030 AND BEYOND

- RELATIVISTIC HEAVY-ION PHYSICS AT NICA
- JINR PARTICIPATION IN FOREFRONT EXTERNAL EXPERIMENTS OFF-SITE
 - LHC, SPS, RHIC, and at facilities under construction, as for example the FAIR facility
- NICA SPIN PHYSICS
- PARTICLE PHYSICS AT THE LHC AND BEYOND
 - Accelerator-based research and frontier accelerator technologies (LHC, SPS, NICA, FAIR, etc)
 - Neutrino physics and astroparticle physics (Baikal-GVD, JUNO, NOvA, DUNE, etc)
 - Multi-messenger astronomy including gravitational wave detection (Baikal-GVD, TAIGA, VIRGO, etc)



Baikal-GVD (Gigaton Volume Detector)

Why High Energy is Needed?



Particle physics have focused on the inner space frontier, pursuing the questions of the construction of matter and the fundamental forces at the smallest scale accessible.



Wavelength of probe radiation should be smaller than the object to be resolved

$$\lambda \ll \frac{h}{p} = \frac{hc}{E}$$


Object	Size	Energy of Radiation
Atom	10 ⁻¹⁰ m	0.00001 GeV (electrons)
Nucleus	10 ⁻¹⁴ m	0.01 GeV (alphas)
Nucleon	10 ⁻¹⁵ m	0.1 GeV (electrons)
Quarks	?	> 1 GeV (electrons)

Radioactive sources give energies in the range of MeV

Need accelerators for higher energies.



"electronic eyes"





- **Luminosity L** is a measure of how many interactions of **cross section s** can be created per unit time


$$L\sigma = \frac{dN}{dt} \quad N = \sigma \int L dt = \sigma L_{\text{int}}$$

- L_{int} is integrated luminosity, an important factor of production for colliders
- $[L] = \text{cm}^{-2} \text{s}^{-1}$, $[L_{\text{int}}] = \text{cm}^{-2}$ (1 barn = 10^{-24}cm^2 ; $1 \text{pb}^{-1} = 10^{36} \text{cm}^{-2}$)

- For equal-sized head-on Gaussian beams in a collider

$$L = \frac{f_{\text{rev}} h N_1 N_2}{4\pi\sigma_x\sigma_y}$$

- $\sigma_{x,y}$ are rms beam sizes, h is number of bunches



Colliding 100 mm 7.5×10^9 proton bunches at 100 kHz for 1 year gives about 1pb^{-1} of integrated luminosity



Highest energies can be reached with proton colliders

Machine	Year	Beams	Energy (\sqrt{s})	Luminosity
SPPS (CERN)	1981	pp	630-900 GeV	$6 \cdot 10^{30} \text{cm}^{-2} \text{s}^{-1}$
Tevatron (FNAL)	1987	pp	1800-2000 GeV	$10^{31}-10^{32} \text{cm}^{-2} \text{s}^{-1}$
SLC (SLAC)	1989	e^+e^-	90 GeV	$10^{30} \text{cm}^{-2} \text{s}^{-1}$
LEP (CERN)	1989	e^+e^-	90-200 GeV	$10^{31}-10^{32} \text{cm}^{-2} \text{s}^{-1}$
HERA (DESY)	1992	ep	300 GeV	$10^{31}-10^{32} \text{cm}^{-2} \text{s}^{-1}$
RHIC (BNL)	2000	pp / AA	200-500 GeV	$10^{32} \text{cm}^{-2} \text{s}^{-1}$
LHC (CERN)	2009	pp (AA)	7-14 TeV	$10^{33}-10^{34} \text{cm}^{-2} \text{s}^{-1}$


Luminosity = number of events/cross section/sec

Limits on circular machines (except the expense)


- Proton colliders: Dipole magnet strength
→ superconducting magnets
- Electron colliders: Synchrotron radiation/RF power

Limits on linear machines

- the device's length



Do we really need more and more accelerator facilities?





1900s: e discovered (cathode ray tube)
 γ interpreted as a particle
 1930s: μ discovered (cosmic rays)
 1950s: ν_e observed (nuclear reactor)
 ν_μ discovered (BNL)
 1960s: 1st evidence for quarks
 u and d observed (SLAC)
 s observed (BNL)
 1970s: *standard model is born*
 c discovered (SLAC, BNL)
 τ observed (SLAC)
 b observed (FNAL)
 1980s: W and Z observed (CERN)
 1990s: t quark observed (FNAL)
 2000s: ν_τ observed (FNAL)

Three Generations of Matter (Fermions)


	I	II	III	
mass →	2.4 MeV	1.27 GeV	171.2 GeV	0
charge →	$\frac{2}{3}$	$\frac{2}{3}$	$\frac{2}{3}$	0
spin →	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
name →	u up	c charm	t top	γ photon
Quarks	4.8 MeV	104 MeV	4.2 GeV	0
	$-\frac{1}{3}$	$-\frac{1}{3}$	$-\frac{1}{3}$	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	d down	s strange	b bottom	g gluon
Leptons	<2.2 eV	<0.17 MeV	<15.5 MeV	91.2 GeV
	0	0	0	0
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	ν_e electron neutrino	ν_μ muon neutrino	ν_τ tau neutrino	Z⁰ weak force
	0.511 MeV	105.7 MeV	1.777 GeV	80.4 GeV
	-1	-1	-1	± 1
	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1
	e electron	μ muon	τ tau	W[±] weak force

Bosons (Forces)

SM is fully completed, expect for Higgs boson



- What about a mass generation mechanism?
 - ✓ Higgs bosons? (Now – Yes)
- Three generation of particles:
 - ✓ why 3 (they can not be fixed in the framework of SM)?
 - ✓ why do we need all of them?




Quarks	u up	c charm	t top
	d down	s strange	b bottom
Leptons	ν_e e- Neutrino	ν_μ μ - Neutrino	ν_τ τ - Neutrino
	e electron	μ muon	τ tau
	I	II	III
The Generations of Matter			

- Yukawa hierachy (explanation of mass patterns for quarks and leptons)
- A lot of free parameters: gauge couplings, Yukawa coupling constants, CKM-mixing angles, Higgs vacuum expectation value, etc (in total, 26)



... why do we believe in something beyond Standard Model?




- Where is a gravity?
- Hierarchy problem
 - fine tuning of higgs mass is needed to “neutralize” contribution from high order corrections
 - huge gap between Electroweak (10^2 GeV) and Planck scale (10^{19} GeV) scales), Gravity/EW $\sim 10^{19}/10^2$ GeV?



Unification of Forces



- Dark Matter: what does it consist of?
- CP-violation in Early Universe




Well, the Standard Model is not a ultimate theory!



1. What is dark matter ?
2. What is dark energy ?
3. How were the heavy elements from iron to uranium made ?
4. Do neutrinos have mass ?
5. Where do ultrahigh-energy particles come from ?
6. Is a new theory of light and matter needed to explain what happens at very high energies and temperatures ?
7. Are there new state of matter at ultrahigh temperate and dentensity ?
8. Are protons unstable ?
9. What is gravity ?
10. Are there additional dimensions ?
11. How did the universe begin ?

What do we hope to see beyond the SM?



How we can escape beyond the Standard Model $SU(3)_c \times SU(2) \times U(1)$?





- Simplest extension of SM (based on SM gauge group)
 - ✓ 4 generation of fermions, q^* , l^* ...
- Extended gauge sector
 - ✓ ExQCD (colorons, axigluons, diquarks etc)
 - ✓ ExEW (W' , Z' , ...)
 - ✓ GUT (leptoquarks)
- Extended Higgs Sector
 - ✓ Higgs Doublet Models (HDM)
 - ✓ Higgs in ED, Higgs-Radion mixing, composite Higgs (type)
- Hierarchy problem
 - ✓ SUSY (s-particles, LSP из RPV/split/GM SUSY...)
 - ✓ Extra dimensions (ED)
 - KK-modes of particles SM, KKPV, FCNC...
 - microscopic black holes (semi-classical, string balls, quantum black holes)
 - ✓ Technicolor (technibosons and technifermions, leptoquarks ...)
 - ✓ Compositeness
- Dark Matter (EFT)

Will be discussed in details someday


How do we intend to observe all this (which accelerator is better?)?

Types of particle accelerators

- ✓ linear and circular accelerator
- ✓ e^+e^- , ep , pp , $ppbar$, pA , AA , ...




First High Energy Physics Experiments:
Beam on fixed target!



Rutherford experiment (1909)


High Energy Physics Experiments since mid 70's:
Colliding beams!



Centre of mass energy squared $s=2E_1m_2$

Centre of mass energy squared $s=4E_1E_2$

...plus secondary beams such as neutrinos...




academic 11 3/6/98 slide 2

Electron-Positron Collider: clean signature

Synchrotron Radiation:

$$P = \frac{2 e^2 c}{3 R^2} \left(\frac{E}{m c^2} \right)^4$$



CERN LEP : $R=4.5\text{km}$, $E_{\text{beam}} \sim 100 \text{ GeV}$

CERN LHC: $R=4.5\text{km}$, $E_{\text{beam}} \sim 7000 \text{ GeV}$

$$\frac{\Delta E(e)}{\Delta E(p)} = \left(\frac{m_p}{m_e} \right)^4 \sim 10^{13}$$



- The simplest answer is .. as much as possible (construction restriction, technologies, price..)
- Different approach is .. more than Tevatron energy of 1.8 TeV (vague claim)

- Real life

- ✓ Higgs boson

- $m_H < 1$ TeV from theory (SM unitarity requires)
 - $m_H > 114.1$ GeV from LEP
 - $156 < m_H < 177$ GeV from Tevatron



need a machine to discover/exclude Higgs from ≈ 120 GeV to 1 TeV

- ✓ $m_{SUSY} \sim$ TeV



need a machine to explore a range of up to a several TeV

- ✓ $m_{BSM} > 0.5$ TeV from theory and from Tevatron



need a energy scan



Factorization hypothesis



Fragmentation function

$$E \frac{d^3 \sigma}{dp^3} \propto f_{a/A}(x_a, Q^2) \otimes f_{b/B}(x_b, Q^2) \otimes \frac{d\hat{\sigma}^{ab \rightarrow cd}}{dt} \otimes D_{h/c}(z_c, Q^2)$$

Parton distributions

Parton density functions (PDFs)

cross section of hard processes




effective centre-of-mass energy $\sqrt{\hat{s}}$ smaller than \sqrt{s} of colliding beams:

$$\left. \begin{aligned} \vec{p}_a &= x_a \vec{p}_A \\ \vec{p}_b &= x_b \vec{p}_B \end{aligned} \right\} p_A = p_B = 7 \text{ TeV} \quad \sqrt{\hat{s}} = \sqrt{x_a x_b s} \approx x \sqrt{s}$$

if $x_a \approx x_b$

→ to produce $m \approx 100 \text{ GeV}$ $x \sim 0.01$
to produce $m \approx 5 \text{ TeV}$ $x \sim 0.35$




Carlo Rubbia

Giorgio Brianti



1984 (Lausanne): the first LHC Working Group for LHC conception

December 16, 1991: LHC Project was approved by CERN Council



- **High Energy** \Rightarrow factor 7 increase w.r.t. past accelerators
- **High Luminosity** (# events/cross section/time) \Rightarrow factor 100 increase

Linac: 50 MeV \rightarrow PSB: 1.4 GeV \rightarrow PS: 28 GeV \rightarrow SPS: 450 GeV \rightarrow LHC: 7 TeV






Quantity	number
Circumference	26 659 m
Dipole operating temperature	1.9 K (-271.3°C)
Number of magnets	9593
Number of main dipoles	1232
Number of main quadrupoles	392
Number of RF cavities	8 per beam
Nominal energy, protons	7 TeV
Nominal energy, ions	2.76 TeV/u (*)
Peak magnetic dipole field	8.33 T
Min. distance between bunches	~7 m
Design luminosity	$10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
No. of bunches per proton beam	2808
No. of protons per bunch (at start)	1.1×10^{11}
Number of turns per second	11 245
Number of collisions per second	600 million

(*) Energy per nucleon

LHC is **100m** underground
 LHC is **27 km** long
 Magnet Temperature is **1.9 Kelvin** = -271 Celsius
 LHC has ~ **9000 magnets**
 LHC: **40 million** proton-proton collisions per second
 LHC: Luminosity **100 fb⁻¹/year** (after start-up phase)




The Large Hadron Collider Project 4 detectors





Onion structure of detector layers placed in B-field

type	tracking	ECAL	HCAL	MUON
γ				
e				
μ				
Jet				
Et miss				



Each layer identifies and measures (or remeasures) the energy of particles unmeasured by the previous layer

No single detector can determine identity and measure energies/momenta of all particles



LHC detectors must have

- Fast response, otherwise too large pile-up. Typical response time 20-50 ns
 - pile-up of 25-50 minimum bias events
 - very challenging readout electronics
- high granularity to minimize probability that pile-up particles be in the same detector element as interesting object
 - large number of electronic channels, high cost
- a robust and redundant Muon system
- the best possible e/g calorimeter ECAL that consistent with Muon System
- a highly efficient Tracking system consistent that with Muon System and ECAL
- a hermetic calorimeter system
- high radiation resistant e.g. in forward calorimeters: up to 10^{17} n/cm² in 10 years of LHC operation
- good PID (particle identification)
- good E, p_T resolution

Precision Muon Spectrometer $\sigma / p_T \sim 10\% \text{ at } 1 \text{ TeV}/c$
 Fast response for trigger
 Good p resolution (e.g., $A/Z' \rightarrow \mu\mu$, $H \rightarrow 4\mu$)



EM Calorimeters $\sigma / E \sim 10\% / \sqrt{E(\text{GeV})}$
 excellent electron/photon identification
 Good E resolution (e.g., $H \rightarrow \gamma\gamma$)

Hadron Calorimeters
 Good jet and E_T miss performance
 (e.g., $H \rightarrow \tau\tau$) $\sigma / E \sim 50\% / \sqrt{E(\text{GeV})} \oplus 0.03$


Inner Detector
 $\sigma / p_T \sim 5 \cdot 10^{-4} p_T \oplus 0.001$
 Good impact parameter res.
 (e.g., $H \rightarrow bb$)




Example : $H \rightarrow \gamma\gamma$



number and p_T of hadrons in a jet have large fluctuations




in some cases: one high- p_T π^0 ; all other particles too soft to be detected




$d(\gamma\gamma) < 10$ mm in calorimeter \rightarrow QCD jets can mimic photons. Rare cases, however:

$$\frac{\sigma_{jj}}{\sigma(H \rightarrow \gamma\gamma)} \sim 10^8 \quad m_{\gamma\gamma} \sim 100 \text{ GeV}$$



ATLAS

<i>Weight</i>	<i>7000 t</i>
<i>Diameter</i>	<i>25 m</i>
<i>Length of toroid</i>	<i>26 m</i>
<i>Total Length</i>	<i>46 m</i>
<i>B-field</i>	<i>2 Tesla</i>



CMS


**CMS
A Compact Solenoidal Detector for LHC**

Total weight : 12,500t.
Overall diameter : 15.00m
Overall length : 21.60m
Magnetic field : 4 Tesla


CMS-PARA-001-11/07/97 JLB,PP

<i>Weight</i>	<i>12 500 t</i>
<i>Diameter</i>	<i>15 m</i>
<i>Length</i>	<i>21.6 m</i>
<i>B-field</i>	<i>4 Tesla</i>

Detector systems are designed to measure:
energy and momentum of photons, electrons, muons, and jets up to a few TeV



What do we know today about the Standard Model from LHC?



■ SM processes:

$\sigma \sim 1/(100 \text{ MeV})^2$



10⁻⁸ !

■ New Physics:

$\sigma \sim 1/(1 \text{ TeV})^2$

During Run 2 the LHC produced 10¹⁶ collisions

Large samples of various particles produced:

- W bosons: 12 billion
- Z bosons: 2.8 billion
- Top quarks: 300 million
- Bottom quarks: 40 trillion
- Higgs bosons: 7.7 million




Summaries of CMS cross section measurements

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsCombined>

Bosons

plots are updated for Summer 2023 Conferences

Overview of CMS cross section results





Higgs Physics

From design



to discovery



4 July 2012

Higgs announcement at CERN






	Int. Luminosity at 7, 8 TeV	m _H [GeV]	Expected [st. dev.]	Observed [st. dev.]
ATLAS	10.7 fb ⁻¹	126.0 ± 0.6	4.6	5.0
CMS	10.4 fb ⁻¹	125.3 ± 0.6	5.9	4.9




During Run 2 of the LHC the experimental collaborations started to employ the combined data for precision measurements of Higgs properties (mass, width, couplings, CP, rare decays)

- All main production mechanisms are observed, including $h \rightarrow b\bar{b}$, $t\bar{t}H$, VH
- Mass of Higgs boson m_h is measured with an accuracy of 0.1% (!)



- Precisions of cross section and branching ratio measurements in combined channel are down to 8.5% level
- We have ~6-30% accuracy for measurements of couplings
- The absolute value of a width $\Gamma_H = 3.2^{+2.4}_{-1.7}$ MeV is getting closer to the SM expectations (4.1 MeV). We still need to improve an accuracy.
- Spin, parity, differential distributions do not contradict the SM






**THE STANDARD MODEL: IT HAS TO BREAK DOWN
AT SOME POINT BUT JUST KEEPS CHUGGING ALONG!**

MCK, COSPA2014

Why we are still expecting the New Physics?






... but the current accuracy of Higgs coupling measurements is still insufficient to reject BSM Higgs hypothesis

EPJC 79 (2019) 421

Summary of Higgs Couplings Measurements

Model	κ_V	κ_b	κ_γ
Singlet Mixing	$\sim 6\%$	$\sim 6\%$	$\sim 6\%$
2HDM	$\sim 1\%$	$\sim 10\%$	$\sim 1\%$
ecoupling MSSM	$\sim -0.0013\%$	$\sim 1.6\%$	$\sim -4\%$
Composite	$\sim -3\%$	$\sim -(3-9)\%$	$\sim -9\%$
Top Partner	$\sim -2\%$	$\sim -2\%$	$\sim +1\%$




The 95% CL upper limit on B_{Undet} is found to be < 0.16

- Measurements precision continuous increasing
- Search for new higgs states and other BSM
 - ✓ BSM effects on in the differential distributions



The properties of the Higgs h_{125} agree fully with SM in decay into

- gauge bosons
- 3rd generation fermions (t/b/ τ)
- and do not conflict with results for the 2nd generation (no deviations in $cc/\mu\mu$ decays after RUN2)




$$\mathcal{B}_{SM}(H \rightarrow e^+e^-) \approx 9 \cdot 10^{-9}$$

$$\mathcal{B}_{SM}(H \rightarrow \mu^+\mu^-) \approx 2,2 \cdot 10^{-4}$$



$\mu\mu: 3\sigma$
 $1.19^{+0.40}_{-0.39} \text{ (stat)}^{+0.15}_{-0.14} \text{ (syst)}$
JHEP 01 (2021) 148




We do not know and will not know until the end of the LHC whether the coupling of the Higgs h_{125} to 1st generation fermions is in a “standard” way or not.

If we have no Extra Higgses! (rare decays are enhanced within Extended Higgs Sectors)






$$R_X \equiv \frac{\mathcal{B}(B \rightarrow X \mu^+ \mu^-)}{\mathcal{B}(B \rightarrow X e^+ e^-)} = 1 \text{ in the SM}$$

All QCD effects cancel in ratios.
Small $\mathcal{O}(1\%)$ radiative corrections.



[arXiv:2103.11769](https://arxiv.org/abs/2103.11769)
Nature 18, 277 (2022)



$$R_K(1.1 < q^2 < 6.0 \text{ GeV}^2 \text{ c}^{-4}) = 0.846^{+0.042+0.013}_{-0.039-0.012}$$

Control uncertainties by measuring double ratios:

$$R_X \equiv \frac{\mathcal{B}(B \rightarrow X \mu \mu)}{\mathcal{B}(B \rightarrow X J/\psi (\rightarrow \mu \mu))} \frac{\mathcal{B}(B \rightarrow X J/\psi (\rightarrow ee))}{\mathcal{B}(B \rightarrow X ee)} = 1_{(SM)}$$

► LHCb searches in B decays (90% CL limits):



$$\mathcal{B}(B_s^0 \rightarrow e \mu) < 5.4 \times 10^{-9} \quad [3 \text{ fb}^{-1} \text{ JHEP03(2018)078}]$$

$$\mathcal{B}(B^+ \rightarrow K e \mu^-) < 7 \times 10^{-9} \quad [3 \text{ fb}^{-1} \text{ hep-ex/1909.01010}]$$

$$\mathcal{B}(B_s^0 \rightarrow \tau \mu) < 3.4 \times 10^{-5} \quad [3 \text{ fb}^{-1} \text{ hep-ex/1905.06614}]$$





$W \rightarrow \mu\nu$ and $W \rightarrow e\nu$ decays.



Science Vol 376, 170-176 (2022)



$$M_W = 80,433.5 \pm 6.4_{\text{stat}} \pm 6.9_{\text{syst}} = 80,433.5 \pm 9.4 \text{ MeV}/c^2$$



for $J=1/2$
 $g = \mu/\mu_B J = 2$
 $a_\mu = (g - 2)/2$

$$a_\mu = a_\mu^{QED} + a_\mu^{Had} + a_\mu^{Weak} + a_\mu^{NewPhysics}$$

$$1,000,000 : 60 : 1.3 : \propto (m_\mu/m_X)^2$$



$$a_\mu^{Th} [2020] = 116\,591\,810(43) \times 10^{-11} \text{ (0.37 ppm)}$$

$$a_\mu^{Exp} [2021] = 116\,592\,061(41) \times 10^{-11} \text{ (0.35 ppm)}$$

$$a_\mu^{Exp} - a_\mu^{Th} = (251 \pm 59) \times 10^{-11} \text{ (4.2}\sigma\text{)}$$


The new experimental result is:
 $g-2 = 0.00233184110 \pm 0.00000000043 \text{ (stat.)} \pm 0.00000000019 \text{ (syst.)}$, 0.2 ppm



Direct Searches for the Physics Beyond the SM

- ✓ Conventional Signals, such as new resonances in dileptons/diphotons/dijets spectra or non-resonant signals, combinations of physics objects (leptons/photons/jets) and MET/ b/t-jets tags, high-multiplicity events, etc

SUSY Extended Gauge Sector Extra Dimensions CI/Excited Fermions/B3G



- ✓ Non-conventional Signals, for example displaced vertices/leptons/lepton-jets/dileptons from Long-Lived Particles or emerging jets/leptons from boosted heavy objects, $m \ll p_T$ (i.e. high- p_T Z/W/h₁₂₅ bosons)

Long-Lived Particles (Dark Matter/Non-standard SUSY/Neutrino Masses/etc)

Extended Higgs and Dark Matter Sectors

BSM-Higgs Physics

- ✓ Searches for the new Higgs states (from extended Higgs sector including SUSY)
- ✓ Probes for the New Physics with h₁₂₅ (Higgs as a tool for new discovery)




Extra Higgses, Dark Matter, Flavour Universality Violation

Precision Tests of SM


- ✓ Measurements of the W/Z, Drell-Yan (+ n jets) x-sections and angular characteristics
- ✓ Search for rare decays of B-mesons
- ✓ Observations of other rare process in top sector within SM (Wtb couplings, CP violating top quark couplings, flavor-changing neutral current interactions of the t-quark and h₁₂₅)



- Heavy Resonances (extended gauge models, extra dimensions, technicolor) \Rightarrow dileptons, dijets, diphotons, $t\bar{t}$, WZ
- Non-Resonant Signals
- Mono-particle + Missing ET (extended gauge models, extra dimensions, technicolor, SUSY) \Rightarrow mono-jet + MET, mono-photon + MET, mono-lepton + MET
- Microscopic Black Holes (extra dimensions) \Rightarrow high-multiplicity events




- Leptoquarks \Rightarrow lepton + jet
- 4th Generation \Rightarrow leptons/jets, dilepton






Overview of CMS EXO results





New Physics ($Z'/Z_{KK}/G_{KK}$) contributions to SM processes









$$R_\sigma = \frac{\sigma(pp \rightarrow Z' + X \rightarrow l^+l^- + X)}{\sigma(pp \rightarrow Z^0 + X \rightarrow l^+l^- + X)}$$

Dileptons, full RUN2 data

JHEP 07 (2021) 208

$W\gamma$, full RUN2 data PLB 826 (2022) 136888



benchmark heavy scalar (vector) triplet bosons with masses between 0.75 (1.15) and 1.40 (1.36) TeV are excluded at 95% CL

What does Brazilian Flag mean?



Dimuon example

$$R_\sigma = \frac{\sigma(pp \rightarrow Z' + X \rightarrow l^+l^- + X)}{\sigma(pp \rightarrow Z^0 + X \rightarrow l^+l^- + X)}$$



Extended gauge models

Models of low-energy gravity (RS1-type scenario of ED)

SM predictions

BSM predictions

SM predictions



Model-independent limits on cross section (in narrow width approximation, NWA)

Channel	Z'_{SSM}		Z'_{ψ}	
	Obs. [TeV]	Exp. [TeV]	Obs. [TeV]	Exp. [TeV]
$e e$	4.72	4.72	4.11	4.13
$\mu^+ \mu^-$	4.89	4.90	4.29	4.30
$e e + \mu^+ \mu^-$	5.15	5.14	4.56	4.55


Channel	$k/\bar{M}_{Pl} = 0.01$		$k/\bar{M}_{Pl} = 0.05$		$k/\bar{M}_{Pl} = 0.1$	
	Obs. [TeV]	Exp. [TeV]	Obs. [TeV]	Exp. [TeV]	Obs. [TeV]	Exp. [TeV]
$e e$	2.16	2.29	3.70	3.83	4.42	4.43
$\mu^+ \mu^-$	2.34	2.32	3.96	3.96	4.59	4.59
$e e + \mu^+ \mu^-$	2.47	2.53	4.16	4.19	4.78	4.81

Example of Dark Matter Searches in Dijets + Dileptons



We consider a model that assumes the existence of a single DM particle that interacts with the SM particles through a spin-1 mediator, which can be either a vector or axial-vector boson.









- vector mediator with small couplings to leptons, $g_{DM} = 1.0$, $g_q = 0.1$, $g_l = 0.01$
- axial-vector mediator with equal couplings to quark and leptons: $g_{DM} = 1.0$, $g_q = g_l = 0.1$



5 parameters:

m_{DM} , m_{Med} , g_{DM} , g_l , g_q

DM-nucleon upper limits on the cross section







First intrigue @ 13 TeV: Excess in Diphotons

CMS-PAS-EXO-16-018

CMS sees excess in diphoton spectrum around ~ 750 GeV on 8+13 TeV data

✓ 3.4σ (local) and 1.6σ (global) for analysis of spin-0 and spin-2 states:




A lot of theoretical works attempt to explain this effect:
more 320 papers since Dec. 2015


Need more data!




• Light $X \rightarrow \gamma\gamma$



• Light $X \rightarrow \tau\tau$



• Light $X \rightarrow \mu\mu$



A. Nikitenko




RUN3 is a perfect judge for these challenges!

Higgs Boson as a Tool to Search for the New Physics


THE HIGGS
BOSON
EXPLAINED




H_{125} exotic decays



Direct Search for BSM: LLP Non-conventional Signals



e.g. for $c\tau = 5$ cm, $\langle\beta\gamma\rangle \sim 30$





LLPs may have decay lengths up to several meters, hence traveling through the inner detector layers without leaving any trace

- a proper lifetime $c\tau_0$ is greater than or comparable to the characteristic size of the (sub)detectors
- small $c\tau_0$ that comparable to the inner tracker size, no displaced tracks \rightarrow "standard" prompt decay
- intermediate $c\tau_0 \rightarrow$ LLP
- very large/infinite large $c\tau_0 \rightarrow$ stable particles, "standard" MET signatures




- inelastic dark matter: relic particles that cannot scatter elastically off of nuclei the dark sector
- particles continue traveling for a long time and traverse several meters (Long-Lived Particles) before tunneling back into our visible universe (quarks or leptons)



Motivation	Top-down Theory	IR LLP Scenario
Naturalness	RPV SUSY GMSB mini-split SUSY Stealth SUSY Axinos Sgoldstinos	BSM \rightarrow LLP <i>(direct production of BSM state at LHC that is or decays to LLP)</i>
Dark Matter	Neutral Naturalness Composite Higgs Relaxion	Hidden Valley ALP SM+S SM+V(+S) exotic Z decays exotic Higgs decays exotic Hadron decays
Baryogenesis	Asymmetric DM Freeze-In DM SIMP/ELDER Co-Decay Co-Annihilation Dynamical DM	HNI
Neutrino Masses	Minimal RH Neutrino with $U(1)_{B-L} Z'$ with $SU(2)_R W_R$ long-lived scalars with Higgs portal from ERS Discrete Symmetries	

<https://arxiv.org/abs/1901.04040>




M. Lisanti

LHC Prospects and beyond






The Present and the Future






LHC dedicated LLP experiments





Phys. Rev. D 99 (2019) 015021

<https://arxiv.org/abs/1901.04040>


CMS/ATLAS




CMS




LHCb



ATLAS



- $m_X = 5 \text{ GeV}$
- $m_X = 20 \text{ GeV}$
- $m_X = 40 \text{ GeV}$
- MATHUSLA
- ATLAS



Future Circular Colliders (100 TeV pp)






Table 1 | Run plan for the FCC-ee

FCC-ee phase	Run duration (yr)	\sqrt{s} (GeV)	L_{int} (ab ⁻¹)	Event statistics
Z ⁰	4	88-95	150	3 × 10 ¹² hadronic Z ⁰ decays
W ⁺ W ⁻	2	158-192	12	3 × 10 ⁸ W ⁺ W ⁻ pairs
Z ⁰ H	3	240	5	10 ⁶ Z ⁰ H events
t \bar{t}	5	345-365	1.5	10 ⁶ t \bar{t} and 6 × 10 ⁴ H $\nu\bar{\nu}$ events
H (optional)	3	125	21	Optional run on H resonance




$$\sqrt{s} = 100 \text{ TeV}$$

The launch point is 2020





Compact Linear Collider (CLIC)



	Collision energy	Integrated luminosity (unpolarized beams)
1st stage	380 GeV	1.0 ab ⁻¹
2nd stage	1500 GeV	2.5 ab ⁻¹
3rd stage	3000 GeV	5.0 ab ⁻¹


Circular Electron Positron Collider (CEPC)



	Collision energy	Integrated luminosity (unpolarized beams)
1st stage	90 GeV	16 ab ⁻¹
2nd stage	180 GeV	2.6 ab ⁻¹
3rd stage	240 GeV	5.6 ab ⁻¹


(CEPC Study Group, arXiv:1809.00285, 1811.10545)

International Linear Collider (ILC)



	Collision energy	Integrated luminosity (unpolarized beams)
1st stage	250 GeV	2.0 ab ⁻¹
2nd stage	500 GeV	4.0 ab ⁻¹
3rd stage	1000 GeV	5.4 ab ⁻¹

(ILC Technical Design Report, arXiv:1306.6327, 1903.01629)




Higgs boson is found



shutterstock.com · 264969203

Standard Model works




Extensive Searches for New Physics


- No significant signals
- A set of hints
- A number of future projects



Particle physics isn't going to die — even if the LHC finds no new particles







Anyway...




The Hitchhiker's Guide to the Galaxy
by Douglas Adams

Оорт первый взглянул на звездное небо и заметил, что Галактика вращается

(с) Г. Проницательный



THANK YOU FOR YOUR ATTENTION!



The ATLAS Experiment




Muon Detector

- air-core toroids, MDT+RPC+TGC
- $\sigma/p_T \sim 2-7\%$
- $|\eta| < 2.7$,
- $|\eta| < 2.5$

EM Calorimetry

- Pb-LAr
- $\sigma/E \sim 10\%/\sqrt{E(\text{GeV})} \oplus 1\%$
- $|\eta| < 3.2$, $|\eta| < 2.5$ (fine granularity)

Length : ~ 46 m
 Radius : ~ 12 m
 Weight : ~ 7000 tons
 Channels: $\sim 10^8$
 L_{cable} : ~ 3000 km



Inner detector

- Si pixels and strips
- Transition Radiation Detector (e/π separation)
- $\sigma/p_T \sim 0.05\% p_T(\text{GeV}) \oplus 0.1\%$;
- $|\eta| < 2.5$, $B=2$ T(central solenoid)

Hadron Calorimeter

- Fe/scintillator (central), Cu/W-LAr (fwd)
- $\sigma/E \sim 50\%/\sqrt{E(\text{GeV})} \oplus 3\%$
- $|\eta| < 3$



CMS DETECTOR

Total weight : 14,000 tonnes
Overall diameter : 15.0 m
Overall length : 28.7 m
Magnetic field : 3.8 T

STEEL RETURN YOKE
12,500 tonnes

SILICON TRACKERS
Pixel (100x150 μm) $\sim 16\text{m}^2 \sim 66\text{M}$ channels
Microstrips (80x180 μm) $\sim 200\text{m}^2 \sim 9.6\text{M}$ channels

$$\frac{\delta p_T}{p_T} \sim 1.0 - 1.5\% @ 100\text{GeV}$$

SUPERCONDUCTING SOLENOID
Niobium titanium coil carrying $\sim 18,000\text{A}$

MUON CHAMBERS
Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers

PRESHOWER
Silicon strips $\sim 16\text{m}^2 \sim 137,000$ channels


FORWARD CALORIMETER
Steel + Quartz fibres $\sim 2,000$ Channels


$$\frac{\sigma_E}{E} = \frac{2.7\%}{\sqrt{E}} \oplus 0.5\% \oplus \frac{0.15}{E}$$


CRYSTAL ELECTROMAGNETIC CALORIMETER (ECAL)
 $\sim 76,000$ scintillating PbWO_4 crystals

$$\frac{\sigma_E}{E} = \frac{120}{\sqrt{E}} \oplus 5\%$$

HADRON CALORIMETER (HCAL)
Brass + Plastic scintillator $\sim 7,000$ channels










CMS Luminosity Information

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/LumiPublicResults>



LHC / HL-LHC Plan





$$L_{inst} = 2.1 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$$



We are here

$$L_{nisy} = 7.5 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$$


- CMS Dataset RUN2**
 - ✓ ~163 fb⁻¹ of proton-proton collisions @ 13 TeV is delivered
 - ✓ 151.78 fb⁻¹ is recorded by CMS (data-taking efficiency ~93%)
- CMS Dataset RUN3**
 - ✓ ~73.4 fb⁻¹ is already delivered @ 13.6 TeV during the RUN3
 - ✓ 63.7 fb⁻¹ is recorded by CMS (data-taking efficiency ~92%)
 - ✓ ~93% of collected data “good for physics” in 2023 (91% in 2022)
 - ✓ number of pp interactions per beam crossing (PU): $\langle \mu \rangle = 52$ for 2023
- ~260 fb⁻¹ it is expected @ 13.6 TeV for the end of the RUN3 (450 fb⁻¹, in total for RUN1/2/3)
- pPb and PbPb Runs (see talk by Serguei Petrushanko)



CMS Data Quality Information

<https://twiki.cern.ch/twiki/bin/view/CMSPublic/DataQuality>

The ALICE Experiment










MoEDAL: Monopole and Exotics Detector at the LHC



Heavy particles which carry “magnetic charge”
Could eg explain why particles have “integer electric charge”


Monopole production




Remove the sheets after some running time and inspect for ‘holes’





TOTEM: measuring the total, elastic and diffractive cross sections
 Add Roman pots (and inelastic telescope) to CMS interaction regions (200 m from IP)
 Common runs with CMS planned



TOTAL and Elastic cross section Measurement





LHCf: measurement of photons and neutral pions in the very forward region of LHC
 Add a EM calorimeter at 140 m from the Interaction Point (of ATLAS)



Connection with cosmic rays





10 September 2008, 9:50, the first LHC beam event was recorded by CMS



The first collisions (3.5 TeV + 3.5 TeV) were happen on March 30th, 2010, at 13-00 (Geneve)

12:52 – CMS, 12:58 – ATLAS, 12:59 – LHCb, 13:01 – ALICE



14:30 Neutral pion decay was detected by CMS