

Measurement and analysis of the $^{13}\text{C}(\alpha, \alpha^0)^{13}\text{C}$ reaction cross-section in the energy range of 2.0 – 7.0 MeV

Tuesday 2 July 2024 18:40 (15 minutes)

The differential cross-sections of the $^{13}\text{C}(\alpha, \alpha^0)^{13}\text{C}$ reaction were measured at three angles (130° , 150° and 170°) in the energy range of 2.0-7.0 MeV. The thin layer of ^{13}C deposited to the beryllium backing was used as a target. The thickness and enrichment of the target were determined by the ion beam analysis methods. The effect of the carbon build-up was taken into account during the data analysis. The measurement results can be used for re-examination of evaluation of the $^{13}\text{C}(\alpha, n)^{16}\text{O}$ reaction cross-section using multi-channel R-matrix calculations.

Section

Experimental and theoretical studies of nuclear reactions

Primary authors: Dr PRUSACHENKO, P.S. (Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research); Prof. GURBICH, A.F. (Institute for Physics and Power Engineering); Dr BOKHOVKO, M.V. (Institute for Physics and Power Engineering)

Presenter: Dr PRUSACHENKO, P.S. (Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research)

Session Classification: Experimental and theoretical studies of nuclear reactions