

Vector K^* mesons in **strong magnetic field** from **SU(3) lattice gauge theory**

O.V. Teryaev,



E.V. Luschevskaya and E.A. Dorenskaya

KI ITEP

Introduction

- ▶ For the vector K^{0*} and $K^{\pm*}$ mesons we explore on the lattice
 - ▶ The dependence of the energy and magnetic properties from spin projection
 - ▶ The dependence of the magnetic characteristics from $\frac{m_s}{m_d}$ ratio
- ▶ For the K^{0*} and \bar{K}^{*0} mesons we calculate
 - ▶ Magnetic moment which is the new effect
 - ▶ The magnetic dipole polarizability
 - ▶ The tensor polarizability which is the measure of lepton asymmetry
- ▶ For the charged vector $K^{\pm*}$ mesons we find
 - ▶ The magnetic moment

Energy of vector K^{*0} and \bar{K}^{*0} mesons




- Lattice parameters: $N_s^3 \times N_t = 18^4$, $a = 0.105 \text{ fm}$, $m_\pi = 367(8) \text{ MeV}$

The lattice data are fitted by equations:

- $E^2(S_Z = 0) = m^2 - 4\pi m\beta_m(eB)^2 - 4\pi m\beta_m^{h1}(eB)^4 - 4\pi m\beta_m^{h2}(eB)^6$ at $eB \in [0: 1.03] \text{ GeV}^2$
- $E^2(S_Z = \pm 1) = m^2 \mp g(eB) - 4\pi m\beta_m(eB)^2$ at $eB \in [0: 1.23] \text{ GeV}^2$

Fits for magnetic moment and polarizability



The best fit for $\frac{m_s}{m_d} \leq 5.7$:

$$E^2 = m^2 - 4\pi m \beta_m (eB)^2$$

The best fit for $\frac{m_s}{m_d} > 5.7$:

$$E^2 = m^2 - g(eB) - 4\pi m \beta_m (eB)^2$$

Magnetic moment and dipole polarizability of the K^{*0} meson for spin $S_z = -1$.

Previous results:

- ▶ Lattice calculations: $g(K^{*0}) = -0.26$, Hedditch et. al. Phys.Rev.D75 094504 (2007).
- ▶ Light cone QCD sum rules $g(K^{*0}) = 0.26 \pm 0.4$,
Aliev et.al., Phys.Lett B678 470 (2009).
- ▶ Field cumulant method $g(K^{*0}) = -0.183$,
M. Badalian and Yu. A. Simonov, Phys. Rev. D 87 074012 (2013).

Our lattice results:

m_s/m_u	g -factor	$\beta_m(\text{GeV}^{-3})$	p-value	$eB(\text{GeV}^2)$
1	-	-0.026 ± 0.006	0.522	[0 : 0.82]
5.7	-	-0.033 ± 0.003	0.384	[0 : 1.03]
8.6	-0.044 ± 0.106	-0.019 ± 0.005	0.728	[0 : 1.23]
11.4	-0.265 ± 0.100	-0.024 ± 0.005	0.642	[0 : 1.23]
14.3	-0.378 ± 0.097	-0.025 ± 0.004	0.710	[0 : 1.23]
20	-0.599 ± 0.076	-0.027 ± 0.003	0.966	[0 : 1.23]
30	-0.816 ± 0.103	-0.024 ± 0.003	0.913	[0 : 1.23]



Lepton asymmetry and tensor polarizability for K^{*0} and \bar{K}^{*0} mesons

m_s/m_u	$\beta_{S_z=+1}(\text{GeV}^{-3})$	$\beta_{S_z=-1}(\text{GeV}^{-3})$	$\beta_{S_z=0}(\text{GeV}^{-3})$	β_t
1	-0.026 ± 0.006	-0.026 ± 0.006	0.185 ± 0.022	-3.2 ± 0.3
5.7	-0.039 ± 0.006	-0.033 ± 0.003	0.232 ± 0.044	-3.4 ± 0.4
8.6	-0.018 ± 0.012	-0.019 ± 0.005	0.212 ± 0.019	-3.6 ± 0.3
11.4	-0.019 ± 0.010	-0.024 ± 0.005	0.230 ± 0.039	-3.7 ± 0.3
14.3	-0.016 ± 0.009	-0.025 ± 0.004	0.206 ± 0.019	-2.8 ± 0.2
20	-0.011 ± 0.009	-0.027 ± 0.003	0.187 ± 0.017	-2.8 ± 0.3
30	-0.0002 ± 0.007	-0.024 ± 0.003	0.167 ± 0.019	-2.5 ± 0.2

- The large negative values of β_t indicate that the longitudinal polarization dominates for the decays of these mesons.
- The dileptons are mainly emitted in the directions close to the perpendicular ones to the magnetic field axis.

$$\beta_t = \frac{\beta_{S_z=+1} + \beta_{S_z=-1} - 2\beta_{S_z=0}}{\beta_{S_z=+1} + \beta_{S_z=-1} + \beta_{S_z=0}}.$$

Energy of the vector $K^{\pm*}$ mesons



► We find the g-factor from the fit (right figure)

$$E^2 = m^2 + eB - gqS_z(eB)$$

where q is the meson charge, e is the electron charge.

Magnetic moment of $K^{\pm*}$ mesons

Previous results:

- ▶ Lattice background field method from 2pt correlation functions: $|g(K^{\pm*})| = 2.36$,
F.X.Lee et.al., Phys. Rev. D78 094502 (2008).
- ▶ Lattice calculations from 3pt corr. functions: $|g(K^{\pm*})| = 2.23$,
Hedditch et. al. Phys.Rev.D75 094504 (2007).
- ▶ Field cumulant method $|g(K^{\pm*})| = 2.194$,
M. Badalian and Yu. A. Simonov, Phys. Rev. D 87 074012 (2013).


Our lattice results:

for $\frac{m_s}{m_d} = 20$ at $eB \in [0:0.62] \text{ GeV}^2$

$$|g(K^{\pm*})| = 2.42 \pm 0.15.$$

for $\frac{m_s}{m_d} = 30$ at $eB \in [0:0.62] \text{ GeV}^2$

$$|g(K^{\pm*})| = 2.68 \pm 0.16.$$



Conclusions-II: magnetic moment but no tachyonic mode due to strangeness

For K^{*0} mesons

- ▶ The g-factor was found
 - ▶ The magnetic moment of the K^{*0} meson is negative in value, that agrees with other lattice results.
 - ▶ The extrapolations to physical pion mass and continuous limit are necessary.
- ▶ The magnetic dipole polarizability was found
- ▶ The tensor polarizability is negative in value
 - ▶ So the dileptons are mainly emitted in the directions close to the perpendicular ones to the magnetic field axis.

For $K^{\pm*}$ mesons

- ▶ The g-factor was calculated