Contribution ID: 1740

Type: Oral

Direct Photon Interferometry in Bi-Bi Collision at $\sqrt{s_{NN}}$ = 9.2 GeV

Tuesday 29 October 2024 13:00 (15 minutes)

Bose-Einstein interferometry of identical hadrons are known to be one of the key experimental tool to study space-time properties of the hot and dense nuclear matter (fireball) in heavy-ion collisions. However, hadron correlations capture the space-time evolution of the source at the freeze-out stage, when final-state hadrons are formed. On the other hand, direct photon correlation are expected to provide important information on the space-time evolution of the fireball on its earliest stages, as direct photons are emitted from the fireball center. Thus, transverse and longitudinal radii of the fireball and their $k_{\rm T}$ -dependence due to fireball radial extension before freeze-out might be measured using two one-dimensional photon-photon correlation functions. The Multi-Purpose Detector (MPD) of the Nuclotron-based Ion Collider facility (NICA) is designed to study the baryon-rich region of the QCD phase diagram, and Bose-Einstein interferometry is an important component of MPD experimental program. In this talk, the possibility to perform measurements of Bose-Einstein photon-photon correlations in Bi-Bi collision at $\sqrt{s_{NN}} = 9.2$ GeV at MPD using UrQMD model are reported.

Primary author: KUSKOV, Vladislav (National Research Centre Kurchatov Institute)
Presenter: KUSKOV, Vladislav (National Research Centre Kurchatov Institute)
Session Classification: High Energy Physics

Track Classification: High Energy Physics