

Calculations of the cell survival rate after irradiating with minibeam arrays of protons and C-12

Thursday 31 October 2024 15:50 (15 minutes)

Beams of protons and C-12 nuclei are very effective in treating deep-seated solid tumours, but collateral damage to surrounding healthy tissues is still unavoidable. It has been proposed [1], to reduce this damage by splitting a homogeneous radiation field into arrays of sub-millimetre-wide beams called minibeam arrays. Such minibeam arrays become wider with depth of penetration due to multiple Coulomb scattering and production of secondary particles in tissues and finally overlap in the target tumour volume [2].

In this work, the propagation of minibeam arrays of protons and C-12 in a water phantom was simulated using the Geant4 toolkit [3] of version 10.3. Electromagnetic processes were modelled with the G4EMStandard_opt3 physics list, the Binary Cascade (BIC) model was used for proton- and neutron-induced nuclear reactions, and the Quantum Molecular Dynamics (QMD) model was used for nucleus-nucleus collisions. The modified microdosimetric kinetic model (MKM) was used to calculate the relative biological efficiency (RBE), biological dose and cell survival rate of human salivary gland (HSG) cells [4] representing tissues with normal radiosensitivity.

The hexagonal and rectangular minibeam arrays with individual beams of 0.3 mm and 0.5 mm FWHM with the centre-to-centre distance of 2 mm were considered and compared with a homogeneous radiation field. In all cases the beam energies were adjusted to obtain 60 mm wide spread-out Bragg peaks (SOBP) centered at 130 mm. A higher average cell survival rate was calculated at the entrance to the phantom and at the dose plateau for the arrays of minibeam arrays of protons and C-12 in comparison to the homogeneous irradiation field. While a limited number of cells close to the minibeam central axes receive very high peak dose and thus inactivated, many cells between the minibeam arrays have a high survival probability. At the same time, the cell survival rate in the target volume with minibeam arrays was found to be similar to the survival rate in the homogeneous field. Peak-to-valley dose ratios (PVDR) and dose-volume histograms were also calculated as functions of the depth in phantom and compared for all considered minibeam irradiation geometries and homogeneous field. The calculations demonstrated the advantages of minibeam arrays with respect to homogeneous radiation fields.

This study was funded by the Russian Science Foundation (RSF) grant No. 23-25-00285 "Modeling of the phys-

References

1. C. Lamirault et al., "Spatially modulated proton minibeam arrays results in the same increase of lifespan as",
2. I. Pshenichnov et al., "Proton and carbon-ion minibeam therapy: from modelling to treatment", Phys. Part. Nucl. 53, 103 (2022).
3. J. Allison et al., "Recent developments in Geant4," Nucl. Instrum. Methods Phys. Res., Sect. A 835, 186 (2017).
4. Y. Kase et al., Microdosimetric measurements and estimation of human cell survival for heavy-ion beams

Primary authors: SVETLICHNYI, Aleksandr (INR RAS); Mr SAVENKOV, Savva (INR RAS); PSHENICHNOV, Igor (Institute for Nuclear Research, Russian Academy of Sciences)

Presenter: SVETLICHNYI, Aleksandr (INR RAS)

Session Classification: Life Science

Track Classification: Life Science