

Kinetics of DNA double-strand breaks formation in mature neurons of primary rat hippocampal cell culture under the action of ionizing radiation with different physical characteristics

Krupnova M. E.¹, Boreyko A. V.^{1,2}, Hramco T. S.^{1,2}, Pahomova N.V.¹, Shamina D.D.^{1,2}, Yasinskaya A.V.^{1,2}

¹Joint Institute for Nuclear Research, Laboratory of Radiation Biology, Dubna, Russia


²Dubna State University, Dubna, Russia

Introduction

The effects of radiation on the central nervous system are of major interest because of several issues:

- the manned deep-space missions

Cosmic radiation




- the widespread use of radiation in medical treatment


Radiation therapy



DNA damage




Cognitive impairment





Double-strand break is one of the most critical types of DNA damage

Hippocampal functions

Rodent
hippocampus



Human
hippocampus



Rodent
hippocampus



Human
hippocampus

Functions:

- Spatial orientation
- Memory
- Learning

Ionizing radiation exposure:

- Inhibits the adult hippocampal neurogenesis
- Reduces the dendritic arborization
- Impairs nerve impulse conduction
- Changes proteins expression and forms mutant proteins
- Causes cognitive impairment
- Violates animal spatial memory

The hippocampus is considered as one of the most radiosensitive brain structures due to the adult neurogenesis processes occurring in it.

Research aim




Investigation of the influence of radiation with different physical characteristics on DNA DSBs formation in mature neurons of rat primary hippocampal cell culture


Materials and methods

Primary rat hippocampal cell culture

DIV-days in vitro



Brain tissue isolation of *Sprague Dawley* rat pups (P0-P1)



Plating of cell suspension



Cell cultivation



Irradiation
DIV10




Materials and methods

- Irradiation
- Fixation 4% PFA
- Immunocytochemical staining for DNA DSBs markers and cell type markers
- Fluorescent microscopy
- Quantitative analysis

Radiation source	E	LET, keV/ μm	Angle, $^{\circ}$	Dose, Gy	Facility
^{60}Co γ -rays	1.25 MeV	0.3	90	3	«Rocus-M» MTC LNR JINR
Protons	150 MeV	-	10	3	Synchrocyclotron MTC LNR JINR
^{15}N	14 MeV/nucleon	175	10	1.25	U400-M, LNR JINR

RIF (γH2AX , $\gamma\text{H2AX}/53\text{BP1}$) – radiation induced foci



γH2AX - DNA DSBs

MAP2 – soma, axon and dendrites


DAPI - cell nuclei

Merge

The kinetics of RIF formation in mature neurons of primary rat hippocampal cell culture

Measured parameters:


- The number and complexity of colocalized γ H2AX/53BP1 foci in neuronal cells after γ -irradiation
- The number and complexity of γ H2AX foci in mature neurons (MAP2+) after proton irradiation
- The number and complexity of γ H2AX foci in mature neurons (MAP2+) after ^{15}N ions irradiation
- The number and foci complexity shaped by nitrogen ions tracks




The kinetics of γ H2AX and 53BP1 foci formation in mature neurons of primary rat hippocampal cell culture after exposure to radiation with different physical characteristics (x1000 magnification).

The kinetics of RIF formation in mature neurons of primary rat hippocampal cell culture


3 Gy γ -rays of ^{60}Co



3 Gy protons



1.25 Gy ^{15}N -ions




Time after irradiation, h


The statistically significant difference between groups was shown with the Mann-Whitney test (* $P=0,001$)

The kinetics of RIF formation in mature neurons of primary rat hippocampal cell culture


3 Gy γ -rays of ^{60}Co



3 Gy protons




1.25 Gy ^{15}N -ions




The statistically significant difference between groups was shown with the Mann-Whitney test (* $P=0,001$)

The kinetics of RIF formation in mature neurons of primary rat hippocampal cell culture




The statistically significant difference between groups was shown with the Mann-Whitney test ($*P=0,001$)

The kinetics of RIF formation in mature neurons of primary rat hippocampal cell culture




The statistically significant difference between groups was shown with the Mann-Whitney test (* $P=0,001$)

The kinetics of RIF formation in mature neurons of primary rat hippocampal cell culture




The statistically significant difference between groups was shown with the Mann-Whitney test ($*P=0,001$)

The structure of complex RIF clusters in mature neurons of primary rat hippocampal cell culture



The X-axis is a number of individual foci in a complex cluster, the Y-axis is a number of foci in clusters with defined complexity calculated relatively to the total number of analyzed RIF clusters in three dimensional space of nuclei.

The structure of complex RIF clusters in mature neurons of primary rat hippocampal cell culture



The X-axis is a number of individual foci in a complex cluster, the Y-axis is a number of foci in clusters with defined complexity calculated relatively to the total number of analyzed RIF clusters in three dimensional space of nuclei.

Conclusions

- The number of RIF is significantly higher after proton-irradiation in comparison with γ -irradiation that could be connected with the features of the proton energy transmission
- The peak of RIF formation shifts as LET increases to the later post-irradiation time
- The number of RIF is 2.4 times higher 1h after nitrogen irradiation compared to gamma irradiation
- 24h h after nitrogen irradiation the RIF number remained preserved in contrast to proton irradiation: 21 γ H2AX foci - ^{15}N -ions, 13 γ H2AX foci – protons
- The foci clusters complexity increases as LET grows that is clearly demonstrated by the dominance of foci clusters with the higher number of individual foci

Thank you for your attention!
