

Contribution of nontrivial angular momenta in bound state energy of triton in Bethe-Salpeter-Faddeev approach.

Three nucleon system(triton) was considered. Relativistic properties of this system were investigated. For this relativistic generalization of Faddeev approach was used. As two particle T-matrix which contained in relativistic Faddeev equation we used solution of Bethe-Salpeter equation. So eventually we had Bethe-Salpeter-Faddeev(BSF) equation for describe three nucleon system. As potential of nucleon-nucleon(NN) interaction we used separable potential. Form factors of this potential taken in Yamaguchi-type function. Using of separable potential in particular allow to reduce integration on two variables into integration on one variable in BSF equation. Six states $1S_0, 3S_1, 3D_1, 3P_0, 3P_1$ and $1P_1$ with different angular momenta were considered. For this we made particle wave decomposition of BSF equation. System of 12 integral equations(for real and imaginary parts of amplitudes of $1S_0, 3S_1, 3D_1, 3P_0, 3P_1$ and $1P_1$ states) was solved with used iteration method. Bound state energy of triton and amplitudes of S,P and D states was found. Amplitudes used for calculation electric and magnetic form factors of triton.

Primary author: Mr YUREV, Sergey (JOINT INSTITUTE FOR NUCLEAR RESEARCH)

Co-author: Dr BONDARENKO, Serge (JINR)

Presenter: Mr YUREV, Sergey (JOINT INSTITUTE FOR NUCLEAR RESEARCH)

Track Classification: Theoretical Physics