



Status of the BM@N experiment



M.Kapishin





5 Countries, 13 Institutions, 217 participants

- *University of Plovdiv, Bulgaria*
- *St.Petersburg University*
- *Shanghai Institute of Nuclear and Applied Physics, CFS, China;*
- *Joint Institute for Nuclear Research;*
- *Institute of Nuclear Research RAS, Moscow*
- *NRC Kurchatov Institute, Moscow combined with Institute of Theoretical & Experimental Physics. NRC KI. Moscow*
- *Moscow Engineer and Physics Institute*
- *Skobeltsyn Institute of Nuclear Physics, MSU, Russia*
- *Moscow Institute of Physics and Technics*
- *Lebedev Physics Institute of RAS, Moscow*
- *Institute of Physics and Technology, Almaty*
- *Physical-Technical Institute Uzbekistan Academy of Sciences, Tashkent*
- *High School of Economics, National Research University. Moscow*







BM@N Experiment



Production of p , d , t in 3.2 AGeV argon-nucleus interactions



Two classes of centrality <40% and >40% based on barrel detector and track multiplicities



Deuterons: dN/dy dependence on y



$$y^* = y_{\text{lab}} - y_{\text{CM}}, \quad y_{\text{CM}} \approx \langle y(\pi) \rangle$$



Ar+C: $\langle y(\pi) \rangle = 1.27$

Ar+Pb: $\langle y(\pi) \rangle = 0.82$

- dN/dy spectrum softer in interactions with heavier target
- DCM-SMM and PHQMD models describe data shape, but are lower in normalization by factor 4

See talks of L.Kovachev and V.Kolesnikov

Deuterons: $\langle m_t \rangle$ dependence on y



- Maximum $\langle m_t \rangle$ at mid-rapidity y^*
- PHQMD model is in better agreement with data at mid-rapidity than DCM-SMM

BM@N preliminary results, papers, conferences



Production of p , d , t in 3.2 AGeV argon-nucleus interactions at the Nuclotron

BM@N preliminary, extension of the paper draft

The BM@N spectrometer at the NICA-Nuclotron facility

The BM@N detector paper for the Xe+CsI run configuration, submitted to NIM A, arxiv:2312.17573

BM@N presented / submitted physics and detector talks at conferences:

Conference Nucleus-2023, Sarov, October 2023

Workshop NICA-2023, December 2023

Scientific session of the Russian Academy of Sciences, Dubna, April 2024

Conference Nucleus-2024, Dubna, July 2024


Plans for BM@N meetings:

2 Collaboration meetings in a year: April - May, October

Analysis and detector meeting: February - March



Configuration of BM@N detector in Xe+Csl run



Detector paper
arxiv:2312.17573





- Magnet SP-41 (J)
- Vacuum Beam Pipe (1)
- ▨ BC1, VC, BC2 (2-4)
- ▨ SiBT, SiProf (5, 6)
- ▨ Triggers: BD + SiMD (7)
- ▨ FSD, GEM (8, 9)
- ▨ CSC 1x1 m² (10)
- ▨ TOF 400 (11)
- ▨ DCH (12)
- ▨ TOF 700 (13)
- ▨ ScWall (14)
- ▨ FD (15)
- ▨ Small GEM (16)
- ▨ CSC 2x1.5 m² (17)
- ▨ Beam Profilometer (18)
- ▨ FQH (19)
- ▨ FHCAL (20)
- ▨ HGN (21)

Xe¹²⁴ + Csl interactions:

main trigger cover centrality < 70-75% (85% events)

min bias trigger (7% events), beam trigger (3% events)

FSD hit reconstruction in Xe run: 4 Si stations




→ Readout cards with defected chips in stations 2, 3 and 4 are replaced


FSD: after replacing

See talk of D.Chemezov


Cosmic tests




Cosmic tests station 3



Cosmic tests station 2




Cosmic tests station 4




Efficiency of Si and GEM detectors in Si run

A.Zinchenko


Si-2 station: X/Y map of efficiency




GEM-3: X/Y map of efficiency






Si-4 station: X/Y map of efficiency





GEM-5: X/Y map of efficiency



Xe+ CsI data : $\Lambda \rightarrow p\pi^-$, $K_s^0 \rightarrow \pi^+\pi^-$, $\Xi^- \rightarrow \Lambda\pi^-$



In 500M events expect: **4M** Λ , **1.2M** K_s^0 and **8K** Ξ^-




See talk of A.Zinchenko

Life time is in agreement with PDG values: **0.2632 ns** for Λ , **0.0895 ns** for K_s^0





Xe+CsI data: π^\pm , K^\pm , p, He3, d, t identification

Total β vs rigidity

ToF-700, S.Metz





BM@N acceptance for Λ , K_s^0 , identified p, d




Centrality selection from fits of the track multiplicity

MEPhI group

- Parametrization of data track multiplicity N_{ch} by MC Glauber model or Negative Binominal Distribution (Γ -fit) with free parameters
- Extract $P(b | N_{ch})$
- Still need to correct for trigger efficiency, changes in central tracker (FST, GEM) efficiency





Γ -fit and MC-Glauber fit are in agreement





Centrality selection in forward detectors: hodoscope and FHCaI

INR RAS group

Color bins – 10% of number of events in each bin



~30% events with Z^2 signal pile-up in the beam hodoscope



Need to subtract pile-up to determine centrality unbiased

Pile-up correction in fragment hodoscope by signal unfolding

→ remain 4% events with unresolved peaks in multi-ion pile-up

Current tasks for the Xe data analysis



Activities since the last Collaboration meeting in November 2023:

- 2 times processing of event reconstruction using DIRAC at Tier MLIT
- Reasonable signals of Λ and K^0_S , life time within 1 sigma from PDG (A.Zinchenko, V.Vasendina, J.Drnoyan, R.Barak)
- Good agreement between data and reconstructed Λ and K^0_S simulation
- Progress in identification of charged particles in ToF-400 and ToF-700 (M.Rumyantsev, I.Zhavoronkova, S.Merts, N.Huhaeva, V.Plotnikov)
- newly processed data could be used for physics analyses of charged mesons and light nuclear fragments
- Analysis of v_1 and v_2 flows for protons (MEPhI)
- Beam pile-up corrections in fragment hodoscope are done, they are needed for the centrality measurement in fragment hodoscope and hadron calorimeter (INR RAS)



Tasks to be completed for physics analyses:

- Centrality measurement with forward detectors (INR RAS) and track multiplicity (MEPhI), need to compare the results of two methods for Λ and K^0_S
- Evaluate trigger efficiency for different centrality classes



• Topics of physics analyses:

- analysis of production of Λ , Ξ^- hyperons, K^0_S , K^\pm , π^\pm mesons, light nuclear fragments and neutrons in Xe+Csl interactions;
- analysis of collective flow of protons, π^\pm , light nuclear fragments
- search for light hyper-nuclei ${}_\Lambda H^3$, ${}_\Lambda H^4$

Repair and tests of GEM detectors



**S.Piyadin
E.Kulish
S.Khabarov
A.Makankin
and support team**





2-coordinate Si-plane based on STS modules

See talk of D.Dementiev

A new Si-plane based on STS modules to be installed between the **Target** and **Forward Si-Tracker**


Motivation: to improve track and momentum resolution for the low-momentum particles





Plan to install and commission the new Si plane in fall 2024

High Granularity Neutron detector

INR RAS, JINR, NRC Kurchatov → plan to construct in 2024-25




Structure of Scint. layer:
array of 11x11 scintillator cells 4 x 4 cm²



2 positions of HGN detector at BM@N: at 10° and 17°

HGN detector parameters: 2 sub-detectors with 8 layers each ($\sim 1.5 \lambda_{int}$)

- 11 x 11 cells in one layer with SiPM read-out
- first layer works as VETO
- next 7 layers: 3cm Cu + 2.5cm scintillator
- FPGA based fast TDC read-out with additional ToT amplitude measurement
- time resolution of one scint. cell ~ 120 ps
- neutron detection efficiency: $> 60\%$ @ 1GeV



Physics run with the Xe beam in 2024-2025



- beam energy scan in the range of 2-3 AGeV
- same central tracker configuration based on silicon FSD and GEM detectors,
- additional 1st vertex plane of silicon STS detectors
- complete replacement of outer drift chambers with cathode strip chambers
- additional ToF-400 modules to extend acceptance by factor 1.5

Preparations for the physics run with the Bi beam

- further development of the central tracker is foreseen: installation of additional stations of silicon FSD detectors
- It is planned to put into operation a 2-coordinate (X/Y) neutron detector of high granularity to measure neutron yields and collective flow

**Thank you
for attention!**

Directed and elliptic flow at BM@N



- Good agreement between reconstructed and model data
- Approximately 250-300M events are required to perform multi-differential measurements of v_n