




Charged fragments-spectators measurements with Scintillation Wall in the BM@N experiment Xe+Csl@3.8 AGeV

Volkov Vadim


INR RAS

16/05/2024



Scintillation Wall for fragments charge measurements and reaction plane estimation



- 36 small inner cells $7.5 \times 7.5 \times 1 \text{ cm}^3$ + 138 big outer cells $15 \times 15 \times 1 \text{ cm}^3$
- light yield for MIP signal – small cells $55 \text{ p.e.} \pm 2.4\%$; big cells $32 \text{ p.e.} \pm 6\%$.
- optional beam hole (covered with 4 small cells for the SRC run)
- covered with a light-shielding aluminum plate
- light collection by WLS fibers
- light readout with SiPM mounted on the PCB at each scint. cell



light collection from tiles



- Hamamatsu MPPC S14160-1310PS
- $1.3 \times 1.3 \text{ mm}^2$
- Number of pixels: 2668
- Gain: 7×10^5
- PDE: 25%





ScWall: design

41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58							
59	60	A	61	62	63	64	B	66	67	68	69	C	71	72	73	74	D	75	76					
77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94							
95	96	97	98	99	100	E	101	1	2	3	4	5	6	7	8	9	10	102	F	103	104	105	106	107
108	109	110	111	112	113	G	114	11	12	13	14	15	16	17	18	19	20	115	H	116	117	118	119	120
121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138							
139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156							
157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174							

- readout divided into 12 sectors each one equipped with single temperature sensor
- each 4 sectors are read by combined electronics unit:
 - One ADC64s2 board
 - Four 16-channels FEE boards
 - Voltage control unit

ScWall average Z^2 distribution with CsI (2%) target, Xe, CCT2 after calibration


3.8 GeV



ScWall stability during the run8

Data for XeCsI are presented for all data on a file-by-file basis for energies of 3.0 and 3.8 GeV for ScWall

+− 3 σ dashed lines are shown





The mean total charge values for ScWall for each file are presented.

Applied cuts: 1 Xe, vertex Z ($-1.5 < Z < 1.5$)

Data taken on March 24

Charges spectra




The charge spectrum on the ScWall is in the range up to $Z = 2$ (small cells).

Large charges leak out into the hole.

In the cells around the hole, charges up to $Z = 4$ can be detected.

The shift of the peaks is due to the Birks effect.




The charge yields in the experiment and in the simulation data for $Z = 2$ are significantly different.




Particle yield difference in the models and data is related to the angular distribution of the particles.

Future upgrade of ScWall

ScWall Z^2 distributions
XeCs@3.26AGeV
 DCM-QGSM-SMM





ScWall Z^2 distributions
BiBi@3AGeV
 DCM-QGSM-SMM



The **range** of charges detected on the ScWall is much **greater** (up to $Z = 5$) in small cells when **thicker cells (25 mm)** are used according to the DCM-QGSM-SMM simulation.

41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58					
59	60	A	62	63	64	65	B	66	67	68	69	C	70	71	72	73	74	D	75	76		
77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94					
95	96	97	98	99	100	101	1	2	3	4	5	6	7	8	9	10	102	103	104	105	106	107
108	109	110	111	112	113	114	11	12	13	14	15	16			19	20						
							21	22	23	24	25	26			29	30						
							31	32	33	34	35	36	37	38	39	40						
121	122	I	123	124	125	126	J	128	129	130	131	K	132	133	134	135	136	137	138			
139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156					
157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174					



Centrality estimators



As an estimator of centrality, FHCAL E_{dep} performs best (similar to the number of tracks).



The scintillation wall can sense centrality, but much worse.

It is possible to use the combined observable of these quantities to determine centrality.



DCM—SMM 3.8 GeV

Centrality estimators and combination of observables




The combined usage of the energy deposition in the FHCal and the total charge on the ScWall gives a narrower distribution.

The centrality accuracy improves only within 1%.

Need to consider autocorrelations with FHCal.
ScWall can be used to estimate systematics.

Centrality estimators: ScWall vs FHCal




The width of the distributions of the presented observables as a dependence of the impact factor shows that the ScWall is significantly inferior to the FHCal.

The difference for the most central events in standard deviation units is about 2 times.


DCM—SMM 3.8 GeV

Centrality estimators: correlations



The best observables for centrality are tracks multiplicity and energy deposition in the FHCAL.


The ScWall can only be used to slightly improve the results.



The combined usage of tracks multiplicity energy deposition in the FHCAL can significantly improve the results.


[Pearson](#) correlation coefficient is used.

Purity and centrality for FHCaI



To obtain the required purity of 80% for the most central class, it is necessary to take classes size of at least 20%.

Flow measurements theory



The azimuthal angle distribution is decomposed in a Fourier series relative to reaction plane angle:

$$\rho(\varphi - \Psi_{RP}) = \frac{1}{2\pi} \left(1 + 2 \sum_{n=1}^{\infty} v_n \cos n(\varphi - \Psi_{RP}) \right)$$

Anisotropic flow: $v_n = \langle \cos [n(\varphi - \Psi_{RP})] \rangle$

Reaction plane is not experimentally measured, we define the symmetry plane (SP) from spectators:

$$Q_1 = \sum_{k=1}^N w_k (\cos \phi_k, \sin \phi_k) = |Q_1| (\cos \Psi_{SP}, \sin \Psi_{SP})$$

Directed flow is measured

$$v_1 = \frac{\langle \cos(\phi - \Psi_{SP}) \rangle}{R_1}$$

Resolution correction factor

$$R_1 = \langle \cos(\Psi_{SP} - \Psi_{RP}) \rangle$$

Comparison of RP resolution from FHCAL and ScWall

Scalar product (SP) method:



$$v_1 = \frac{\langle u_1 Q_1^{F1} \rangle}{R_1^{F1}} \quad v_2 = \frac{\langle u_2 Q_1^{F1} Q_1^{F3} \rangle}{R_1^{F1} R_1^{F3}}$$

Where R_1 is the resolution correction factor

$$R_1^{F1} = \langle \cos(\Psi_1^{F1} - \Psi_1^{RP}) \rangle$$

Symbol "F2(F1,F3)" means R_1 calculated via (3S resolution):

$$R_1^{F2(F1,F3)} = \frac{\sqrt{\langle Q_1^{F2} Q_1^{F1} \rangle \langle Q_1^{F2} Q_1^{F3} \rangle}}{\sqrt{\langle Q_1^{F1} Q_1^{F3} \rangle}}$$



3 vectors (F1, F2, F3 and S1, S2, S3) each from FHCAL and ScWall were selected and the resolutions were compared.

The ScWall symmetry plane is more fluctuating. Hence SP has lower resolution, and requires more statistics for flow calculations.

Conclusion

- The ScWall performance during run8 was demonstrated
 - ScWall was stable during run8
 - Charge spectra up to $Z = 2$, in central small cells up to $Z = 4$
 - Upgrade (25 mm cells) can greatly improve charge separation
- ScWall centrality and RP are compared with FHCAL, Hodoscope and other various variables
 - ScWall is weakly correlated with centrality
 - And has worse capability for RP determination
 - Can be used for systematics studies
- The ScWall can be used to measure the charged fragment-spectator yields. Such data are important for further constraints on the models.

Thank you for your attention!

backup

ScWall Cell 25 mm, 10 mm


ScWall Z² distributions (RECO, w/o vertex selection)

BiBi@3A GeV, DCM-QGSM-SMM UNIGEN,
 With magnetic field Map_1900 scale 0.834
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh= -1cm, rotY 4.2 deg
 ScWall hole 697.4 cm, Xsh=65.65cm, Ysh=-0.43 cm
 air in cave, Magnet, vac Tubes
 99978ev,


Geometry: air in cave, Magnet, vac. tubes, vac Wall before ScWall, ScWall, Hodo

Not correct nb of entries


Small cells






Large cells






All cells



Cell thickness
10 mm



Cell thickness
25 mm




ScWall ratio Z^2 distributions (RECO, w/o vertex selection)

Geometry: air in cave, Magnet, vac. tubes, vac Wall before ScWall, ScWall, Hodo

BiBi@3A GeV, DCM-QGSM-SMM UNIGEN,
 With magnetic fieldMap_1900 scale 0.834
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh= -1cm, rotY 4.2 deg
 ScWall hole 697.4 cm, Xsh=65.65cm, Ysh=-0.43 cm
 air in cave, Magnet, vac Tubes
 99978ev, vacBox


hZ2ScWall_small

h3	
Entries	645
Mean	17.04
RMS	10.01



hZ2ScWall_small

h3	
Entries	645
Mean	17.04
RMS	10.01




ScWall Z² distributions (BiBi@3.26GeV)


BiBi@3A GeV, DCM-QGSM-SMM UNIGEN,
 With magnetic field Map_1900 scale 0.834
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh= -1cm, rotY 4.2 deg
 ScWall hole 697.4 cm, Xsh=65.65cm, Ysh=-0.43 cm
 air in cave, Magnet, vac Tubes
 99978ev, IDEAL geometry

Ideal geometry: air in cave, Magnet, vac. tubes, vac Wall before ScWall, ScWall, Hodo


Small cells






Large cells



All cells



BiBi@3AGeV, ideal geometry
 All cells thickness **10 mm**
 RECO, no vertZ cut




BiBi@3AGeV, ideal geometry
 All cells thickness **25 mm**
 RECO, no vertZ cut


ScWall Z^2 distributions (XeCs@3.26AGeV)

XeCs@3.26A GeV, DCM-QGSM-SMM UNIGEN,
 With magnetic fieldMap_1900 scale 0.929
 FHCAL, Hodo rotY 1.6 deg, 4.2 deg
 ScWall hole 697.4 cm, Xsh=65.65cm, Ysh=-0.43 cm
 ScWall hole 741.5 cm, Xsh=68.7cm
 58804ev, 199976ev, FULL geometry


Small cells






Large cells



All cells



Full geometry
 ScWall at 741.5 cm, 58804 ev
 Small cells thickness **10 mm**
 RECO, with vertZ cut (+-1.5 cm)



Full geometry
 ScWall at 697.4 cm, 199976ev
 Small cells thickness **25 mm**
 RECO, with vertZ cut (+-1.5 cm)

ScWall Z² distributions




BiBi@3A GeV, DCM-QGSM-SMM UNIGEN,
With magnetic fieldMap_1900 scale 0.834
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh= -1cm, rotY 4.2 deg
 ScWall hole 697.4 cm, Xsh=65.65cm, Ysh=-0.43 cm
 air in cave, Magnet, vac Tubes
99978ev, IDEAL geometry

XeCs@3.26A GeV, DCM-QGSM-SMM UNIGEN,
With magnetic fieldMap_1900 scale 0.929
 FHCAL, Hodo rotY 4.2 deg
 ScWall hole 697.4 cm, Xsh=65.65cm, Ysh=-0.43 cm
199976ev, FULL geometry




Small cells

Large cells




All cells



BiBi@3AGeV, ideal geometry
 All cells thickness 10 mm
 RECO, no vertZ cut




BiBi@3AGeV, ideal geometry
 All cells thickness 25 mm
 RECO, no vertZ cut



XeCs@3.26AGeV, full geometry
 Small cells thickness 25 mm
 RECO, with vertZ cut (+-1.5 cm)

Common Events Percentage Matrix by Class Indices




Event selection






≥ 2 tracks in vertex reconstruction

Single Xe ion selected with Beam Counter BC1S

With cuts on vertex Z ($-1.5\text{cm} < Z < 1.5\text{cm}$)





Charge distribution in ScWall cells (CCT2)







Charge distribution over the scintillation wall. A peaks corresponding to charges $Z = 1, 2$ can be clearly seen.

Charge distribution in ScWall cells



- Comparison of the charge distributions over the scintillation wall for the two energies at 3.0 and 3.8 GeV for the CCT2 trigger.
- The two cell types (small and big) are presented separately.
- It can be seen that the distributions are very similar, with a slight difference in the second peak.

ScWall multiplicity distributions of charged particles for different centrality classes



ScWall multiplicity refers to the number of fired cells in the wall.



Multiplicity is sensitive to centrality -> can be used as estimator. Green, red and blue reflect the most central, semi-central and semi-peripheral arbitrary classes of events.

~50% of minbias events, need to be checked with sim ($b < 10$ fm).

- Cuts:
- BC1S (1 Xe)
 - Z^2 (ScWall) > 0.4
 - vertex Z (-1.5 < Z < 1.5)
 - Z^2 (FQH) < 50




MBT

Multiplicity in ScWall / multiplicity in BD




Multiplicity correlates with energy deposition in the calorimeter, and anticorrelates with multiplicity in BD.

Ambiguity in multiplicity vs number of global tracks





Multiplicity distribution of charged particles in ScWall



Multiplicity is sensitive to charges on the wall for both energies. The peak corresponding to the single charge is clearly prominent.


This dependency can be used for comparison with Monte Carlo models (DCM-QGSM-SMM etc.)

ScWall QA




ScWall $Z^2 > 0.5$

hMultScWall_with_vertZ_cut



ScWall $0.5 < Z^2 < 1.5$

hMultScWall_with_vertZ_cut_Z2_EQ_1



ScWall multiplicities with different number of global tracks in evt


XeCs@3.26A GeV, DCM-QGSM-SMM, UNIGEN
Scale 0.929
 FHCal 977.8 cm, Xsh=65.3 cm, Ysh=-0.8cm, rotY 1.6 deg
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh=-1cm, rotY 1.6 deg
 ScWall hole 741.5 cm, Xsh=68.7cm
 air in cave, Magnet, **all BMN detectors**
 VacZdcWall 200x200cm before nDet 12x12cm 27.3deg
Simul - 58992 ev, RECO - 58804 ev

XeCs@3.8A GeV, PHQMD, UNIGEN
Scale 0.929
 FHCal 977.8 cm, Xsh=65.3 cm, Ysh=-0.8cm, rotY 1.6 deg
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh=-1cm, rotY 1.6 deg
 ScWall hole 741.5 cm, Xsh=68.7cm
 air in cave, Magnet, **all BMN detectors**
 VacZdcWall 200x200cm before nDet 12x12cm 27.3deg
Simul - 281163 ev, RECO - 279140 ev, no etaCut


DCMSMM

Simulation
 (after RECO,
 with reconstructed vertex
 Z cut -1.5cm - 1.5 cm)

hMultScWall_with_vertZ_cut



hMultScWall_with_vertZ_cut_Z2_EQ_1




PHQMD

W/o cut on number of global tracks

	DCMSMM	PHQMD
Number of global tracks > 5	15	15
Number of global tracks > 12	19	19
Number of global tracks > 13	21	21
Number of global tracks > 14	23	23
Number of global tracks > 15	25	25

ScWall small cells 3.8 GeV



Runs 7830 – 7885

360k events

3.8 GeV

FHCal

7839, 7840, 7850, 7856, 7905, 7907, 7950, 7969,
7970, 7972, 7973, 7979, 7997, 8066, 8077, 8111,
8129, 8184, 8186, 8216, 8247, 8289, 8304

Hodo

7839, 7840, 7897, 7901, 7969, 7970, 7972, 7973,
8014, 8063, 8075, 8081, 8088, 8131, 8167, 8175,
8215, 8216, 8247, 8307, 8308

ScWall

7839, 7840, 7900, 7969, 7970, 7972, 7973, 8059,
8167, 8216, 8219, 8307, 8308

3.0 GeV

FHCal

8312, 8323, 8341, 8414, 8419




Hodo

8312, 8321, 8334, 8341, 8395




ScWall

8312, 8421

3.8 GeV



3.0 GeV



Simulation and experiment comparison (ScWall multiplicity)

XeCs@3.26A GeV, DCM-QGSM-SMM, UNIGEN
Scale 0.929
 FHCAL 977.8 cm, Xsh=65.3 cm, Ysh=-0.8cm, rotY 1.6 deg
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh=-1cm, rotY 1.6 deg
 ScWall hole 741.5 cm, Xsh=68.7cm
 air in cave, Magnet, **all BMN detectors**
 VacZdcWall 200x200cm before nDet 12x12cm 27.3deg
Simul - 58992 ev, RECO - 58804 ev

XeCs@3.8A GeV, PHQMD, UNIGEN
Scale 0.929
 FHCAL 977.8 cm, Xsh=65.3 cm, Ysh=-0.8cm, rotY 1.6 deg
 Hodo 970.2 cm, Xsh=64.9 cm, Ysh=-1cm, rotY 1.6 deg
 ScWall hole 741.5 cm, Xsh=68.7cm
 air in cave, Magnet, **all BMN detectors**
 VacZdcWall 200x200cm before nDet 12x12cm 27.3deg
Simul - 281163 ev, RECO - 279140 ev, no etaCut

Simulation
 (after RECO, with reconstructed vertexZ cut
 -1.5cm – 1.5 cm
 no trigger cut, MB
DrawNormalized()

Experiment
 (run 8
 XeCsI@3.8 AGeV,
 MBT trigger
DrawNormalized() Vadim)

ScWall $Z^2 > 0.5$

ScWall $0.5 < Z^2 < 1.5$




ScWall $Z^2 > 0.5$

ScWall $0.5 < Z^2 < 1.5$




All cells

Small cells

Large cells



DCMSMM (b < 10 fm) & experiment





PHQMD (b < 9 fm) & experiment

ScWall $Z^2 > 0.5$

ScWall $0.5 < Z^2 < 1.5$

DCMSMM

ScWall multiplicities
with different impact
parameter cuts



XeCs@3.26A GeV, DCM-QGSM-SMM, UNIGEN

Scale 0.929

FHCal 977.8 cm, Xsh=65.3 cm, Ysh=-0.8cm, rotY 1.6 deg

Hodo 970.2 cm, Xsh=64.9 cm, Ysh=-1cm, rotY 1.6 deg

ScWall hole 741.5 cm, Xsh=68.7cm

air in cave, Magnet, **all BMN detectors**

VacZdcWall 200x200cm before nDet 12x12cm 27.3deg



Simul - 58992 ev, RECO - 58804 ev

Simulation

(after RECO,

with reconstructed vertex

Z cut -1.5cm - 1.5 cm



W/o impact parameter cut

Impact parameter < 11 fm

Impact parameter < 10 fm

Impact parameter < 9 fm

Impact parameter < 8 fm



Impact parameter < 7 fm

ScWall $Z^2 > 0.5$

ScWall $0.5 < Z^2 < 1.5$



PHQMD

ScWall multiplicities
with different impact
parameter cuts



XeCs@3.8A GeV, PHQMD, UNIGEN
Scale 0.929
FHCAL 977.8 cm, Xsh=65.3 cm, Ysh=-0.8cm, rotY 1.6 deg
Hodo 970.2 cm, Xsh=64.9 cm, Ysh=-1cm, rotY 1.6 deg
ScWall hole 741.5 cm, Xsh=68.7cm
air in cave, Magnet, **all BMN detectors**
VacZdcWall 200x200cm before nDet 12x12cm 27.3deg
Simul - 281163 ev, RECO - 279140 ev, no etaCut

Simulation
(after RECO,
with reconstructed vertex
Z cut -1.5cm – 1.5 cm



W/o impact parameter cut
Impact parameter < 11 fm
Impact parameter < 10 fm
Impact parameter < 9 fm
Impact parameter < 8 fm
Impact parameter < 7 fm