

Global polarization and spin alignment measurements

Jinhui Chen

Fudan University

The 2nd China-Russia Joint Workshop on NICA Facility


2024/9/10-13

Outline


- Brief introduction of the global polarization
- Measurements in heavy-ion collisions and prospects
 - Hyperon polarization
 - Vector meson spin alignment
- Summary

Global polarization in HIC

Liang, Wang Phys. Rev. Lett. **94**, 102301(2005); Phys. Lett. B **629**, 20 (2005)





Large OAM L is deposited in the interaction region



- The initial momentum gradient should result in a net angular momentum (shear) in this direction that will be transferred to quark spin via spin-orbit interaction, this effect may not be washed out during interaction and hadronization
- Spin-vorticity coupling Betz, Gyulassy, Torrieri Phys. Rev. C **76**, 044901 (2007); Becattini, Piccinini, Rizzo Phys. Rev. C **77**, 024906 (2008)
- Connection to classical world, the Barnett effect, a fraction of the L associated with the body rotation is transformed into the spin L of the electron

Experimental measurements: Λ

- The global quark polarization along L have many observable consequences in non-central HIC
- Λ are self-analyzing, proton tends to be emitted along the spin direction of the Λ




$$\frac{dN}{d \cos \theta^*} = \frac{1}{2} (1 + \alpha_H |\mathcal{P}_H| \cos \theta^*)$$

$$P_H = \frac{8}{\pi \alpha_H} \frac{\langle \sin(\Psi_1^{\text{obs}} - \phi_p^*) \rangle}{\text{Res}(\Psi_1)}$$

Experimental measurements: Λ (cont.)


STAR Col. Phys. Rev. C **76**, 024915 (2007)



from 2007 to 2017



Lambda hyperons show a positive polarization of the order of a few percent




R. Bellwied for E896 Col. QM2001, Nucl. Phys. A 698, 499 (2002)

“The produced Λ are polarized at freeze-out in the heaviest collision system”

...

“The STAR detector will be capable of repeating these measurements at RHIC energies”




Experimental measurements: Λ (cont.2)

- The fluid vorticity was estimated from the data using hydrodynamics relation


$$\omega \approx k_B T (\bar{P}_{\Lambda'} + \bar{P}_{\bar{\Lambda}'}) / \hbar$$

- The collision energy-average polarization data from STAR BES-I indicate a vorticity of $(9 \pm 1) \times 10^{21} \text{ s}^{-1}$
 \rightarrow experimental access to the vortical structure of the QGP



- Late-stage B field

$$|B| \approx \frac{T_s |P_{\bar{\Lambda}} - P_{\Lambda}|}{2|\mu_{\Lambda}|}$$




- High precision BES-II data of 19.6 and 27 GeV


- $-0.018 \pm 0.127 \text{ (stat.)} \pm 0.024 \text{ (syst.)}$
- $0.109 \pm 0.118 \text{ (stat.)} \pm 0.022 \text{ (syst.)}$

$$B < 9.4 \times 10^{12} \text{ T and } B < 1.4 \times 10^{13} \text{ T}$$

Measurements of Λ and Ξ, Ω



- Measurements in different Exps. -didn't see the "drop" trend?



Deng et al., Phys. Rev. C **101**, 064908 (2020)
Guo et al., Phys. Rev. C **104**, L041902 (2021)...

- Measurements extend to multistrange

STAR Col. Phys. Rev. Lett. **126**, 162301 (2021)



$$\langle P_{\Xi} \rangle = 0.47 \pm 0.10(\text{stat}) \pm 0.23(\text{syst})\%$$

$$\langle P_{\Omega} \rangle = 1.11 \pm 0.87(\text{stat}) \pm 1.97(\text{syst})\%$$


Prospects at NICA-MPD

Heavy ion colliding beams: upto Au @ $\sqrt{s_{NN}} = 4 - 11$ GeV \rightarrow MPD

- The energies are exactly located at the place to understand the transitions?



Figs. from Zebo




Andronic et al., Phys. Letts. B **697** (2011) 203

- The detectors are powerful to identify hyperons with good signal to background ratio, and maybe the light hypernuclei polarization feasible?

Experimental measurements: φ, K^*

- Vector meson ($J=1^-$) spin alignment




- ✓ Spin tensor polarization
- ✓ Different probabilities among three spin states
- ✓ Only ρ_{00} is measurable

$$|11\rangle = |\uparrow\uparrow\rangle$$


$$|10\rangle = \frac{1}{\sqrt{2}}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)$$

$$|1-1\rangle = |\downarrow\downarrow\rangle$$




$$\rho^V = \begin{pmatrix} \rho_{11} & \rho_{10} & \rho_{1-1} \\ \rho_{01} & \rho_{00} & \rho_{0-1} \\ \rho_{-11} & \rho_{-10} & \rho_{-1-1} \end{pmatrix}$$

$$\frac{dN}{d(\cos\theta^*)} \propto (1 - \rho_{00}) + (3\rho_{00} - 1)\cos^2\theta^*$$



$$\rho_{00} < 1/3$$




$$\rho_{00} > 1/3$$


Experimental measurements: ϕ, K^* (cont.)

STAR Col. Phys. Rev. C **77**, 061902 (2008)


ALICE Col. Phys. Rev. Lett. **125**, 012301 (2020)



- Early data of Au+Au 200 GeV suffer from large uncertainties
- Updated measurements seem to provide evidence of spin-orbital angular momentum interactions, but production plane and random plane also show deviation at small p_T



New Measurements ϕ, K^*0 @ non-central collisions



- New measurements extend the study to lower energies with high statistics, @200 GeV, a factor of ~ 50 more event statistics analyzed.



- We see that the signal for the ϕ meson occurs mainly within ~ 1.0 - 2.4 GeV/c; at larger p_T the results can be regarded as being consistent with $1/3$ within $\sim 2\sigma$ or less.

* 1st order EP: ZDC or BBC

* 2nd order EP: TPC

STAR Col. Nature **614**, 244 (2023)

New Measurements φ, K^{*0} @non-central collisions




- K^{*0} is a combination of K^{*0} and anti- K^{*0}
- Independent analysis
- Different from the φ meson data, the K^{*0} data is largely consistent with $1/3$, within statistics and systematical uncertainties


STAR Col. Nature **614**, 244 (2023)

Study the fine structure vs. centrality

STAR Col. Nature **614**, 244 (2023)




Gao, Chen, Deng, Liang, Wang, Wang
Phys. Rev. C **76**, 044901 (2007)



At high energies (≥ 62.4 GeV) for ϕ , and (≥ 39 GeV) for K^{*0} , ρ_{00} in central collisions tends to $\leq 1/3$. This might be caused by transverse local spin alignment and a contribution from the helicity polarization of quarks.

Results mid-central & averaged over p_T



- 1) ϕ -meson is significantly above 1/3 for $\sqrt{s} \leq 62$ GeV
- 2) K^* is largely consistent with 1/3
- 3) Averaged over 62 GeV and below:
 - 0.3541 ± 0.0017 (stat.) ± 0.0018 (sys.) for ϕ
 - 0.3356 ± 0.0034 (stat.) ± 0.0043 (sys.) for K^*

* Different approaches are used in the combinatorial bg. analysis

STAR Col. Nature **614**, 244 (2023)

Expectations of ρ_{00} from theory



Physics Mechanisms	(ρ_{00})
c_Λ : Quark coalescence vorticity & magnetic field ^[1]	< 1/3 (Negative $\sim 10^{-5}$)
c_ε : Vorticity tensor ^[1]	< 1/3 (Negative $\sim 10^{-4}$)
c_E : Electric field ^[2]	> 1/3 (Positive $\sim 10^{-5}$)
Fragmentation ^[3]	> or, < 1/3 ($\sim 10^{-5}$)
Local spin alignment and helicity ^[4]	< 1/3
Turbulent color field ^[5]	< 1/3
c_ϕ : Vector meson strong force field ^[6]	> 1/3

$$\rho_{00}^\phi \approx \frac{1}{3} + c_\omega + c_\varepsilon + c_{EM} + c_\phi + c_{LV} + c_h + c_{TC} + c_{\text{shear}}$$

- [1]. Yang et al., Phys. Rev. C **97**, 034917 (2018) [2]. Sheng et al., Phys. Rev. D **101**, 096005 (2020)
 [3]. Xia et al., Phys. Lett. B **817**, 136325 (2021) [4]. Gao, Phys. Rev. D **104**, 076016 (2021)
 [5]. Muller, Yang, Phys. Rev. D **105**, L011901 (2022) [6]. Li, Liu, arXiv:2206.11890,
 Wagner, Weickgenannt, Speranza, arXiv:2207.01111

The local correlation or fluctuation of ϕ fields is the dominant mechanism for the observed ϕ -meson ρ_{00}

Sheng, et al., Phys. Rev. Lett. **131**, 042304 (2023)



The small Λ vs. large ϕ -meson signal

Lv, Yu, Liang, Wang, Wang, PRD **109**, 114003 (2024)





$$\rho_{00}^V = \frac{1}{C_V} \left\{ 1 + c_{xx}^{(q_1 \bar{q}_2)} + c_{yy}^{(q_1 \bar{q}_2)} - c_{zz}^{(q_1 \bar{q}_2)} + P_{q_1 x} P_{\bar{q}_2 x} + P_{q_1 y} P_{\bar{q}_2 y} - P_{q_1 z} P_{\bar{q}_2 z} \right\}$$

$$P_\Lambda(\alpha_\Lambda) = \bar{P}_{sz} - \frac{\bar{c}_{iz}^{(uds)} + \bar{c}_{iz}^{(us)} \bar{P}_{di} + \bar{c}_{iz}^{(ds)} \bar{P}_{ui}}{1 - \bar{c}_{ii}^{(ud)} - \bar{P}_{ui} \bar{P}_{di}}$$

$$\langle P_\Lambda \rangle \sim \langle P_s \rangle,$$



$$\langle \rho_{00}^\phi \rangle \sim \frac{1 - \bar{c}_{zz;\phi}^{(s\bar{s})} - \langle P_s \rangle^2}{3 + \bar{c}_{zz;\phi}^{(s\bar{s})} + \langle P_s \rangle^2},$$

$$\langle c_{zz}^{\Lambda\bar{\Lambda}} \rangle \sim \bar{c}_{zz;\Lambda\bar{\Lambda}}^{(s\bar{s})} + \langle P_s \rangle^2,$$



X.N. Wang, Chirality 2024 7/22-26, Romania

“Strong-field induced hyperon spin correlation”






*quark-antiquark spin correlations in ee/pp (helicity frame)

Chen, Goldstein, Jaffe, Ji, NPB **445**, 380 (1995); Chen, Yang, Zhou, Liang, PRD **95**, 034009 (2017); Zhang, Wei, PLB **839**, 137821 (2023)

From φ to other mesons

ALICE Col. Phys. Rev. Lett. **131**, 042303 (2023)

ALICE Col. QM2023





ALI-PREL-547532

- Forward rapidity J/ψ
 $\rho_{00} < 1/3$ at LHC
- Midrapidity J/ψ $\rho_{00} \sim 1/3$ at RHIC
- D^{*+} shows a clear p_T dependence

→ The underlying physics seems not converged?





Study the rapidity dependence



RHIC & LHC data : strong rapidity dependence

Prospects at NICA-MPD for mesons

- The different species of vector meson spin alignment from RHIC/LHC seems not converged, independent measurements will be very helpful to understand the underlying physics
- NICA-MPD can identify the vector mesons well, thus will be excellent experiment to measure the light flavor spin alignment



Figs. from Zebo

Summary

- Spin polarization opens a new avenue to investigate heavy-ion collisions
- Global hyperon polarization is observed with the order of a few percent. It represents a measure of the average value of the global quark polarization in the system
- Global vector meson spin alignment is observed with a surprisingly large pattern for ϕ -meson. It represents a local fluctuation/correlation between quark and anti-quark polarization
- Measurements as a function of collision energies, different hadron species are on-going, rich physics to be explored, and the NICA-MPD will be very powerful to establish the feature of high-baryon density region