


# Highlights and Prospects of the COMPASS, AMBER and LHCspin Experiments at CERN: Spin and TMD Structure of Hadrons

Bakur Parsamyan




March 12, 2025  
JINR Dubna, Russia




# Nucleon spin structure: TMD

- 1964 Quark model







- 1969 Parton model



- 1973 asymptotic freedom and QCD




- 1976 large transverse single spin asymmetry in forward  $\pi^\pm$  production



# Nucleon spin structure: TMD


- 1964 Quark model




- 1969 Parton model



- 1973 asymptotic freedom and QCD




- 1976 large transverse single spin asymmetry in forward  $\pi^\pm$  production




- TMD Sivers and Collins or

- Twist-3 Sivers (Efremov-Teryaev-Qui-Sterman (ETQS) function)




# Nucleon spin structure: TMD


- 1964 Quark model



- 1969 Parton model





- 1973 asymptotic freedom and QCD



- 1976 large transverse single spin asymmetry in forward  $\pi^\pm$  production


- 1978 intrinsic transverse motion of quarks and azimuthal asymmetries




(SLAC) Phys. Rev. Lett. 31, 786 (1973)  
(EMC) Phys. Lett. B 130 (1983) 118,  
(EMC) Z. Phys. C34 (1987) 277  
(EMC) Z. Phys. C52, 361 (1991).  
(E665) Phys. Rev. D48 (1993) 5057  
(ZEUS) Eur. Phys. J. C11, 251 (1999)  
(ZEUS) Phys. Lett. B 481, 199 (2000)  
(H1) Phys. Lett. B654, 148 (2007)

# Nucleon spin structure: TMD


- 1964 Quark model



- 1969 Parton model





- 1973 asymptotic freedom and QCD



- 1976 large transverse single spin asymmetry in forward  $\pi^\pm$  production

- 1978 intrinsic transverse motion of quarks and azimuthal asymmetries





# Nucleon spin structure (twist-2): collinear approach $\leftrightarrow$ TMDs

quark			
	<b>U</b>	<b>L</b>	<b>T</b>
<b>U</b>	$f_1^q(x)$ number density		
<b>L</b>		$g_1^q(x)$ helicity	
<b>T</b>			$h_1^q(x)$ transversity



quark			
	<b>U</b>	<b>L</b>	<b>T</b>
<b>U</b>	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders <b>T-odd</b>
<b>L</b>		$g_1^q(x, \mathbf{k}_T^2)$ Helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
<b>T</b>	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers <b>T-odd</b>	$g_{1T}^q(x, \mathbf{k}_T^2)$ Kotzinian-Mulders worm-gear T	$h_1^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity

- PDFs – universal (process independent) objects; T-odd PDFs – conditionally universal



# Nucleon spin structure (twist-2): TMDs


quark

	<b>U</b>	<b>L</b>	<b>T</b>
<b>U</b>	number density		Boer-Mulders
<b>L</b>		helicity	worm-gear L
<b>T</b>	Sivers	Kotzinian-Mulders worm-gear T	transversity pretzelosity

- spin of the nucleon; 
 - spin of the quark 
 -  $\mathbf{k}_T$

quark

	<b>U</b>	<b>L</b>	<b>T</b>
<b>U</b>	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders <b>T-odd</b>
<b>L</b>		$g_1^q(x, \mathbf{k}_T^2)$ Helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
<b>T</b>	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers	$g_{1T}^q(x, \mathbf{k}_T^2)$ Kotzinian-Mulders worm-gear T	$h_1^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity



# SIDIS x-section and TMDs at twist-2

$$\frac{d\sigma^{LO}}{dx dy dz dp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L})$$

$$\times \left\{ \begin{array}{l} 1 + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + S_L \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h + S_L \lambda \sqrt{1-\varepsilon^2} A_{LL} \\ \\ + S_T \left[ \begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_S)} \sin(\phi_h - \phi_S) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_S)} \sin(3\phi_h - \phi_S) \end{array} \right] \\ \\ + S_T \lambda \left[ \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_S)} \cos(\phi_h - \phi_S) \right] \end{array} \right\}$$


	quark		
	<b>U</b>	<b>L</b>	<b>T</b>
<b>U</b>	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders <b>T-odd</b>
<b>L</b>		$g_1^q(x, \mathbf{k}_T^2)$ Helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
<b>T</b>	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers <b>T-odd</b>	$g_{1T}^q(x, \mathbf{k}_T^2)$ Kotzinian-Mulders worm-gear T	$h_{1T}^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity

$A_{UU}^{\cos 2\phi_h} \propto h_1^{\perp q} \otimes H_{1q}^{\perp h}$	Boer-Mulders (T-odd)
$A_{UT}^{\sin(\phi_h - \phi_S)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$	Sivers (T-odd)
$A_{UT}^{\sin(\phi_h + \phi_S)} \propto h_1^q \otimes H_{1q}^{\perp h}$	Transversity
$A_{UT}^{\sin(3\phi_h - \phi_S)} \propto h_{1T}^{\perp q} \otimes H_{1q}^{\perp h}$	Pretzelosity

$$\otimes \equiv \mathbb{C}[wfD] = x \sum_q e_q^2 \int d^2 \mathbf{k}_T d^2 \mathbf{p}_\perp \delta^{(2)}(z \mathbf{k}_T + \mathbf{p}_\perp - \mathbf{P}_h) w(\mathbf{k}_T, \mathbf{p}_\perp) f^q(x, \mathbf{k}_T^2) D_q^h(x, p_\perp^2)$$

12 March 2025

B. Parsamyan



# Single-polarized Drell-Yan x-section and twist-2 TMDs

$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS})$$


$$\times \left\{ \begin{array}{l} 1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\varphi_{CS}} \cos 2\varphi_{CS} \\ + S_L \sin^2 \theta_{CS} A_L^{\sin 2\varphi_{CS}} \sin 2\varphi_{CS} \\ + S_T \left[ A_T^{\sin \varphi_S} \sin \varphi_S + D_{[\sin^2 \theta_{CS}]} \left( A_T^{\sin(2\varphi_{CS}-\varphi_S)} \sin(2\varphi_{CS}-\varphi_S) + A_T^{\sin(2\varphi_{CS}+\varphi_S)} \sin(2\varphi_{CS}+\varphi_S) \right) \right] \end{array} \right\}$$

where  $D_{[\sin^2 \theta_{CS}]} = \sin^2 \theta_{CS} / (1 + \cos^2 \theta_{CS})$

		quark	
		<b>U</b>	<b>L</b>
nucleon	<b>U</b>	$f_1^q(x, \mathbf{k}_T^2)$ number density	$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders <b>T-odd</b>
	<b>L</b>	$g_1^q(x, \mathbf{k}_T^2)$ Helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
	<b>T</b>	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers <b>T-odd</b>	$g_{1T}^q(x, \mathbf{k}_T^2)$ Kotzinian-Mulders worm-gear T
		$h_1^q(x, \mathbf{k}_T^2)$ transversity	$h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity


$A_U^{\cos 2\varphi_{CS}} \propto \underline{h}_{1,\pi}^{\perp q} \otimes \underline{h}_{1,p}^{\perp q}$	Boer-Mulders (T-odd)
$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes \underline{f}_{1T,p}^{\perp q}$	Sivers (T-odd)
$A_T^{\sin(2\varphi_{CS}-\varphi_S)} \propto \underline{h}_{1,\pi}^{\perp q} \otimes \underline{h}_{1,p}^q$	Transversity
$A_T^{\sin(2\varphi_{CS}+\varphi_S)} \propto \underline{h}_{1,\pi}^{\perp q} \otimes \underline{h}_{1T,p}^{\perp q}$	Pretzelosity

SIDIS  $\leftrightarrow$  Drell-Yan sign-change of the  
T-odd TMD PDFs  
Fundamental quest: COMPASS, STAR, SpinQuest, LHCspin, etc.




# Polarized SIDIS and Drell-Yan: universality

Semi-inclusive DIS




T-odd TMD PDFs  
↔  
sign change

Drell-Yan process




# Polarized SIDIS and DY – factorization and kinematic regions

## Semi-inclusive DIS



T-odd TMD PDFs  
 sign change

## Drell-Yan process



High  $q_T$  – Collinear factorization  
 Low  $q_T$  – TMD factorization

$$q_T \geq Q$$

Current fragmentation  
 Collinear factorization

High  $x_F$  – Current fragmentation  
 Low  $x_F$  – Target fragmentation

Target fragmentation  
TMD factorization  
 Fracture Functions

Soft region

$$q_T \ll Q$$


Current fragmentation  
TMD factorization

PDFs, FFs

$$x_F$$


# Main TMD tools – universality and synergies

## Semi-inclusive DIS




T-odd TMD PDFs  
sign change

## Drell-Yan process




## Fragmentation Functions

## Electron-positron annihilation



## Parton Distribution Functions

## pp, pA-scattering, jet production, etc.




Cleanest access to hadronization/fragmentation

Hybrid collinear-TMD approach. The wealth of pp data allows studies of:  
TMD universality, evolution, expected factorization breaking



# Main TMD tools – list of experiments (non exhaustive)

## Semi-inclusive DIS




ePIC Jefferson Lab @ 22 GeV

## Drell-Yan process




AMBER  
Apparatus for Meson and Baryon Experimental Research

## Electron-positron annihilation





## pp, pA-scattering, jet production, etc.




# Main TMD tools – list of experiments (non exhaustive)

## Semi-inclusive DIS




ePIC H Jefferson Lab @ 22 GeV


## Drell-Yan process



## Electron-positron annihilation




## pp, pA-scattering, jet production, etc.





# COMPASS-AMBER-LHCspin FTs at CERN



LHCb  
LHCf




L+C  
spin



physics in common  
dimuon production in  $hh$



L+C R&D  
spin





# COMPASS timeline



- CERN SPS north area – M2 beamline
- Fixed target experiment
- Approved in 1997
- Taking data since 2002 (20 years)
- The Analysis Phase started in 2023

33 institutions from 15 countries: ~ 200 members




4 new groups joined COMPASS in 2023-2024  
UConn (US), AANL (Armenia), NCU (Taiwan), Bochum (Germany)

COMPASS Phase-II addendum  
(d-quark  $h_1$  and PRM $\rightarrow$ AMBER)

Spectroscopy  
SIDIS L/T  
Primakoff  
DVCS(pilot)  
Drell-Yan  
DVCS  
Drell-Yan  
SIDIS T


B. Parsamyan

# COMPASS timeline



- CERN SPS north area – M2 beamline
- Fixed target experiment
- Approved in 1997
- Taking data since 2002 (20 years)
- The Analysis Phase started in 2023

33 institutions from 15 countries: ~ 200 members






# COMPASS Physics Program

This talk



## Nucleon structure


- Hard scattering of  $\mu^\pm$  and  $\pi^-$  off (un)polarized P/D targets
- Inclusive and Semi-Inclusive DIS
- Drell-Yan and J/ $\psi$  production
- Study of nucleon spin structure
  - Longitudinal and Transverse
- Collinear and TMD pictures
- Parton distribution functions and fragmentation functions
- Last COMPASS measurement:  
2022 run – transverse SIDIS



# COMPASS-AMBER timeline




**A<sub>000</sub>BER**  
Apparatus for Meson and Baryon  
Experimental Research



- CERN SPS north area – M2 beamline
- Fixed target experiment
- Approved in 1997
- Taking data since 2002 (20 years)
- The Analysis Phase started in 2023

33 institutions from 15 countries: ~ 200 members



# AMBER timeline




- CERN SPS north area – M2 beamline
- Successor of COMPASS
- Approved in 2019
- Taking data since 2023
- Phase-I is planned to continue after LS3

36 institutions from 14 countries: ~ 150 members








COMPASS++/AMBER  
proposal  
[CERN-SPSC-2019-022](https://cds.cern.ch/record/2704422)



# AMBER measurements 2023-2024: Drell-Yan

## $\pi^\pm, K^\pm$ induced dimuon production: Drell-Yan, J/ $\psi$ (and $\psi'$ )

- Study of pion and kaon PDFs
  - Crucial input for the study of the Emergent Hadron Mass (EHM)
- Possibility to collect unique balanced  $\pi^+/\pi^-$  induced DY data
- Measurement of  $\lambda, \mu$  and  $\nu$  (DY, J/ $\psi$ )
- J/ $\psi$  production mechanisms ( $q\bar{q}, gg$ )
- Carbon and tungsten targets
- Improved vertex/mass resolution
- Updated setup, new TL DAQ



[CERN Accelerator Complex schedule to 2041](#)

# COMPASS experimental setup




## COmmon Muon Proton Apparatus for Structure and Spectroscopy



CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV  $p$  from SPS
  - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
  - $h^-$  beam: 97%  $\pi^-$ , 2%  $K^-$ , 1%  $p$
  - $h^+$  beam: 75%  $\pi^+$ , 24%  $p$ , 1%  $K^+$
- 160 GeV tertiary muon beams
  - $\mu^\pm$  longitudinally polarized




Large-acceptance forward spectrometer

- Precise tracking (350 planes)  
SciFi, Silicon, MicroMegas, GEM, MWPC, DC, Straw, Muon walls
- PID - CEDARs, RICH, calorimeters, MWs

Various targets:

- Polarized solid-state  $NH_3$  or  ${}^6LiD$
- Liquid  $H_2$
- Solid-state nuclear targets (e.g. Ni, W, Pb)



# COMPASS experimental setup: Phase II (SIDIS program)




## COmmon Muon Proton Apparatus for Structure and Spectroscopy



CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS


- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV  $p$  from SPS
  - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
  - $h^-$  beam: 97%  $\pi^-$ , 2%  $K^-$ , 1%  $p$
  - $h^+$  beam: 75%  $\pi^+$ , 24%  $p$ , 1%  $K^+$
- 160 GeV tertiary muon beams
  - $\mu^+$  longitudinally polarized



- Polarized solid-state  $NH_3$  or  ${}^6LiD$
- Two or three oppositely polarized cells
- Longitudinal and transverse polarization




# AMBER Phase I: $\bar{p}$ cross-section, 2023 setup

## Apparatus for Meson and Baryon Experimental Research



CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS




- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- $h^+$  beam: 60, 80, 100, 160, 190 and 250 GeV/c;  
Intensity:  $25 \cdot 10^3$  h/s
- Beam PID: 2 CEDAR detectors
- Target: He (2023), H/D (2024)
- Data-taking ~2 months/year
- Dedicated trigger and beam-killer systems



vertex distribution



# COMPASS experimental setup: Phase II (DY program)




## COmmon Muon Proton Apparatus for Structure and Spectroscopy



CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS



- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)



- Primary beam - 400 GeV  $p$  from SPS
  - impinging on Be production target (T6)
- 190 GeV secondary hadron beams
  - $h^-$  beam: 97%  $\pi^-$ , 2%  $K^-$ , 1%  $p$
  - $h^+$  beam: 75%  $\pi^+$ , 24%  $p$ , 1%  $K^+$
- 160 GeV tertiary muon beams
  - $\mu^\pm$  longitudinally polarized



## Hadron absorber




# AMBER Phase I-II: DY program setup

## Apparatus for Meson and Baryon Experimental Research






CERN SPS North Area (building 888)

Two-stage spectrometer LAS+SAS

- Large Angle Spectrometer (SM1 magnet)
- Small Angle Spectrometer (SM2 magnet)




- Secondary  $h^\pm$  beam:  $(\pi^\pm, K^\pm, p/\bar{p})$
- Improved beam PID (CEDARs)
  - enabling kaon physics
- Isoscalar target (C 3x25 cm) and W
- Vertex detector: improve Z and  $M_{\mu\mu}$  resolution
- New trigger-less DAQ,
- Revised setup, new detectors



# LHCspin experiment


JINST 3 (2008) S08005  
IJMPA 30 (2015) 1530022

0.45 - 7 TeV




2.76 TeV

pp:  $\sqrt{s} \simeq 29 - 115$  GeV  
Ap:  $\sqrt{s} \simeq 72$  GeV




New collaborators are Welcome!


## Timeline



# SMOG2 (unpolarized data)




beam-gas  
collisions




beam-beam  
collisions

5 mm radius x 200 mm length




## Timeline



2024 2025 2026 2027

SMOG2




2030

LHC LS3  
installation of the  
apparatus at the IR4

LHCb spin R&D

As a group independent  
from LHCb collaboration

LHC Run4  
data taking at the IR4




2033


LHC LS4  
installing the  
target at LHCb

LHCb spin

LHCspin Run5+6  
data taking at LHCb




# SMOG2 (unpolarized data) gas target: H<sub>2</sub>, D<sub>2</sub>, N<sub>2</sub>, O<sub>2</sub>, He, Ne, Ar, Kr, Xe




Negligible impact on the beam lifetime ( $\tau_{beam-gas}^{p-H_2} \sim 2000$  days,  $\tau_{beam-gas}^{Pb-Ar} \sim 500$  h)

## Timeline




# SMOG2 (unpolarized data)





Negligible impact on the beam lifetime


## Timeline



# Towards polarized measurements at LHCb: IR4




Install a compact “LHCb independent” apparatus in IR4  
purpose: R&D to have a “plug & play” PGT for Run5  
perform unique first ever physics measurements  
perform service measurements related to LHC




# LHCspin experiment

[arXiv:1807.00603]




of course, the Sivers effect / sign-change

## Timeline



# LHCspin experiment


[arXiv:1807.00603]






arXiv:1807.00603


Channel	Events / week	Total yield
$J/\psi \rightarrow \mu^+ \mu^-$	$1.3 \times 10^7$ !!	$1.5 \times 10^9$
$D^0 \rightarrow K^- \pi^+$	$6.5 \times 10^7$	$7.8 \times 10^9$
$\psi(2S) \rightarrow \mu^+ \mu^-$	$2.3 \times 10^5$	$2.8 \times 10^7$
$J/\psi J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$ (DPS)	8.5	$1.0 \times 10^3$
$J/\psi J/\psi \rightarrow \mu^+ \mu^- \mu^+ \mu^-$ (SPS)	$2.5 \times 10^1$	$3.1 \times 10^3$
Drell Yan ( $5 < M_{\mu\mu} < 9 \text{ GeV}$ )	$7.4 \times 10^3$	$8.8 \times 10^5$
$\Upsilon \rightarrow \mu^+ \mu^-$	$5.6 \times 10^3$	$6.7 \times 10^5$
$\Lambda_c^+ \rightarrow p K^- \pi^+$	$1.3 \times 10^6$	$1.5 \times 10^8$

- Precise *spin asymmetry* on  $J/\Psi \rightarrow \mu^+ \mu^-$  and  $D^0 \rightarrow K^- \pi^+$  for  $pH^\uparrow$  collisions in just few weeks
- Inclusive quarkonia production in (un)polarized pp interactions - ideal observable to access gTMDs
- Flavor separation using H/D, EMC effect
- Spin physics in heavy-ion collisions and a lot more!








*Drell-Yan*



*SIDIS*

# Single-polarized Drell-Yan cross-section at twist-2 (LO)




$$\frac{d\sigma^{LO}}{dq^4 d\Omega} \propto F_U^1 (1 + \cos^2 \theta_{CS})$$

$$1 + D_{[\sin^2 \theta_{CS}]} A_U^{\cos 2\varphi_{CS}} \cos 2\varphi_{CS}$$
  

$$+ S_L \sin^2 \theta_{CS} A_L^{\sin 2\varphi_{CS}} \sin 2\varphi_{CS}$$
  

$$\times \left\{ \begin{array}{l} A_T^{\sin \varphi_S} \sin \varphi_S \\ + S_T \left[ A_T^{\sin(2\varphi_{CS} - \varphi_S)} \sin(2\varphi_{CS} - \varphi_S) \right. \\ \left. + D_{[\sin^2 \theta_{CS}]} \left( A_T^{\sin(2\varphi_{CS} + \varphi_S)} \sin(2\varphi_{CS} + \varphi_S) \right) \right] \end{array} \right\}$$



$A_U^{\cos 2\varphi_{CS}} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^{\perp q}$	Boer-Mulders (T-odd)
$A_T^{\sin \varphi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$	Sivers (T-odd)
$A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$	Transversity
$A_T^{\sin(2\varphi_{CS} + \varphi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1T,p}^{\perp q}$	Pretzelosity

SIDIS  $\leftrightarrow$  Drell-Yan sign-change of the T-odd TMD PDFs


COMPASS phase-II proposal submitted in 2010 (Drell-Yan, DVCS,...)

Predictions for a large Sivers effect in Drell-Yan and J/psi at COMPASS  $\rightarrow$  sign change test



# Sivers effect: Drell-Yan and J/ $\psi$

Sivers DY TSA

$$A_T^{\sin\phi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$



- The Drell-Yan Sivers asymmetry tends to be positive ( $\sim 1.5$  s.d.)




Sivers SIDIS TSA

$$A_{UT}^{\sin(\phi_h - \phi_s)} \propto f_{1T}^{\perp q} \otimes D_{1q}^h$$

COMPASS proton Sivers measurements


- Clear signal in the matching  $Q^2$  ranges

# Sivers effect: Drell-Yan and J/ $\psi$





Sivers DY TSA

$$A_T^{\sin \phi_S} \propto f_{1,\pi}^q \otimes f_{1T,p}^{\perp q}$$




- The Drell-Yan Sivers asymmetry tends to be positive ( $\sim 1.5$  s.d.)




- J/ $\psi$  Sivers asymmetry is compatible with zero (within  $\sim 1\%$ )
- Predictions for a large Sivers effect in Drell-Yan and J/ $\psi$  at COMPASS
- Hint that J/ $\psi$  production might go via gg fusion in COMPASS?**
- Access to small gluon TMDs?




# Transversity TSA: Drell-Yan and J/ $\psi$






Transversity DY TSA  
 $A_T^{\sin(2\phi_{CS} - \phi_S)} \propto h_{1,\pi}^{\perp q} \otimes h_{1,p}^q$




- The Drell-Yan Transversity asymmetry is negative (~2 s.d.)




# 3D unpolarized Drell-Yan cross section on NH<sub>3</sub> and W



- First new results in 30 years!
- Data from light/heavy targets
  - NH<sub>3</sub>-He, Al, W
  - Nuclear dependence
- 1D/2D/3D representations
- $x_F:q_T:M$
- Unique data to access collinear and TMD distributions  
e.g. pion TMD PDF





# 3D unpolarized Drell-Yan cross section on NH<sub>3</sub> and W



recent global fit and projections for COMPASS

MAP collaboration  
Phys. Rev. D. 107, 014014






- First new results in 30 years!
- Data from light/heavy targets
  - NH<sub>3</sub>-He, Al, W
  - Nuclear dependence
- 1D/2D/3D representations
- $x_F:q_T:M$
- Unique data to access collinear and TMD distributions  
e.g. pion TMD PDF
- To be included in future global fits (MAP, JAM, etc.)

# AMBER – $\pi^\pm, K^\pm$ induced dimuon production on C/W

- Unique complementary measurements:  $\pi^\pm, K^\pm$ 
  - Cross-sections, pion and kaon PDFs
    - Data for both collinear and TMD PDF studies
  - Drell-Yan, J/ $\psi$  and potentially  $\psi'$  channels
  - Study of nuclear effects with C and W


<b>AMBER</b>	75 cm C	190	$\pi^+$	<b>1200000</b>
			$\pi^-$	<b>1800000</b>
<b>J/<math>\psi</math> events</b>	12 cm W	190	p	<b>1500000</b>
			$\pi^+$	<b>500000</b>
<b>J/<math>\psi</math> events</b>	12 cm W	190	$\pi^-$	<b>700000</b>
			p	<b>700000</b>

Experiment	Target type	Beam energy (GeV)	Beam type	Beam intensity (part/sec)	DY mass ( $\text{GeV}/c^2$ )	DY events
E615	20 cm W	252	$\pi^+$	$17.6 \times 10^7$	4.05 – 8.55	5000
			$\pi^-$	$18.6 \times 10^7$		30000
NA3	30 cm H <sub>2</sub>	200	$\pi^+$	$2.0 \times 10^7$	4.1 – 8.5	40
			$\pi^-$	$3.0 \times 10^7$		121
NA10	6 cm Pt	200	$\pi^+$	$2.0 \times 10^7$	4.2 – 8.5	1767
			$\pi^-$	$3.0 \times 10^7$		4961
NA10	120 cm D <sub>2</sub>	286	$\pi^+$	$65 \times 10^7$	4.2 – 8.5	7800
		140	$\pi^-$		4.35 – 8.5	3200
COMPASS 2015 COMPASS 2018	110 cm NH <sub>3</sub>	190	$\pi^-$	$7.0 \times 10^7$	4.3 – 8.5	35000
					4.0 – 8.5	52000
AMBER	75 cm C	190	$\pi^+$	$1.7 \times 10^7$	4.3 – 8.5	<b>21700</b>
		190	$\pi^-$	$6.8 \times 10^7$	4.3 – 8.5	<b>67000</b>
	12 cm W	190	$\pi^+$	$0.4 \times 10^7$	4.3 – 8.5	<b>8300</b>
		190	$\pi^-$	$1.6 \times 10^7$	4.0 – 8.5	<b>11700</b>






$$\Sigma_{val} = \sigma^{\pi^- C} - \sigma^{\pi^+ C}: \text{only valence-valence}$$

$$\Sigma_{sea} = 4\sigma^{\pi^+ C} - \sigma^{\pi^- C}: \text{no valence-valence}$$






# AMBER – $\pi^\pm, K^\pm$ induced dimuon production on C/W




## Compared to COMPASS




- Light isoscalar target (carbon) instead of  $\text{NH}_3\text{-He}$  mix
- Improved mass resolution ( $\sim 100 \text{ MeV}/c^2$ )
  - Lower background  $\rightarrow$  enlarge DY mass range
  - $J/\psi$  and  $\psi(2S)$  studies
- Wider beam choice:  $\pi^\pm, K^\pm, p/\bar{p}$ , CEDARs (PID)
- Unique complementary measurements:  $\pi^\pm, K^\pm$
- Higher beam intensity (RP upgrades)
- Revised spectrometer, Triggerless DAQ



# LHC – SMOG2 early data – fantastic precision



A wealth of unique data collected in 2024






- Recent SIDIS highlights from COMPASS

# Hadron multiplicities; $h^\pm$ , $\pi^\pm$ and $K^\pm$ (2016 data) collinear

A set of complex corrections:

- Acceptance, rad. corrections,  
PID, diffractive VMs, etc.



New radiative corrections (DJANGOH)


[hep-ex/2410.12005](https://arxiv.org/abs/hep-ex/2410.12005) soon in PRD

# Hadron multiplicities; $h^\pm$ , $\pi^\pm$ and $K^\pm$ (2016 data)

collinear


A set of complex corrections:

- Acceptance, rad. corrections, PID, diffractive VMs, etc.



New radiative corrections (DJANGOH)

[hep-ex/2410.12005](https://arxiv.org/abs/hep-ex/2410.12005) soon in PRD



## Diffractive VM production

- In DIS  $\gamma^*$  interacts with a single quark
- DVMP -  $\gamma^*$  fluctuates into a VM
  - VM then interacts diffractively with the nucleon through multiple gluon exchange
- DVMP correction: two MC samples are used
- SIDIS
- LEPTO 6.5 MC (diffractive contributions off)

## Diffractive $\rho^0$ and $\phi$ mesons

- HEPGEN generator

Diffractive events enhance at low  $x$  and  $Q^2$

- Pions from  $\rho^0$  decay (at high  $z$ )

For pions maximum correction can reach even 50%

- Kaons from  $\phi$  decay ( $0.4 < z < 0.6$ )

For kaons maximum correction ~24%

for ( $z \approx 0.6$  and  $Q^2 \approx 1$  (GeV/c) $^2$ ).

# Cahn effect in SIDIS: Diffractive VMs contribution

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$$f_1^q(x, k_T^2)$$



number density

As of 1978 – simplistic kinematic effect:

- non-zero  $k_T$  induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments



$$F_{UU}^{\cos\phi_h} = \frac{2M}{Q} C \left\{ -\frac{\hat{h} \cdot p_T}{M_h} \left( \cancel{x} \cancel{h} H_{1q}^{\perp h} + \frac{M_h}{M} f_1^q \frac{\tilde{D}_q^{\perp h}}{z} \right) - \frac{\hat{h} \cdot k_T}{M} \left( \cancel{x} \cancel{f}^{\perp q} D_{1q}^h + \frac{M_h}{M} h_1^{\perp q} \frac{\tilde{H}_q^h}{z} \right) \right\}$$

# Cahn effect in SIDIS: Diffractive VMs contribution

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_h d\phi_s} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$$f_1^q(x, k_T^2)$$




number density

As of 1978 – simplistic kinematic effect:


- non-zero  $k_T$  induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
  - So far, no comprehensive interpretation
- A set of complex corrections:
  - Acceptance, diffractively produced VMs, radiative corrections (RC), etc.



VM fractions



# Cahn effect in SIDIS: DVMs

COMPASS, EPJC (2023) 83 924

SDMEs

$\gamma^* \rightarrow \rho^0$  spin components

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect


$f_1^q(x, k_T^2)$   
number density

As of 1978 – simplistic kinematic effect:

- non-zero  $k_T$  induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
  - So far, no comprehensive interpretation
- A set of complex corrections:
  - Acceptance, diffractively produced VMs, radiative corrections (RC), etc.



Kinematic dependences of SDMEs  
Measured (1D), not yet implemented in HEPgen

# Cahn effect in SIDIS: DVMs

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$   
number density


As of 1978 – simplistic kinematic effect:

- non-zero  $k_T$  induces an azimuthal modulation


“Visible” hadron pairs

- Both hadrons reconstructed
- Removed by selecting:  $z_t = z_{h+} + z_{h-} < 0.95$

COMPASS, EPJC (2023) 83 924



VM contribution “amplitudes”



Only “average” SDMEs are implemented in HEPgen  
They seem to describe the data well

# Cahn effect in SIDIS: DVMs

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_S} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$   
number density

As of 1978 – simplistic kinematic effect:

- non-zero  $k_T$  induces an azimuthal modulation


“Visible” hadron pairs

- Both hadrons reconstructed
- Removed by selecting:  $z_t = z_{h+} + z_{h-} < 0.95$


“Invisible” hadron pairs

- Only one hadron out of two is reconstructed
- Subtraction done at the level of  $\phi_h$  using simulated HEPGEN distribution for VMs
- HEPGEN and Lepto are normalized to the data using  $E_{\text{miss}}$  distribution, SIDIS tail is subtracted

COMPASS, EPJC (2023) 83 924



VM corrections, applied




# Cahn effect in SIDIS: DVMs and RCs

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \times (1 + \underbrace{\sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h}_{\text{Cahn effect}} + \dots)$$



Cahn effect

$f_1^q(x, k_T^2)$   
number density




As of 1978 – simplistic kinematic effect:

- non-zero  $k_T$  induces an azimuthal modulation

As of 2023 – complex SF (twist-2/3 functions)

- Measurements by different experiments
- Complex multi-D kinematic dependences
  - So far, no comprehensive interpretation
- A set of complex corrections:
  - Acceptance, diffractively produced VMs, radiative corrections (RC), etc.
- Strong  $Q^2$  dependence – unexplained
  - Do not seem to come from RCs
  - Transition TMD  $\leftrightarrow$  collinear regions?


VM corrections, applied



# Azimuthal effects in unpolarized SIDIS

$$\frac{d\sigma}{dxdydzdp_T^2d\phi_hd\phi_s} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L}) \\ \times (1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h + \dots)$$

Target spin independent part of the cross-section: three asymmetries




Cahn effect  
Different for  $h^+$ ,  $h^-$   
Non-trivial  $Q^2$  dependence

Boer-Mulders effect  
Collins-like behavior  
( $h^+h^-$  - mirror symmetry)

Beam-spin asymmetry  
higher-twist effect  
non-zero, positive trend

Working on 3D kinematic dependences

# SIDIS TSAs: Collins effect and Transversity





$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[ -\frac{\hat{h} \cdot p_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results HERMES/COMPASS  
( $Q^2$  is different by a factor of  $\sim 2$ -3)
- New deuteron data crucial to constrain  $d$ -quark transversity




Point-by-point extraction framework  
A. Martin et al. PRD 91, 014034 (2015)  
A. Martin et al. PRD 95, 094024 (2017)

COMPASS 2022 run – highly successful data-taking!

- 2<sup>nd</sup> COMPASS deuteron measurements conducted in 2022: unique SIDIS data for the next decades

# SIDIS TSAs: Collins effect for $K^0$









$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) + \dots \right\}$$


$$F_{UT}^{\sin(\phi_h + \phi_s)} = C \left[ -\frac{\hat{\mathbf{h}} \cdot \mathbf{p}_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results HERMES/COMPASS  
( $Q^2$  is different by a factor of  $\sim 2$ -3)
- New deuteron data crucial to constrain  $d$ -quark transversity



# Dihadron Collins effect and Transversity





$$\frac{d^7 \sigma}{d \cos \theta d M_{hh} d \phi_R d z d x d y d \phi_S} =$$

$$\frac{\alpha^2}{2\pi Q^2 y} \left( (1 - y + \frac{y^2}{2}) \sum_q e_q^2 f_1^q(x) D_{1,q}(z, M_{hh}^2, \cos \theta) + \right.$$

$$\left. S_\perp (1 - y) \sum_q e_q^2 \frac{|\mathbf{p}_1 - \mathbf{p}_2|}{2M_{hh}} \sin \theta \sin \phi_{RS} h_1^q(x) H_{1,q}^\triangleleft(z, M_{hh}^2, \cos \theta) \right)$$


$$A_{UT}^{\sin \phi_{RS}} = \frac{|\mathbf{p}_1 - \mathbf{p}_2|}{2M_{hh}} \frac{\sum_q e_q^2 h_1^q(x) H_{1,q}^\triangleleft(z, M_{hh}^2, \cos \theta)}{\sum_q e_q^2 f_1^q(x) D_{1,q}(z, M_{hh}^2, \cos \theta)}$$



COMPASS 2022 run – highly successful data-taking!

- 2<sup>nd</sup> COMPASS deuteron measurements conducted in 2022: unique SIDIS data for the next decades
- New results – dihadron Collins-like asymmetries
- Access to collinear transversity PDF; Non-zero trend at large  $x$
- Precision comparable with proton results


# COMPASS 2022 run: new unique deuteron data




## Pavia group fits

Bacchetta, Delcarro, Pisano, Radici, *in preparation*

analysis of statistical error  
with replica method (200)  
68% confidence level




JAM Collaboration, PRD **106** (2022) 3, 034014



**COMPASS 2022 deuteron run**

S. Bhattacharya, Z. B. Kang, A. Metz, G. Penn and D. Pitonyak  
PRD **105** (2022) 3, 034007



# Conclusions

- COMPASS - longest-running CERN experiment (20 years of data-taking)
- A wealth of Spectroscopy, (SI)DIS, Drell-Yan, DVCS, HEMP data collected across the years
  - Petabytes of data available for analysis
- Wide and unique kinematic domain accessing low  $x$  and large  $Q^2$ 
  - Will remain unique for at least another decade
  - Highly successful SIDIS deuteron in 2022 run, promising preliminary results
- Since 2023 the experiment entered the Analysis Phase (4 new groups joined the Analysis Phase)
  - The spectrometer has been transferred to the AMBER collaboration
- AMBER took its first data in 2023-2024 (Antiproton production) and is preparing for the PRM run
- AMBER phase one comprises unique Drell-Yan measurements (after LS3)
- Long AMBER program is being developed: Phase-II proposal is being drafted
- LHCspin is a project for an FT experiment at LHCb operating with a polarized gas target
  - Successful data collection and proof of principle using SMOG2 unpolarized target
  - Important knowledge and wealth of data collected
  - R&D project in scope of small experiment in IR4 is planned to prepare the polarized target
  - LHCspin at LHCb to be operational during run 5.
- Altogether these FTs at CERN provide unique set of unprecedented measurements complementary to EIC, JLab22, NICA-SPD, etc.


# SIDIS x-section and TMDs at twist-2

$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_s} = \left[ \frac{\alpha}{xyQ^2} \frac{y^2}{2(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x} \right) \right] (F_{UU,T} + \varepsilon F_{UU,L})$$

$$1 + \sqrt{2\varepsilon(1+\varepsilon)} A_{UU}^{\cos\phi_h} \cos\phi_h + \varepsilon A_{UU}^{\cos 2\phi_h} \cos 2\phi_h \\ + \lambda \sqrt{2\varepsilon(1-\varepsilon)} A_{LU}^{\sin\phi_h} \sin\phi_h \\ + S_L \left[ \sqrt{2\varepsilon(1+\varepsilon)} A_{UL}^{\sin\phi_h} \sin\phi_h + \varepsilon A_{UL}^{\sin 2\phi_h} \sin 2\phi_h \right] \\ + S_L \lambda \left[ \sqrt{1-\varepsilon^2} A_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} A_{LL}^{\cos\phi_h} \cos\phi_h \right]$$



$$\times \left\{ + S_T \left[ \begin{array}{l} A_{UT}^{\sin(\phi_h - \phi_s)} \sin(\phi_h - \phi_s) \\ + \varepsilon A_{UT}^{\sin(\phi_h + \phi_s)} \sin(\phi_h + \phi_s) \\ + \varepsilon A_{UT}^{\sin(3\phi_h - \phi_s)} \sin(3\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin\phi_s} \sin\phi_s \\ + \sqrt{2\varepsilon(1+\varepsilon)} A_{UT}^{\sin(2\phi_h - \phi_s)} \sin(2\phi_h - \phi_s) \end{array} \right] \right. \\ \left. + S_T \lambda \left[ \begin{array}{l} \sqrt{(1-\varepsilon^2)} A_{LT}^{\cos(\phi_h - \phi_s)} \cos(\phi_h - \phi_s) \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos\phi_s} \cos\phi_s \\ + \sqrt{2\varepsilon(1-\varepsilon)} A_{LT}^{\cos(2\phi_h - \phi_s)} \cos(2\phi_h - \phi_s) \end{array} \right] \right\}$$

	quark		
	<b>U</b>	<b>L</b>	<b>T</b>
<b>U</b>	$f_1^q(x, \mathbf{k}_T^2)$ number density		$h_1^{\perp q}(x, \mathbf{k}_T^2)$ Boer-Mulders <b>T-odd</b>
<b>L</b>		$g_1^q(x, \mathbf{k}_T^2)$ Helicity	$h_{1L}^{\perp q}(x, \mathbf{k}_T^2)$ worm-gear L
<b>T</b>	$f_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ Sivers <b>T-odd</b>	$g_{1T}^q(x, \mathbf{k}_T^2)$ Kotzinian-Mulders worm-gear T	$h_{1T}^q(x, \mathbf{k}_T^2)$ transversity $h_{1T}^{\perp q}(x, \mathbf{k}_T^2)$ pretzelosity





# Polarized SIDIS and DY – factorization and kinematic regions

## Semi-inclusive DIS



JAM, JHEP 04 (2022) 084




# AMBER measurements 2023-2024: $\bar{p}$ production cross-section

## $\bar{p}$ production measurement

- $\bar{p}$  detected in the cosmic rays
  - produced in CR collisions
  - dark matter signature


Understanding the  $\bar{p}$  flux:

- Accurate determination of the CR-component
- Accuracy of the  $\bar{p}$ -production models is at  $\sim 20\%$  level



## Motivation for AMBER 2023-2024 runs

New measurements needed to determine the  $\bar{p}$  -production from cosmic-ray collisions accurately





# AMBER measurements 2023-2024: proton charge radius

## The proton-radius puzzle


- Discrepancies between the charge-radius of the proton extracted from:
  - Electron-proton scattering
  - Hydrogen spectroscopy
  - Muonic-hydrogen spectroscopy

## AMBER PRM


- Elastic muon-proton scattering
  - 100 GeV/c muon beam
  - Active-target Hydrogen TPC for proton detection



# 3D unpolarized Drell-Yan cross section on NH<sub>3</sub> and W




- First new results in 30 years!
- Data from light/heavy targets
  - NH<sub>3</sub>-He, Al, W
  - Nuclear dependence
- 1D/2D/3D representations  
 $x_F:q_T:M$
- Unique data to access collinear and TMD distributions  
e.g. pion TMD PDF



Experiment	target	number of events	systematic uncertainty	datapoints (M, x_F)
COMPASS (2018 data)	NH <sub>3</sub> -He	36000	~5%	110
	Al	6000	~15%	50
	W	43000	~15%	50
NA10	W	155000	6.50%	59
E615	W	36000	16%	168

# SIDIS TSAs: Collins effect and Transversity








$$\frac{d\sigma}{dxdydzdp_T^2 d\phi_h d\phi_S} \propto (F_{UU,T} + \varepsilon F_{UU,L}) \left\{ 1 + \dots + S_T \varepsilon A_{UT}^{\sin(\phi_h + \phi_S)} \sin(\phi_h + \phi_S) + \dots \right\}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = C \left[ -\frac{\hat{h} \cdot \mathbf{p}_T}{M_h} h_1^q H_{1q}^{\perp h} \right]$$



- Measured on P/D in SIDIS and in dihadron SIDIS
- Compatible results HERMES/COMPASS  
( $Q^2$  is different by a factor of  $\sim 2$ -3)
- New deuteron data crucial to constrain  $d$ -quark transversity




Highly successful  
Run in 2022!



# Sivers TMD PDF: sign change


STAR, arXiv:[2308.15496](https://arxiv.org/abs/2308.15496) [hep-ex]

The RHIC Cold QCD program: arXiv:[2302.00605](https://arxiv.org/abs/2302.00605) [nucl-ex]



COMPASS, PRL 133 (2024) 071902



- COMPASS Drell-Yan  
 $M_{\mu\mu} \in [4; 9] \text{ GeV}/c^2$



- SIDIS $\leftrightarrow$ Drell-Yan ( $W, Z$ )  
sign change of T-odd TMD PDFs
- Difficult measurement
    - Low x-section, background
  - Sivers TMD PDF
  - Pioneering measurements
    - COMPASS (Drell-Yan): 2015, 2018
    - STAR ( $W, Z$ ): 2011, 2017, 2022
  - COMPASS data favors the sign change
    - Useful input to constrain the fits





# Boer-Mulders TMD PDF: sign change




$$\text{DY: } A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto - \left\{ h_{1,\pi^-}^{\perp\bar{u}} \otimes h_{1,p}^u \right\} < 0 \Rightarrow h_{1,\pi^-}^{\perp\bar{u}} > 0$$

# Boer-Mulders TMD PDF: sign change



$$\text{DY: } A_T^{\sin(2\varphi_{CS} - \varphi_S)} \propto -\left\{ h_{1,\pi^-}^{\perp\bar{u}} \otimes h_{1,p}^u \right\} < 0 \Rightarrow h_{1,\pi^-}^{\perp\bar{u}} > 0$$







$$\text{DY: } A_T^{\sin 2\varphi_{CS}} \propto \left\{ h_{1,\pi^-}^{\perp\bar{u}} \otimes h_{1,p}^{\perp u} \right\} > 0 \stackrel{\text{sign-change}}{\Leftrightarrow} \text{SIDIS: } h_{1,p}^{\perp u} < 0$$

$h_{1,p}^{\perp u} < 0 \rightarrow \text{SIDIS fits}$   
V. Barone, et al.  
PRD 82 (2010) 114025

- COMPASS data favors proton Boer-Mulders TMD PDF sign-change

# COMPASS→AMBER: Vertex detector improvements



# COMPASS data taking campaigns

Beam	Target	year	Physics program
$\mu^+$	Polarized deuteron ( ${}^6\text{LiD}$ )	2002	
		2003	80% Longitudinal   20% Transverse SIDIS
		2004	
		2006	Longitudinal SIDIS
	Polarized proton ( $\text{NH}_3$ )	2007	50% Longitudinal   50% Transverse SIDIS
$\pi   K   p$	$\text{LH}_2, \text{Ni}, \text{Pb}, \text{W}$	2008 2009	Spectroscopy
$\mu^+$	Polarized proton ( $\text{NH}_3$ )	2010	Transverse SIDIS
		2011	Longitudinal SIDIS
$\pi   K   p$	Ni	2012	Primakoff
$\mu^\pm$	$\text{LH}_2$	2012	Pilot DVCS & HEMP & unpolarized SIDIS
$\pi^-$	Polarized proton ( $\text{NH}_3$ )	2014	Pilot Drell-Yan
		2015	
		2018	Transverse Drell-Yan
$\mu^\pm$	$\text{LH}_2$	2016 2017	DVCS & HEMP & unpolarized SIDIS
$\mu^+$	Polarized deuteron ( ${}^6\text{LiD}$ )	2021 2022	Transverse SIDIS

# CERN LHC and NA schedules

## CERN Accelerator Complex schedule to 2041

