



POLYTECH

Peter the Great
St. Petersburg Polytechnic
University



PNPI

Identified Hadrons Measurements (π , K , p) in XeW@2.5 GeV (MPD-FXT)




M.Malaev, D. Ivanishchev, V.Riabov

* We acknowledge support from Russian Ministry of Education and Science, state assignment for fundamental research
(code FSEG-2025-0009)

Outline




- ❖ Light charged hadrons spectra ($\pi/K/p$) needed
- ❖ Simplified approach based on n-sigma method for TPC/TOF:
 - ✓ limited pT range at higher momenta
 - ✓ minimization of model-dependent corrections
 - ✓ robust \rightarrow most appropriate for the first-day analysis & results
- ❖ Before: results in [BiBi@9.2](#) collisions
- ❖ Today: analysis details and results in [XeW@2.5](#) AGeV (fixed target mode)
- ❖ Data: Request 36, 15M UrQMD events, Xe-W($T = 2.5$ GeV/n, FTX)

Generated spectra



❖ Almost no difference for pions; NO antiprotons; considerable asymmetry for kaons

Normalized Spectra



❖ Spectra are soft \rightarrow low p_T part is very crucial



Fraction of spectrum lost (p_T dependence)

❖ The fraction of total spectrum lost if spectrum starts from p_T in the most left column

St p_T	Proton	Pi+	Pi-	K+	K-
0.05	0.005	0.037	0.037	0.007	0.014
0.10	0.019	0.131	0.134	0.029	0.053
0.15	0.042	0.259	0.266	0.064	0.114
0.20	0.074	0.394	0.404	0.112	0.193
0.25	0.114	0.514	0.526	0.170	0.279
0.30	0.162	0.616	0.630	0.236	0.370
0.35	0.215	0.700	0.713	0.306	0.462
0.40	0.272	0.767	0.779	0.378	0.547
0.45	0.333	0.820	0.831	0.450	0.623
0.50	0.395	0.861	0.870	0.520	0.691
0.55	0.457	0.893	0.901	0.586	0.747
0.60	0.518	0.918	0.924	0.648	0.798
0.65	0.576	0.937	0.942	0.703	0.840
0.70	0.631	0.952	0.956	0.753	0.874
0.75	0.681	0.963	0.966	0.796	0.900
0.80	0.727	0.972	0.974	0.833	0.920
0.85	0.768	0.979	0.980	0.865	0.938
0.90	0.804	0.984	0.985	0.891	0.951
0.95	0.835	0.988	0.989	0.914	0.962
1.00	0.863	0.991	0.991	0.932	0.970



PID strategy (π , P)

- Event selection: centrality 0-90%
- Track selection:
 - TPC-hits > 24
 - DCA-to-PV < $2\sigma_{x,y,z}$
 - $|y_{\text{cms}}| < 0.5$
- Two quasi-independent measurements for π/p :
 - 1st: **(TPC-TOF)**
 - TPC 2σ -PID selection for a given specie (π/p)
 - If track is 2σ -matched to TOF then TOF 2σ -PID selection for a given specie (π/p)
 - 2nd: **(TOF-TPC)**
 - TOF 2σ -PID selection for a given specie (π/p)
 - TPC 2σ -PID selection for a given specie (π/p)
- Spectra are reconstructed while purity > 95%:
 - spectra are corrected for impurities \rightarrow impose 50% uncertainty for the correction value = $0.5 * 5\% = 2.5\%$ pT-correlated systematic uncertainty for spectra
- **TPC-TOF** and **TOF-TPC** spectra are combined for final results for minimum total uncertainties









PID strategy (Kaons)

- Event selection: centrality 0-90%
- Track selection:
 - TPC-hits > 28
 - DCA-to-PV < $2\sigma_{x,y,z}$
 - $|y_{\text{cms}}| < 0.5$
- Two quasi-independent measurements for K:
 - 1st: **(TPC-TOF)**
 - TPC 1σ -PID selection for a given specie (K)
 - If track is 2σ -matched to TOF then TOF 1σ -PID selection for a given specie (K)
 - TPC 3σ -veto-PID for other species (for K - e/ π /p veto)
 - 2nd: **(TOF-TPC)**
 - TOF 1σ -PID selection for a given specie (K)
 - TPC 1σ -PID selection for a given specie (K)
 - TOF 3σ -veto-PID for other species (for K - e/ π /p veto)
- Spectra are reconstructed while purity > 90%:
 - spectra are corrected for impurities \rightarrow impose 50% uncertainty for the correction value = $0.5 * 5\% = 2.5\%$
 - pT-correlated systematic uncertainty for spectra
- **TPC-TOF** and **TOF-TPC** spectra are combined for final results for minimum total uncertainties









Pions (TPC+TOF→TPC; TOF+TPC→TOF)

- ❖ Accepted p_T range is defined by purity $> 95\%$ → whole range is fine for π^- and limits p_T range to ~ 1.2 GeV/c for π^+









Protons

❖ Accepted p_T range is defined by purity $> 95\%$ \rightarrow whole range is fine









Kaons

- ❖ Accepted p_T range is defined by purity $> 90\%$ \rightarrow limits p_T range to ~ 1.2 GeV/c for K^+ and $\sim 0.9-1$ GeV/c for K^-



Particle sources




❖ Fractions of primaries in the measured spectrum (primaries – produced at dist < 1 cm from PV)



❖ Clean raw spectrum



Combined spectra – transition point

❖ Relative statistical uncertainties for **TPC-TOF** and **TOF-TPC** spectra





Set transition point (p_T):



- ✓ K – 0.25 GeV/c
- ✓ P – 0.7 GeV/c
- ✓ π – 0.5 GeV/c





Final Spectra, pions



- ❖ Start at $p_T \sim 50 \text{ MeV}/c$
- ❖ Measured spectra sample $\sim 96\%$ of the total yields, loose 4% at low p_T





Final Spectra, protons





- ❖ Start at $p_T \sim 100 \text{ MeV/c}$
- ❖ Measured spectra sample $\sim 97\%$ of the total yields, loose $\sim 3\%$ at low p_T , high p_T reach limited only by statistics

Final Spectra, kaons



- ❖ Start at $p_T \sim 100(150)$ MeV/c for $K^+(K^-)$
- ❖ Measured spectra sample $\sim 96(85)\%$ of the total yields for $K^+(K^-)$, main losses at low p_T



Summary

- ❖ A very straightforward approach for $\pi/K/p$ measurements is proposed \rightarrow good for the first-day measurements
- ❖ Generated spectra are soft \rightarrow reconstruction of low p_T part is very crucial
- ❖ Losses at high p_T negligible and/or statistic dependent
- ❖ Losses at low p_T is $\sim 3-5\%$ for pions, protons and K^+ , $\sim 15\%$ for K^-