

Probing the *D*euteron short-range *S*pin *S*tructure in the (d,p) reactions using polarized deuteron beam at Nuclotron

DSS - project

V.A.Baskov², E.V.Chernykh¹, I.Dobrin³, V.A.Dronov², D.Enache³, S.G.Genchev⁴, Yu.V.Gurchin¹, A.Yu.Isupov¹, M.Janek⁵, E.V.Karpechev⁶, J.-T.Karachuk³, A.N.Khrenov¹, A.B.Kurepin⁶, N.A.Kurepin⁶, P.K.Kurilkin¹, V.P.Ladygin^{1†}, N.B.Ladygina¹, D.Lipchinski³, A.N.Livanov¹, A.I.Lvov², Y.Maeda⁷, G.Martinska⁸, O.Mezhenska⁸, K.Miki⁹, S.Nakai⁹, S.Ohta¹⁰, S.M.Piyadin¹, V.V.Polansky², J.Popovichi³, S.G.Reznikov¹, D.Sakai⁹, K.Sekiguchi⁹, S.Shibuya⁹, S.S.Sidorin², K.A.Skazytkin⁶, Ya.T.Skhomenko^{1,11}, V.V.Syschenko¹¹, G.Tarjányiová⁵, A.A.Terekhin¹, A.V.Tishevsky¹, T.Uesaka¹², J.Urban⁸, M.Veverichek⁵, A.Watanabe⁹, I.E.Vnukov¹¹, I.S.Volkov^{1,11}.

(1) *Joint Institute for Nuclear Research, Dubna, Russia*

(2) *Lebedev Physical Institute of the Russian Academy of Sciences, Moscow, Russia*

(3) *Advanced Research Institute for Electrical Engineering, Bucharest, Romania*

(4) *Laboratory for Technical Developments and Applications of the BAS, Sofia, Bulgaria*

(5) *Physics Department, University of Žilina, Žilina, Slovak Republic*

(6) *Institute for Nuclear Research of the Russian Academy of Sciences, Moscow, Russia*

(7) *Department of Applied Physics, University of Miyazaki, Miyazaki, Japan*

(8) *P.J.Šafarik University, Košice, Slovak Republic*

(9) *Department of Physics, Tohoku University, Sendai, Japan*

(10) *Center for Nuclear Study, University of Tokyo, Tokyo, Japan*

(11) *Belgorod State National Research University, Belgorod, Russia*

(12) *RIKEN Nishina Center, Saitama, Japan*

† *Contact person: vladygin@jinr.ru*

Project Leaders: M.Janek⁵, V.P.Ladygin¹, K.Sekiguchi⁹

Abstract

The main goal of the project is to study the deuteron and three-nucleon systems short-range spin structure via the measurements of the polarization observables in the dp -, pd - and dd - reactions at Nuclotron.

The beam energy scan of the analyzing powers A_y , A_{yy} and A_{xx} in dp - elastic scattering between 400 MeV and 1800 MeV has been performed in 2016-2017. The commissioning measurements of the analyzing powers iT_{11} and T_{20} in dp - non-mesonic breakup at 400 MeV in the complanar geometry has been also performed.

In the next term DSS collaboration is planning to measure the deuteron analyzing powers A_y , A_{yy} and A_{xx} in dp - non-mesonic breakup for different kinematic conditions at 300-500 MeV. The energy scan of the deuteron A_y , A_{yy} and A_{xx} analyzing powers in dp - elastic scattering in the deuteron energy range of 300-700 MeV with the step of 50 MeV is also planned. The availability of the polarized proton beam allows to perform the measurements of the nucleon analyzing power A_y^p in pd - non-mesonic breakup for different kinematic conditions at 150-250 MeV/nucleon, as well as the energy scan of the proton analyzing power A_y^p in pd - elastic scattering at 150-1000 MeV/nucleon at Internal Target Station.

The feasibility study of dd - collisions at the intermediate energies at Internal Target Station are planned since these investigations can provide the important information of the structure of the three- and four- nucleons forces.

In the framework of the project the deuteron beam polarimetry will be developed at Internal Target Station and at the extracted deuteron beam at Nuclotron. R&D works on the deuteron and proton polarimeters for NICA and other polarization techniques are planned.

1 Introduction

The main goal of the *DSS* experiment is to study the spin structure of two-nucleon (2N) and three-nucleon (3N) short-range correlations via the measurements of the polarization observables in the deuteron induced reactions (*pd*-, *dp*- and *dd*- interactions) at Nuclotron.

Short range correlations (SRC) of nucleons in nuclei is the subject of intensive theoretical and experimental works during last years. Since SRC have densities comparable to the density in the center of a nucleon which is about $\rho \sim 5\rho_0$ ($\rho_0 \approx 0.17 \text{ fm}^{-3}$), they can be considered as the drops of **cold dense nuclear matter** [1]. These studies explore a new part of the phase diagram and very essential to understand the evolution of neutron stars.

The results obtained at BNL [2], SLAC [3] and JLAB [4, 5] clearly demonstrate that: (i) more than 90% all nucleons with momenta $k \geq 300 \text{ MeV}/c$ belong to 2N SRC; (ii) probability for a given proton with momenta $300 \leq k \leq 600 \text{ MeV}/c$ to belong to *pn* correlation is ~ 18 times larger than for *pp* correlations; (iii) probability for a nucleon to have momentum $\geq 300 \text{ MeV}/c$ in medium nuclei is $\sim 25\%$; (iv) 3N SRC are present in nuclei with a significant probability [6]. However, still many open questions persist and further investigations are required both from the experimental and theoretical sides. For instance, the experimental data on the spin structure of 2N ($I=1$) and 3N SRC are almost absent.

The main tools to study SRCs at hadronic facilities can be deuteron structure investigations at large internal momenta allowing to explore 2N SRC with $I = 0$; ^3He structure to understand the role of 2N SRC with $I = 1$ and 3N SRC; nuclei breakup $A(p, pp)X$, $A(p, pn)X$, $A(p, ppp)X$ etc. with the detection of few nucleons in the final state. The greatest importance is the study of the spin effects in these reactions because the data on the SRCs spin structure are scarce. Nuclotron and NICA will allow to investigate the spin effects for multi-nucleon correlations in a wide energy range.

The model of 2N and 3N correlations at low and moderate energies (below pion threshold production) can be built from the boson-nucleon picture of strong interaction. During last several years a new generation of nucleon-nucleon potentials are built (Nijmegen, CD-Bonn, AV-18 etc.). These potentials reproduced the NN scattering data up to 350 MeV with very good accuracy. But these potentials cannot reproduce triton binding energy (underbinding is 0.8 MeV for CD-Bonn), deuteron-proton elastic scattering and breakup data. Incorporation of three nucleon forces (3NF), when the interaction depends on the quantum numbers of the all three nucleons, allows to reproduce triton binding energy and unpolarized deuteron-proton elastic scattering and breakup data (see [7] and references

therein). The contribution of 3NF is found to be up to 30% in the vicinity of Sagara discrepancy for deuteron-proton elastic scattering at intermediate energies [8, 9]. However, the use of different 3NF models in Faddeev calculations can not reproduce polarization data intensively accumulated during last decade at different facilities [8]–[14].

On the other hand, pd - elastic scattering cross section data obtained already at 250 MeV [11] can not be reproduced by the Faddeev calculations with the inclusion of modern 3NF. The authors stated that the reason of this discrepancy can be neglecting by new type of short-range 3NF. At higher energies, Faddeev calculations fail to reproduce the cross section at the angles larger than 90° . The relativistic multiple scattering calculations [15, 16] give much better agreement with the data at the angles between 60° and 130° . It is shown that the double scattering dominates over the single scattering starting from $\sim 70^\circ$. The deviation of the data on the calculations at backward angles are related with the manifestation of s - type of Fujita-Miyazawa 3NF. Some discrepancy exists around 90° , which can be connected with new type of SR 3NF. These forces can be built within approaches beyond one-boson-exchange. For instance, in the dressed bag model [17] 3NF comes from the interaction between intermediate six-quark state dressed by σ -field and the third nucleon. The description of 2N and 3N correlations at the energies higher than several hundreds MeV/nucleon should be obtained within QCD [1].

The scientific program of the *DSS* project for 2016-2018 included several directions.

- The systematic measurements of the cross section, A_y , A_{yy} and A_{xx} analyzing powers in dp - elastic scattering at the energies 270-2000 MeV [18, 19] at Internal Target Station (ITS) [20] at Nuclotron.
- The measurements of the cross section and analyzing powers in dp - non-mesonic breakup at the energies below 500 MeV for different kinematic configurations of two final protons.
- The start of the measurements of the polarization observables in the ${}^3\text{He}(d, p){}^4\text{He}$ reaction with new PIS (depending on the availability of polarized deuteron beam and ${}^3\text{He}$ target).
- The calibration of the ITS polarimeter at the energies 270-2000 MeV [21] with the expected error bars for analyzing powers ± 0.02 . The calibration of the external polarimeter at the energy 1600 MeV.
- The continuation of the theoretical analysis of the observables in hadronic reactions with the participation of light nuclei.

The cross section measurements has been performed using unpolarized beam at Nuclotron. The polarization measurements at ITS has been performed using rather low intensity of polarized deuteron beam. Actually, these measurements were be performed during the commissioning run with new PIS [22]. The priority was given to the measurements of the angular dependence of A_y , A_{yy} and A_{xx} analyzing powers in dp - elastic scattering at the energies between 700 and 2000 MeV.

The experiment has been approved in June 2009, prolonged in June 2011 for the period 2013-2015 and in June 2014 for the period 2016-2018 with the first priority. During last period **polarized deuteron and proton beams** with the intensity up to $5 \cdot 10^8$ and $2.5 \cdot 10^7$ ppp at ITS were available.

2 Experiment DSS at Internal Target Station

At intermediate energies, intensive studies on three-nucleon force effects are proceeded via the deuteron-proton and deuteron-neutron elastic scattering and breakup reactions. At energies $T_d \leq 300$ MeV, Faddeev calculations can provide good description of cross section data, by introducing a three nucleon force, over the whole angular range. However, their reproduction of polarization observables are not as good as that for cross section data, which may be regarded as an insufficiency of our knowledge on spin-dependence of three nucleon force. To clarify the situation of the spin-dependence, it is undoubtedly promising to perform systematic study of polarization observables in the region of so called "cross section minimum" which spans $\theta_{\text{cm}} = 90^\circ\text{--}130^\circ$. At higher energies, for example $T_d = 500$ MeV, the Faddeev calculation deviates from the cross section data at backward angles. This may be due to relativistic effects and/or new three-nucleon force other than Fujita-Miyazawa type. Subthreshold pion production may play a role. At higher energies 2N and 3N SRC can manifest. Measurement of energy dependences of polarization observables in the region of cross section minimum can give an irreplaceable clue to the problem.

The ITS setup is well suited for study of energy dependence of polarization observables for the deuteron-proton elastic scattering and deuteron breakup reaction with the detection of two protons at large angles (see Fig.1).

The measurements of the cross section, deuteron analyzing powers A_y , A_{yy} and A_{xx} in dp - elastic scattering at $T_d = 300\text{--}2000$ MeV in the vicinity of cross section minimum using Internal Target Station (ITS) at Nuclotron has been proposed [19]. Earlier the measurements of the cross section and analyzing powers for the both dp - elastic scattering and dp - breakup reaction have been proposed at $T_d = 300\text{--}500$ MeV [18].




Figure 1: Schematic view of the ITS [20] and detection system for dp - elastic scattering studies.

3 Experimental results obtained at ITS in 2015-2018

The target CH_2 of 10 mkm thick is used for the measurements at ITS. The yield from carbon content of the CH_2 target is estimated in separate measurements using carbon wire. The monitoring of the intensity is done from the detection of pp - quasielastic scattering at 90° in cms by the scintillation counters placed in the horizontal plane. The detection of the dp - elastic events is done by the coincidence measurements of the proton and deuteron. The detectors are placed in the horizontal plane only for the cross section measurements and in the both horizontal and vertical planes for the analyzing powers measurements. The selection of the dp - elastic events is done by the correlation of the energy losses in plastic scintillators for deuteron and proton and their time-of-flight difference. The interaction point for each event is reconstructed by the target position monitor [23] and used in the data analysis.

3.1 Cross section dp - elastic scattering at 1000-1800 MeV

The procedure of the $CH_2 - C$ subtraction for scattering angle $\theta_{cm} = 75^\circ$ at 500 MeV/n is shown in Fig. 2. The CH_2 distribution and the normalized C spectrum are shown in the left panel by the solid- and dashed-line histograms, respectively. The results of the $CH_2 - C$ subtraction is given in the right panel of Fig. 2.

The measurements of the cross section have been performed at as a first stage of these studies Nuclotron. Fig.3 demonstrates the angular behaviour of the deuteron-proton elastic scattering cross section. The solid symbols are the data obtained at Nuclotron at 700 MeV/nucleon [24]. They are compared with the world data obtained at 650 and 800




Figure 2: Procedure of $CH_2 - C$ subtraction for $\theta_{cm} = 75^\circ$ at the energy of 500 MeV/n. A - is the CH_2 - and normalized C - distributions given by the solid and dashed histograms, respectively, vertical solid lines - is the interval of the normalization. B - is the result of $CH_2 - C$ subtraction, vertical dashed lines are the gates indicating the domain of the dp -elastic scattering events.




Figure 3: The angular behaviour of the deuteron-proton elastic scattering cross section. The solid and open symbols are the data obtained at Nuclotron at 700 MeV/nucleon [24] and the world data obtained at 650 and 800 MeV/nucleon. The lines are the calculations performed in the framework of the relativistic multiple scattering model [25].

MeV/nucleon and shown by open symbols. The theoretical calculations were performed in the relativistic multiple scattering expansion formalism [15, 16, 25]. The four contributions are taken into account: one-nucleon-exchange, single- and double- scattering, and Δ -isobar excitation. The presented approach was applied earlier to describe the differential cross sections at deuteron energies between 500 and 1300 MeV in a whole angular range [25]. The dashed line in Fig 3 is the result of the calculations taking into account one-nucleon exchange (ONE) and single-scattering (SS) diagrams only, while the dotted line is obtained when double-scattering (DS) term is added into the consideration [15, 16]. The full calculation with Δ - isobar excitation included is given by the solid line [25]. One can see that the differential cross section is described quite well up to the scattering angles of $\sim 60^\circ$ taking into account only the ONE and SS terms. If we consider the dp -elastic scattering at the angles larger than 60° , it is necessary to include the DS term into consideration. It should be noted the double-scattering contribution into the reaction amplitude increases with the deuteron energy growing and may change the value of the differential cross section on a few orders in comparison with the result obtained without inclusion of the DS-term. The Δ - excitation begins to manifest itself at the angle equal to about 120° and describes the behaviour of the experimental data at the angle above 140° where the differential cross section sharply increases. It has been demonstrated also that the contribution of the Δ - isobar mechanism grows with the initial deuteron energy. It is negligible at $T_d = 500$ MeV and very significant at higher energies [25]. The deviation of the data from the calculations at large angles can be related with the contribution of more complicated reaction mechanisms like three- nucleon SRCs or excitation of heavy baryonic resonances.




Figure 4: The preliminary results on the angular behaviour of the dp - elastic scattering cross section at 500 MeV/nucleon, 750 MeV/nucleon and 900 MeV/nucleon shown in the left, middle and right panels, respectively. The solid and open symbols are the data obtained at Nuclotron and the world data. The dashed and solid lines are the calculations performed in the framework of the relativistic multiple scattering model without and with the Δ -isobar excitation [25].

The preliminary results on the angular behaviour of the dp - elastic scattering cross section at 500 MeV/nucleon, 750 MeV/nucleon and 900 MeV/nucleon shown in the left, middle and right panels of Fig.4, respectively. The data obtained at Nuclotron are shown by the solid symbols are compared with the world data shown by the open symbols. The dashed and solid lines are the calculations performed in the framework of the relativistic multiple scattering model without and with the Δ -isobar excitation [25]. One can see that taking into account the Δ -isobar excitation improves the agreement with the experimental data. However, the deviations are still large.




Figure 5: Differential cross section of dp -elastic scattering at the fixed scattering angle $\theta_{cm} \sim 75^\circ$, $\sim 82^\circ$, $\sim 111^\circ$ as a function of the \sqrt{s} is shown in the left, middle and right panels, respectively. Full symbols are the data obtained at the Nuclotron, open symbols are the world data. Lines are the result of the world data approximated by the function $\sim s^{-16}$.

At high energies and large transverse momenta the constituent counting rules (CCR) [26] predict a $1/s^{n-2}$ dependence of the differential cross section for the binary reaction, where n is the total number of the fundamental constituents involved in the reaction. The differential cross sections at high energy and large transverse momenta for the exclusive processes can be written as:

$$\left(\frac{d\sigma}{dt}\right)_{AB \rightarrow CD} \sim s^{2-n} \cdot f(\cos \theta^*) \quad (1)$$

where n is the total number of the fundamental degrees of freedom (quarks, leptons and photons) involved both in the initial and the final states of the $AB \rightarrow CD$ process, θ^* is the center of mass angle, s and t are the Mandelstam variables. For instance, n is equal to 18 for dp - elastic scattering .

Differential cross section of dp -elastic scattering at the fixed scattering angles in the cms $\theta_{cm} \sim 75^\circ$, $\sim 82^\circ$, $\sim 111^\circ$ as a function of the \sqrt{s} is shown in the left, middle and right panels of Fig.5. Full symbols are the data obtained at the Nuclotron, open symbols are the world data. Lines are the result of the world data approximated by the

function $\sim s^{-16}$ coming from the constituent counting rules (CCR) [26]. The data at fixed scattering angles in the cms are in a qualitative agreement with the behavior of the world data. The discrepancy between data and CCR is decreasing with the scattering angle increasing.

The results on the dp - elastic scattering differential cross section were obtained with unpolarized deuteron beam at Nuclotron at 500 MeV/n, 700 MeV/n [24], 750 MeV/n, 900 MeV/n and 1000 MeV/n [27]. The final paper is under preparation.

3.2 Analyzing powers in dp - elastic scattering data

The upgraded setup at ITS has been used to measure the vector A_y and tensor A_{yy} and A_{xx} analyzing powers in dp - elastic scattering between 400 MeV and 1800 MeV using polarized deuteron beam from new source of polarized ions SPI [22]. These measurements were performed using internal target station at Nuclotron [20] with new control and data acquisition system [28]. The existing setup [21] has been upgraded by new VME based DAQ, new MPod based high voltage system, new system of monitors etc.

New SPI [22] has been used to provide polarized deuteron beam. In the current experiment the spin modes with the maximal ideal values of $(P_z, P_{zz}) = (0,0)$, $(-1/3, +1)$ and $(-1/3, +1)$ were used. The deuteron beam polarization has been measured at 270 MeV [21]. The dp - elastic scattering events at 270 MeV were selected using correlation of the energy losses and time-of-flight difference for deuteron and proton detectors. The values of the beam polarization for different spin have been obtained as weighted averages for 8 scattering angles for dp - elastic scattering in the horizontal plane only. The typical values of the beam polarization were ~ 65 - 75% from the ideal values.

After deuteron beam polarization measurements at 270 MeV, the beam has been accelerated up to the required energy T_d between 400 MeV and 1800 MeV. The scintillation detectors were positioned in the horizontal and vertical plane in accordance with the kinematic of dp - elastic scattering for the investigated energy. The main part of the measurements were performed using CH_2 target. Carbon target was used to estimate the background. The selection of the dp - elastic events is done by the correlation of the energy losses in plastic scintillators for deuteron and proton and their time-of-flight difference. The normalized numbers of dp -elastic scattering events for each spin mode were used to calculate the values of the analyzing powers A_y , A_{yy} and A_{xx} .

The angular dependence of the deuteron vector analyzing power A_y at the deuteron kinetic energy T_d of 400 MeV is presented in the left panel of Fig. 6. The full squares are the results of the present experiment at ITS at Nuclotron. Open symbols are the




Figure 6: The angular dependencies of the deuteron analyzing powers A_y , A_{yy} and A_{xx} at the deuteron kinetic energy T_d of 400 MeV shown in the left, middle and right panels, respectively. The full squares are the preliminary results of the present DSS experiment at ITS at Nuclotron. Open symbols are the world data. Lines are the calculation within relativistic multiple scattering model [15, 16].

data obtained at Saclay and IUCF. Dashed and solid lines are the calculations performed within relativistic multiple scattering model [15, 16] considering ONE+SS terms only and with the DS contribution added, respectively. Note that the contribution of the Δ - isobar mechanism is negligible at this energy [25]. One can see good agreement of new data obtained at Nuclotron with the data from earlier experiments. The relativistic multiple scattering model [15, 16] describes the data up to $\sim 90^\circ$ only, while it fails to reproduce the data at larger angles. The considering of the DS term does not improve the agreement. The angular dependencies of the tensor analyzing powers A_{yy} and A_{xx} at the deuteron kinetic energy T_d of 400 MeV are presented in the middle and right panels of Fig. 6, respectively. The full squares are the results of the present DSS experiment at ITS at Nuclotron. Open circles and triangles are the data obtained at Saclay and IUCF, respectively. The lines are the same as show in the left panel. The consideration of the ONE+SS terms allows to describe the behaviour of the A_{yy} analyzing power up to $\sim 80^\circ$ only. The DS term gives a significant contribution at larger angles, however, its taking into account does not remove the discrepancy of the calculation with the data. The A_{xx} behaviour is not described by the model [15, 16] over the whole angular range. The reason of the deviation can be the neglecting by the 3N SRCs.

The energy dependencies of the vector A_y and tensor A_{yy} analyzing powers at 70° in the cms are presented as a function of the transverse momentum P_T in Figs 7 and 8, respectively. The full circles are the preliminary results of the present experiment. The full squares are the data obtained at ITS at Nuclotron in 2005 [29, 30]. Open symbols are the world data. Both A_y and A_{yy} analyzing powers change the sign at $P_T \sim 600$ MeV/c and have the tendencies at larger P_T to reach the positive and negative constant




Figure 7: The energy dependence of the vector analyzing power A_y at 70° in the cms. The full circles are the preliminary results of the DSS experiment. The full squares are the data obtained at ITS at Nuclotron in 2005 [29, 30]. Open symbols are the world data.




Figure 8: The energy dependence of the tensor analyzing power A_{yy} at 70° in the cms. The full circles are the preliminary results of the DSS experiment. The full squares are the data obtained at ITS at Nuclotron in 2005 [29, 30]. Open symbols are the world data.

values, respectively. These features of the data indicate the serious deviation of the spin structure of the 2N SRCs on the standard description of the nucleon-nucleon interaction. Further theoretical investigations are required to understand the behaviour of the data at large P_T .

3.3 dp - non-mesonic breakup at 400 MeV

The study of dp - breakup reaction in different kinematic configurations gives an opportunity to select the regions of phase space where the observables are sensitive mostly to 2NF or 3NF. The predictions of the tensor A_{yy} analyzing power and cross section for dp - non-mesonic breakup at 400 MeV for different kinematic configurations of the final protons [31] demonstrate such sensitivity to the contribution of 3NF.

The dp - non-mesonic breakup reaction will be investigated using ΔE - E techniques for the detection of two final protons [32, 33]. Monte-Carlo simulation shown the feasibility of such techniques for the energies $T_d \leq 500$ MeV [34]. Each detector consists of 2 scintillation counters: the first and the second ones are with 1 cm and 20 cm scintillators in length, respectively. The diameter of the E -counter scintillator is 10 cm. Two Photomultiplier tubes PMTs-85 are positioned opposite to each other at the outside cylindrical surface of the thin ΔE - scintillator. At the bottom end of the E - scintillator a photomultiplier tube

PMT-63 is positioned. High Voltage system for PMT-63 is described elsewhere [35, 36].

The use 8 ΔE - E detectors simultaneously allows to measure the observables for many kinematic configurations. The angles definition is given in the top panel of Fig.9.




Figure 9: Top: Θ_1 , Θ_2 and Φ_{12} - polar and azimuthal angles of particle scattered from the target. Bottom: Amplitude spectra for the PMT-85 and PMT-63 obtained for the one of the arm in the pp quasi elastic kinematics at 90° cm and deuteron energy of 200 MeV/n.

The energy calibration of the ΔE - E detectors have been performed using pp - quasielastic scattering at 3 energies: 150 MeV/n, 200 MeV/n and 250 MeV/n [37]. Mean amplitudes are obtained from two PMT-85s which are connected to the same ΔE scintillator. Amplitude spectra for the PMT-85 and PMT-63 obtained for one of arm in the pp quasi elastic kinematics at 90° cm and deuteron energy of 200 MeV/n are shown in the bottom panel of Fig. 9. Then PMT-85s and PMT-63 amplitudes are corrected using LED amplitude information. Finally, calibration coefficients are obtained by solving of system of linear equations.

$\Delta E - E$ correlation obtained for the second arm in the pp quasi elastic kinematics at 90° cm and deuteron energy of 200 MeV/n is shown in Fig.10. Dashed curve represents the $\Delta E - E$ correlation obtained from geant4 simulation [38] at the same kinematic configuration. One can see good agreement between obtained data and simulation. Good agreement is also observed at 150 MeV/n and 250 MeV/n. For future data taking calibration can be performed at only one energy, preferably 200 MeV/n, with using $\Delta E - E$ dependence from simulation as calibration curve. E_1 and E_2 energy distributions obtained




Figure 10: Top: $\Delta E - E$ correlation obtained for the one of arm in the pp - quasi elastic kinematics at 90° cm and deuteron energy of 200 MeV/n. Dashed curve represents the $\Delta E - E$ correlation obtained from geant4 simulation [38] at the same kinematic configuration. Bottom: Energy E_1 , E_2 , obtained in the pp quasi elastic kinematics at 90° cm and deuteron energy of 200 MeV/n. Polyethylene spectra is represented by non-shaded histogram, Carbon spectra by shaded one.

in the pp quasi elastic kinematics at 90° cm and deuteron energy of 200 MeV/n are shown in Fig.??, also. Polyethylene spectra is represented by non-shaded histogram and Carbon spectra by shaded one. The Carbon contribution for 300, 400 and 500 MeV is relatively small for the case of calibration measurements and the peak position in Polyethylene and subtracted spectra have the same position within experimental accuracy. Therefore, the calibration can be performed on Polyethylene only.

Missing mass spectra with gauss fit (941 ± 11 MeV) obtained by subtracting of Carbon content from Polyethylene one at angles of 31° and 43° for the particles which are stopped in scintillator at deuteron energy of 200 MeV/n are shown in Fig.11. Mean value of missing mass spectra for chosen configuration are in agreement with rest mass of neutron. Value of 11 MeV can be assumed as the energy resolution of the detector.

The analysis of the data obtained in dp - non-mesonic breakup with unpolarized beam for complanar and space-star configurations is in progress [39].

In 2015-2016 all FEU-85 and FEU-63 PMTs have been replaced by the Hamamatsu PMTs. The commissioning data on the vector and tensor analyzing powers in dp - non-mesonic breakup for quasi-elastic and complanar geometry have been obtained at 400 MeV in 2016 [40].




Figure 11: Missing mass spectra with gauss fit (941 ± 11 MeV) obtained by subtracting of Carbon content from Polyethylene one at angles of 31° and 43° for the particles which are stopped in scintillator at deuteron energy of 200 MeV/n.




Figure 12: Angular dependence of the vector analyzing power A_y at energy of 200 MeV/n. Data obtained at Nuclotron at 72.3° and 76.5° in cm are represented by full circles, triangles represent data from [41]. Line is the result of the partial-wave analysis [42].




Figure 13: Polarizations values P_z and P_{zz} for spin modes $(P_z, P_{zz}) = (-1/3, -1)$ and $(-1/3, +1)$ during the runs in 2016 — 2017 yy.

Analyzing power A_y at 72.3° and 76.5° in the cms was measured at 200 MeV/n under pp - quasielastic scattering conditions. Obtained values at 72.3° and 76.5° are 0.10 ± 0.02 and 0.11 ± 0.06 , respectively. Results are in agreement with world pp - elastic scattering data [41] within experimental errors as well as with the result of the partial-wave analysis [42] shown by the solid line in Fig.12.

Values of the vector iT_{11} and tensor T_{20} analyzing powers at polar angles of 34.8° and 36.8° and difference in azimuthal angles of 135° were found to be 0.47 ± 0.10 and 0.02 ± 0.20 , respectively. [40].

Therefore, the feasibility of the studies of the analyzing powers in dp - non-mesonic breakup at intermediate energies at ITS with the detection of two protons has been established.

3.4 Deuteron and proton polarimetry at Nuclotron

The polarimeter based on the use of dp - elastic scattering at large angles ($\theta_{\text{cm}} \geq 60^\circ$) at 270 MeV [21], where precise data on analyzing powers [9, 10] exist, and placed at internal target station (ITS) at Nuclotron [20] has been proposed as the reference deuteron polarimeter at Nuclotron-NICA. The accuracy of the determination of the deuteron beam polarization achieved with this method is better than 2% because of the values of the analyzing powers were obtained for the polarized deuteron beam, which absolute polarization

Table 1: The vector and tensor polarization for different spin modes of SPI [22].

Spin mode(P_z, P_{zz})	P_z	dP_z	P_{zz}	dP_{zz}
(-1/3,+1)	-0.254	0.022	0.637	0.039
(-1/3,-1)	-0.223	0.017	-0.621	0.030
(-2/3,+1)	-0.489	0.026	0.631	0.045
(+2/3,0)	0.427	0.021	0.061	0.037
(0,+1)	0.030	0.027	0.880	0.049
(0,-2)	0.046	0.015	-1.469	0.031

had been calibrated via the $^{12}\text{C}(d, \alpha)^{10}\text{B}^*[2^+]$ reaction [43]. This polarimeter can work in the counting mode in the energy range of 300–2000 MeV and, therefore, can be used for permanent beam polarization monitoring [19].

The vector and tensor polarizations were measured seven, six and four times in the parts at November-2016, at December-2016 and at February-2017, respectively. The values have small statistical and systematics errors. They are rather stable within each part of the experiment. The results of the measurements and approximation are presented in Fig. 13. All the results are within two standard deviations from these constants. One can see that the beam polarization values are quite stable within more than 200 hour of the SPI operation. On the other hand, SPI demonstrates good reproducibility of the polarization values for different sets of the data after long interruptions.

The polarimeter [21] has been used also to tune the SPI [22] operation for pure tensor spin modes (0,-2) and (0,+1), for pure vector spin mode (+2/3,0) and for the spin mode (-2/3,+1) with both vector and tensor components. The preliminary results are presented in table 1. One can see, that the typical values of the beam polarization were $\sim 65\text{-}75\%$ from the ideal values for all 6 spin modes of SPI.

The classical method to measure the proton beam polarization at intermediate and high energies is the use of the left-right pp - elastic or quasi-elastic scattering (see, for instance, [44] and references therein). The maximal value of the analyzing power at the energies below 1000 MeV is close to $\sim 40^\circ$ in cms [42], that corresponds roughly $14\text{-}15^\circ$ in the laboratory. Unfortunately, this angle is inaccessible due to design of ITS and detector support.

This method has been modified to increase the polarimeter figure of merit. For this purpose the measurements of the proton beam polarization using several pairs of the detectors placed in the kinematic coincidences corresponding to pp - elastic scattering in the horizontal plane (orbit plane of Nuclotron) have been proposed. The feasibility of the proposed method at ITS has been checked in November 2016 run at Nuclotron using

polarized deuteron beam with the energy of 500 MeV/nucleon.

The unpolarized and polarized proton beam provided by SPI [22] has been accelerated in March 2017 run up to 500 MeV. The typical intensity of the beam was $\sim 1.5 \cdot 10^8$ ppp and $\sim 2\text{-}3 \cdot 10^7$ ppp for unpolarized and polarized cases, respectively. SPI provided proton beam polarization using WFT 1 \rightarrow 3 with ideal value of the polarization $P=-1$. The polarization of the proton beam has been obtained using the data from eight pairs of the detectors placed in the kinematic coincidences. The values of the analyzing power for pp elastic scattering were taken from SAID PWA [42]. The weighted average values of the proton beam polarization were found as 0.017 ± 0.021 and -0.354 ± 0.022 for unpolarized and polarized cases, respectively. Such value is large enough to perform the measurements of the nucleon analyzing power A_y^p in pd - elastic scattering.

The high energy tensor-vector polarimeters will be based on the dp - elastic scattering at forward angles where both tensor and vector analyzing powers have large values [45, 46]. The possibility to select the dp - elastic scattering events using timing and amplitude information from the scintillation counters detected deuterons and protons in the kinematic coincidence has been demonstrated at 1600 MeV and 2000 MeV [47]. The detection system based on the double sided silicon strip detector developed at LHEP JINR [48] can be used for further removing of the carbon content of the CH_2 target. The useful events will be selected by the proton and deuteron scattering angles correlation and coplanarity condition. These selection criteria will significantly reduce the the contribution from the carbon content of the CH_2 target. The forward deuteron and recoil proton will be detected by two silicon strip detectors in coincidence.

3.5 Analysis of the polarization effects in few-nucleon systems

The relativistic multiple scattering model [15, 16] calculations for dp - elastic scattering has been applied to describe the experimental data between 500 MeV and 2000 MeV. The same approach has been developed for the analysis of the $dd \rightarrow {}^3\text{He}n$ reaction at intermediate energies [49]. This model describes the cross section [50], tensor analyzing power T_{20} [51] and angular dependencies of the deuteron analyzing powers [52, 53, 54]. The model [15, 16] has been developed for the dp - elastic scattering process by taking into account the explicit Δ - isobar excitation [25]. The next step in the model development in 2019-2021 is to consider the ρ - exchange in the Δ - isobar formfactor. This is very important for the description of the polarization data at large scattering angles.

3.6 Publications, talks, thesis

1. The results obtained within *DSS* project in 2015-2018 are published in 10 papers in the regular journals [24],[25], [27], [35], [36], [37], [39], [40], [54], [55] and in 1 electronic preprint [56].
2. The results have been presented at the international conferences DSPIN-2015, HS-2015, IBSHEPP-XXIII, EFB23, MESON2016, DSPIN-2017 and others in ~ 20 talks [57]-[75].
3. During 2015-2018 2 Master Thesis [76, 77] were defended.
4. Several talks has been given at the annual Nuclotron users workshops.

4 Request for 2019-2021.

The realization of the *DSS*- project requires beam time at Nuclotron and other resources from JINR side.

4.1 The schedule

The collaboration is planning in 2019-2021 to focus on

1. the analysis of the data obtained in 2015-2018 yy. with unpolarized (d, C, Ar and Kr) and polarized (d and p) beams.
2. preparation of the measurements and data taking with nuclear beams in 2020-2021 yy.
3. preparation of the polarization measurements and data taking with polarized deuterons and protons in 2021 y.
4. developments for the deuteron and proton polarimetry at Nuclotron and NICA.

DSS collaboration in 2019-2021 yy. will focus on the studies using ITS. The main goal is the measurements of the analysing powers in *dp*- and *pd*- elastic scattering and in *dp*- and *pd*- non-mesonic breakup. We plan also to perform short methodical measurements for the $dd \rightarrow {}^3H p$ at 400 MeV as the first step in the studies of the four-nucleon systems. In parallel the proton and deuteron polarimetry will be developed. For these purposes we expect for 1 or 2 runs with polarized deuteron and proton beams from new PIS [22]) in 2021 y. Some methodical measurements for the development of the polarimetry will be required using ion beams in 2020 y.

4.2 The beam time request

New PIS [22] will allow to have a variety of spin modes. In the experiment we are planning to use the spin modes with the following ideal values of (p_z, p_{zz}) : (0,0), (0,-2), (2/3,0) and (-1/3,+1).

Unfortunately, due to rather low intensity of the polarized deuteron beam, the estimated error bars in the analyzing powers in dp - elastic scattering obtained during 2016-2017 yy. was only about ~ 0.05 (not ~ 0.02 as it was expected). On the other hand, low energy part (300 MeV -700 MeV) was not measured. Therefore, during next term the calibration data at the energies of 300 MeV–700 MeV will be obtained with an interval of 50 MeV. These data are very important to study the 3NF spin structure. The calibration of the deuteron beam ITS polarimeter at high energies will be performed at 1.6 GeV only, where the data on the full set of analyzing powers exist [46], together with the calibration of the polarimeter at the extracted beam.

The required beam time for the calibration was evaluated in ref.[19] for the solid angle of each detector of 2.2 msr. The cross section decreases with a beam energy by a factor of 20 from 270 MeV to 2.0 GeV. This decrease is compensated by an increase of luminosity due to small energy loss of higher energy particle and by an increase of repetition rate of Nuclotron. Consequently, the expected yield is almost independent from the beam energy.

Since we are limited by the use of thin CH_2 target the maximal available intensity is $\sim 5 \cdot 10^9$ /spill. For determination of tensor analyzing power with a statistical uncertainty of $\Delta A = 0.02$, one needs 10^4 counts for one detector. This takes ~ 12 hours for 4 polarization modes including the time for tuning of the beam, changing of the detector position etc. Our request is **100** hours for 2021.

The studies of dp - non-mesonic breakup will be performed after the measurements of dp - elastic scattering. For these purposes we need the maximal available intensity of $\sim 5 \cdot 10^9$ /spill. The major goal is to perform the measurements in the complanar geometry at 400 MeV [31]. The estimated time for this purpose is **100** hours for 2021.

The proton beam polarization was measured in 2017 as ~ 0.35 [74]. This value can be increased by the tuning of the spin precessor of SPI [22]. We are planning to perform the measurements of the nucleon analyzing power A_y^p in pd - elastic scattering in the energy range between 100 MeV/n and 1000 MeV/n using the coincidence techniques. We expect the error bar of $\Delta A_y^p \sim 0.02$ in the measurements at 14 proton energies: 100, 150, 200, 250, 300, 350, 400, 440, 500, 600, 700, 800, 900 and 1000 MeV. The estimated beam time request is **150** hours for the intensity of $\sim 5 \cdot 10^9$ /spill. For this task we need to prepare CD_2 target, new mechanics and new scintillation counters. This task is strongly overlap

with the goal of the proton beam polarimetry based on pp - quasielastic scattering, which will require **50** hours in 2021.

In 2019 we need about 200 hours of the beam time with ions to test the setup for pp - and pd - measurements. Part of this time can be in parasitic mode.

Total beam time request for 2019-2021 is presented in table 2.

Table 2: Beam-time request for DSS - project.

Year	Particles	Time (hours)
2019	-	-
2020	ions	300*
2021	$\vec{d}+\vec{p}$	200+200
TOTAL		700

* part of the beam time in parasitic mode.

4.3 JINR budget request

The JINR budget money request for 2019-2021 is 120k\$ (see table 3). We expect 84k\$ for the equipment (registration electronics, HV system, mechanics, computing etc.) and 36k\$ for the travel and staying expences. Also we expect \sim 30k\$ from RFBR.

5 Conclusions

During 2019-2021 we expect to obtain the following results.

1. The data analysis on the systematic measurements of the cross section, A_y , A_{yy} and A_{xx} analyzing powers in dp - elastic scattering at the energies 270-1800 MeV will be performed. Several papers will be prepared.
2. The data on the deuteron and proton beam polarization during 2016-2017 will be analyzed and published.
3. New detection system for the measurements with polarized protons will be prepared and tested in 2020 y.
4. The deuteron analyzing powers in dp - elastic scattering will be obtained at the energies of 300 MeV–700 MeV with an interval of 50 MeV. These data are very important to study the 3NF spin structure.

5. The vector-tensor polarimeter will be calibrated at 300 MeV–700 MeV and 1600 MeV with the expected error bars for analyzing powers ± 0.02 .
6. First data on the analyzing powers in dp - non-mesonic breakup in the complanar geometry at 200 MeV/n will be obtained.
7. The efficient proton beam polarimetry based on pp - elastic scattering will be developed at the energies of 100 MeV–1000 MeV.
8. The first measurements of the nucleon analyzing power in pd - elastic scattering will be performed at the energies of 100 MeV–1000 MeV.
9. The first measurements of the nucleon analyzing power in dp - non-mesonic breakup in the complanar geometry at 200 MeV/n will be started.
10. The studies of the $dd \rightarrow {}^3\text{H}p$ reaction will be started at 200 MeV/n.
11. The theoretical analysis of the observables in hadronic reactions with the participation of light nuclei will be continued.

References

- [1] L. Frankfurt, M. Sargsian, M. Strikman, Int.J.Mod.Phys.**A23** (2008) 2991.
- [2] E. Piassetzky, M. Sargsian, L. Frankfurt, M. Strikman, J.W. Watson, Phys.Rev.Lett. **97** (2006) 162504.
- [3] L.L. Frankfurt, M.I. Strikman, D.B. Day, M.M. Sargsian, Phys.Rev. **C48** (1993) 2451.
- [4] K.Sh. Egiyan et al., Phys.Rev. **C68** (2003) 014313.
- [5] K.S. Egiyan et al., Phys.Rev.Lett. **96** (2006) 082501.
- [6] L. Frankfurt, M. Sargsian, and M. Strikman, AIP Conf.Proc.**1056**, 322 (2008).
- [7] W. Glöckle, H. Witala, D. Hüber, H. Kamada, J. Golak, Phys.Rep. **274** (1996) 107.
- [8] N. Sakamoto et al., Phys.Lett. **B367** (1996) 60.
- [9] K. Sekiguchi et al., Phys.Rev. **C65** (2002) 034003.
- [10] K. Sekiguchi et al., Phys.Rev. **C70**, 014001 (2004).
- [11] K. Hatanaka, Y. Shimizu et al., Phys.Rev. **C66**, 044002 (2002).

- [12] R. Bieber et al., Phys. Rev. Lett. **84**, 606 (2000).
- [13] K. Ermisch et al., Phys. Rev. Lett. **86**, 5862 (2001).
- [14] K. Ermisch et al., Phys. Rev. **C68**, 051001 (2003).
- [15] N.B. Ladygina, Phys.Atom.Nucl. **71** (2008), 2039; N.B. Ladygina, e-Print: arXiv:0805.3021[nucl-th].
- [16] N.B. Ladygina, Eur.Phys.J. A42 (2009) 91.
- [17] V.I. Kukulin et al., J. Phys. G: Nucl. Part. Phys. **30**, 287 (2004).
- [18] Ladygin V.P.// Proc. of the XV-th Int.Sem. on High Energy Physics Problems, 25-29 September 2000, Dubna, Russia, Eds. A.M.Baldin, V.V.Burov and A.I.Malakhov, Dubna:JINR. 2001. V.2. P.301.
- [19] T. Uesaka, V.P. Ladygin et al., Part.Nucl.Lett. 3[114] N^o5 (2006), 58.
- [20] A.I. Malakhov, et al., Nucl.Instr.Meth. in Phys.Res. **A440** (2000) 320; Yu.S. Anisimov, et al., Proc. of the 7-th Int. Workshop on Relativistic Nuclear Physics, 25-30 August 2003, Stara Lesna, Slovak Republic, p.117.
- [21] P.K. Kurilkin et al., Nucl.Instr.Meth. in Phys.Res. **A642** (2011) 45.
- [22] V.V. Fimushkin et al., J.Phys.Conf.Ser. **678** (2016) 012058.
- [23] Yu.V.Gurchin et al., Phys.Part.Nucl.Lett. **4** (2007) 263.
- [24] A.A.Terekhin et al., Differential Cross Section for Elastic DeuteronProton Scattering at the Energy of 700 MeV per Nucleon, Phys.Atom.Nucl. **80** (2017) 1061.
- [25] N.B.Ladygina, Delta excitation in deuteron-proton elastic scattering, Eur.Phys.J. **A52** (2016) 199.
- [26] V.A. Matveev et al. Lett. Nuovo Cimento **7** (1973) 719.
- [27] A.A.Terekhin et al., Study of the dp-elastic scattering at 2 GeV, Phys.Part.Nucl.Lett. **12** (2015) 695.
- [28] A.Yu.Isupov et al., Nucl.Instrum.Meth. **A698** (2013) 127.
- [29] P.K. Kurilkin et al., Phys.Lett. B **715** (2012) 61.
- [30] P.K. Kurilkin et al., Phys.Part.Nucl.Lett. **8** (2011) 1081.
- [31] J. Kuros-Zolnierchuk et al., nucl-th/0203017v1 (2002).

- [32] S.M. Piyadin et al., Phys.Part.Nucl.Lett. **8** (2011) 107.
- [33] S.M. Piyadin et al., Int.J.Mod.Phys. A26 (2011) 683.
- [34] V.P. Ladygin and S. Nedev, Part. and Nucl. Lett. **2[111]-2002** (2002) 13.
- [35] S.M.Piyadin et al., The PMT high-voltage power system for experiments at the JINR Nuclotron, Prib.Tech.Exp. **59** (2016) 155 (in Russian).
- [36] S.M.Piyadin et al., The system of high-voltage power for the detection equipment of the DSS experiment at the Nuclotron JINR, Phys.Part.Nucl.Lett. **14** (2017) 160.
- [37] M.Janek et al., Calibration procedure of the $\Delta E - E$ detectors for dp breakup investigation at Nuclotron, Phys.Part.Nucl.Lett. **15** (2018) 76.
- [38] M.Janek et al., Phys.Part.Nucl.Lett. **11** (2014) 552.
- [39] M.Janek et al., Investigation of the dp breakup and dp elastic reactions at intermediate energies at Nuclotron, Few Body Syst. **58** (2017) 40.
- [40] M.Janek et al., iT11 and T20 analyzing powers of $dp \rightarrow ppn$ reaction at 400 MeV investigated at Nuclotron, Communications: Scientific Letters of the University of Zilina **19** (2017) 62.
- [41] S.Rahm et al., Phys. Rev. C **57** (1998) 1077.
- [42] <http://gwdac.phys.gwu.edu>
- [43] K. Suda et al., Nucl.Instr.Meth. in Phys.Res. **A572** (2007) 745.
- [44] L.S. Azhgirey et al., Nucl.Instrum.Meth. in Phys.Res. **A497** (2003) 340.
- [45] M. Bleszynski et al., Phys.Lett. **B87** (1979) 178.
- [46] V. Ghazikhanian et al., Phys.Rev. **C43** (1987) 1532.
- [47] Yu.V. Gurchin et al., Phys.Part.Nucl.Lett. **8** (2011) 566.
- [48] A.V. Isaev et al., PTE **1** (2011) 43.
- [49] N.B. Ladygina, $dd \rightarrow {}^3\text{He}n$ reaction at intermediate energies, Few Body Syst. **53** (2012) 253.
- [50] G. Bizard et al., Phys. Rev. **C22** (1980) 1632.
- [51] V.P. Ladygin et al., Phys.Lett. **B589** (2004) 47.

- [52] M. Janek et al., Eur.Phys.J. **A33** (2007) 39.
- [53] A.K. Kurilkin et al., Phys.Rev. **C87** (2013) 051001(R).
- [54] A.K.Kurilkin et al., The vector A_y and tensor A_{yy} , A_{xx} , A_{xz} analyzing powers of the $dd \rightarrow {}^3\text{H}p$ reaction at energy 200 MeV, Phys.Atom.Nucl. **78** (2015) 918.
- [55] P.G.Akishin, A.Yu.Isupov, A.N.Khrenov, P.K.Kurilkin, V.P.Ladygin, S.M.Piyadin, N.D.Topilin, Optimization of a large aperture dipole magnet for baryonic matter studies at Nuclotron, Phys.Part.Nucl.Lett. **12** (2015) 305.
- [56] A.Yu.Isupov et al., Setup for the Nuclotron beam time structure measurements, ePrint: arXiv:1505.05822 [physics.ins-det].
- [57] A.A.Terekhin et al., The differential cross section in the dp-elastic scattering at the energies from 500 to 1000 MeV/nucleon, PoS BaldinISHEPPXXII (2015) 099.
- [58] P.K.Kurilkin et al., Investigation of the light nuclei spin structure from hadronic channels at Nuclotron, PoS BaldinISHEPPXXII (2015) 100.
- [59] S.M.Piyadin et al., Investigation of the dp-non-mesonic breakup reaction at 300-500 MeV at Nuclotron, PoS BaldinISHEPPXXII (2015) 101.
- [60] A.A.Terekhin et al., The study of dp-elastic scattering at the energies from 500 to 1000 MeV/nucleon, Int.J.Mod.Phys.Conf.Ser. **39** (2015) 1560096.
- [61] M.Janek et al., dCH2 and dC spectra obtained for some kinematic configurations at intermediate energies, Int.J.Mod.Phys.Conf.Ser. **39** (2015) 1560095.
- [62] S.M.Piyadin et al., Experimental data on the $dp \rightarrow ppn$ reaction at 300-500 MeV energy of deuteron obtained at ITS at Nuclotron, Int.J.Mod.Phys.Conf.Ser. **39** (2015) 1560097.
- [63] M.Janek et al., Simulation of ${}^3\text{He}(d, p){}^4\text{He}$ reaction and recent results of the experimental investigation of dp breakup reaction, J.Phys.Conf.Ser. **678** (2016) 012032.
- [64] V.P.Ladygin et al., Study of polarization effects at Nuclotron, J.Phys.Conf.Ser. **678** (2016) 012031.
- [65] V.P. Ladygin et al., The Deuteron Beam Polarimetry at Nuclotron-NICA, Int.J.Mod.Phys.Conf.Ser. **40** (2016) 1660087.
- [66] V.P. Ladygin et al., The Spin Studies in Few Body Systems at Nuclotron, Int.J.Mod.Phys.Conf.Ser. **40** (2016) 1660074.

- [67] M.Janek et al., Experimental investigation of $dp \rightarrow ppn$ reaction at intermediate energies at Nuclotron, EPJ Web Conf. **130** (2016) 07007.
- [68] N.B.Ladygina, Delta excitation in dp-elastic scattering at the intermediate energies, EPJ Web Conf. **138** (2017) 08007.
- [69] A.A.Terekhin et al., The differential cross section for dp-elastic scattering at 500-900 MeV/nucleon, EPJ Web Conf. **138** (2017) 03012.
- [70] M.Janek et al., Investigation of reactions using polarized and unpolarized deuteron beam at Nuclotron: Current state and perspectives, EPJ Web Conf. **138** (2017) 01028.
- [71] A.Yu.Isupov, Online polarimetry of the Nuclotron internal deuteron and proton beams, J.Phys.Conf.Ser. **938** (2017) 012019.
- [72] Ya.T.Skhomenko et al., Measurement of the deuteron beam polarization at internal target at Nuclotron for DSS experiment, J.Phys.Conf.Ser. **938** (2017) 012022.
- [73] V.P.Ladygin et al., First results on the energy scan of the vector A_y and tensor A_{yy} and A_{xx} analyzing powers in deuteron-proton elastic scattering at Nuclotron, J.Phys.Conf.Ser. **938** (2017) 012007.
- [74] V.P.Ladygin et al., First results on the measurements of the proton beam polarization at internal target at Nuclotron, J.Phys.Conf.Ser. **938** (2017) 012008.
- [75] M.Janek et al., Dp breakup reaction investigation using polarized and unpolarized deuteron beam, J.Phys.Conf.Ser. **938** (2017) 012005.
- [76] Yu.T.Skhomenko, Master Thesis, Belgorod SNRU , Belgorod (2017).
- [77] A.V.Tishevsky, Master Thesis, Dubna University, Dubna (2017).

Table 3: Estimate of the JINR budget expences in 2019-2021 for the *DSS*- project: Probing the *Deuteron* short-range *Spin Structure* in the (d,p) reactions using polarized deuteron beam at Nuclotron.

N^0	Name of the expences	Total cost	2019	2020	2021
	Direct expences	120.0	45.0	45.0	30.0
1.	Accelerator, Nuclotron	700	0	300*	400
2.	Computer(type)				
3.	Computer network				
4.	Constructor bureau (hours)	300	100	100	100
5.	Model shop (hours)	500	200	200	100
6.	Matherials	40.0	17.0	17.0	6.0
7.	Equipment	44.0	16.0	16.0	12.0
8.	Payment of SEW, made on the agreements				
9.	Travel expences, including	36.0	12.0	12.0	12.0
	a) countries of the non-rouble zone	20.0	7.0	7.0	6.0
	b) countries of the rouble zone	16.0	5.0	5.0	6.0
	c) travels on the contracts				

*- part of the beam time can be delivered in parasitic mode.