# Cosmology: Observations and Hot Big Bang Theory

Dmitry Gorbunov

Institute for Nuclear Research of RAS, Moscow



XXVI Baikal Summer School
on Physics of Elementary Particles and Astrophysics

JINR-ISU,
Irkutsk region, Bol'shie Koty

# Standard Model: Success and Problems

Gauge fields (interactions): $\gamma$, $W^{\pm}$, $Z$, $g$

Three generations of matter: $L = \begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$, $e_R$; $Q = \begin{pmatrix} u_L \\ d_L \end{pmatrix}$, $d_R$, $u_R$

- Describes
  - all experiments dealing with electroweak and strong interactions
- Does not describe
  - Neutrino oscillations
  - Dark matter ($\Omega_{DM}$)
  - Baryon asymmetry ($\Omega_B$)
  - Inflationary stage
  - Dark energy ($\Omega_\Lambda$) dynamical ??
  - Strong CP
  - Gauge hierarchy: ? (No new scales!)
  - Quantum gravity

Try to explain all above

# “Natural” units in particle physics

$$\hbar = c = k_B = 1$$

measured in GeV: energy $E$, mass $M$, temperature $T$

$m_p = 0.938$ GeV, $\quad 1\ \mathrm{K} = 8.6 \times 10^{-14}$ GeV

measured in $\mathrm{GeV}^{-1}$: time $t$, length $L$

$1\ \mathrm{s} = 1.5 \times 10^{24}\ \mathrm{GeV}^{-1}, \quad 1\ \mathrm{cm} = 5.1 \times 10^{13}\ \mathrm{GeV}^{-1}$

Gravity (General Relativity): $V(r) = -G\frac{m_1 m_2}{r}$ $\qquad [G] = M^{-2}$

$M_{\mathrm{Pl}} = 1.2 \times 10^{19}\ \mathrm{GeV} = 22\ \mu\mathrm{g}$ $\qquad G \equiv \frac{1}{M_{\mathrm{Pl}}^2}$

# “Natural” units in cosmology (and astrophyics)

$$1\ \text{Mpc} = 3.1 \times 10^{24}\ \text{cm}$$

$1\ \text{AU} = 1.5 \times 10^{13}$ cm — mean Earth-to-Sun distance

$1\ \text{ly} = 0.95 \times 10^{18}$ cm — distance light travels in one year

$1\ \text{yr} = 3.16 \times 10^{7}$ s

$1\ \text{pc} = 3.3\ \text{ly} = 3.1 \times 10^{18}$ cm — distance to object which has a parallax angle of one arcsec



100 AU — Solar system size

1.3 pc — nearest-to-Sun stars

1 kpc — size of dwarf galaxies

50 kpc — distance to dwarves

0.8 Mpc — distance to Andromeda

1-3 Mpc — size of clusters

15 Mpc — distance to Virgo



# Universe is expanding

## Doppler redshift of light



$L \propto a(t)$

$n \propto a^{-3}(t)$

$H(t) = \frac{\dot{a}(t)}{a(t)}$

Hubble parameter

Hubble Law

$H(t_0)\, r = v_r$

# Expansion: redshift $z$ $\lambda_{abs.}/\lambda_{em.} \equiv 1+z$

## $z \ll 1$ Hubble law : $z = H_0 r$

$$H_0 = h \cdot 100 \frac{\text{km}}{\text{s} \cdot \text{Mpc}} \quad h = 0.705 \pm 0.015$$

standard candles









# Universe is homogeneous and isotropic

redshift

$$z \equiv \frac{\lambda_{\text{detector}}}{\lambda_{\text{source}}} - 1$$




$v_r = c\,z$

# The Universe: age & geometry & energy density

$$[H_0] = L^{-1} = t^{-1}$$

time scale: $t_{H_0} = H_0^{-1} \approx 14 \times 10^9$ yr — age of our Universe

spatial scale: $l_{H_0} = H_0^{-1} \approx 4.3 \times 10^3$ Mpc — size of the visible Universe

$t_{H_0}$ is in agreement with various observations

homogeneity and isotropy in 3d:

flat, spherical or hyperbolic

Observations: "very" flat $l_{H_0}/R_{curv} < 0.1$

order-of-magnitude estimate: $v = rH_0, \quad GM(r)/r^2 \sim G\rho_0 r \sim v^2/r = r/H_0^2$

flat Universe

$$\rho_c = \frac{3}{8\pi} H_0^2 M_{\rm Pl}^2 \approx 0.53 \times 10^{-5} \frac{\rm GeV}{\rm cm^3} \longrightarrow 5 \text{ protons in each } 1\,\text{m}^3$$

# Universe is occupied by “thermal” photons




$T_0 = 2.726$ K

the spectrum (shape and normalization!) is thermal

$n_\gamma = 411\ \mathrm{cm}^{-3}$

# Conclusions from observations

## The Universe is homogeneous, isotropic, hot and expanding...






## Conclusions

- interval between events gets modified

$$\Delta s^2 = c^2 \Delta t^2 - a^2(t) \Delta \mathbf{x}^2$$

in GR expansion is described by the Friedmann equation

$$\left(\frac{\dot{a}}{a}\right)^2 = H^2(t) = \frac{8\pi}{3} G \rho^{\text{energy}}_{\text{density}}$$

$$\rho^{\text{energy}}_{\text{density}} = \rho_{\text{radiation}} + \rho_{\text{matter}} + \dots$$

- in the past the matter density was higher, our Universe was "hotter" filled with electromagnetic plasma

$$\rho_{\text{matter}} \propto 1/a^3(t), \quad \rho_{\text{radiation}} \propto 1/a^4(t), \quad \rho_{\text{curvature}} \propto 1/a^2(t)$$

certainly known up to $T \sim 1\,\text{MeV} \sim 10^{10}\,\text{K}$



# Universe content from astrophysics

## Rotational curves






X-rays from centers of galaxy clusters

## Gravitational lensing



“Bullet” cluster

# Universe content from cosmology

## Standard candles




## Angular distance




## Nucleosynthesis







Large Scale Structures




Baryon acoustic oscillations




CMB anisotropy

# Dark Matter Properties $p = 0$

(If) particles:

1. stable on cosmological time-scale
2. nonrelativistic long before RD/MD-transition (either Cold or Warm, $v_{RD/MD} \lesssim 10^{-3}$ at $T \sim 1$ eV)
3. (almost) collisionless
4. (almost) electrically neutral

## If were in thermal equilibrium: $M_X \gtrsim 1$ keV

If not: for bosons

$\lambda = 2\pi/(M_x v_x)$, in a galaxy $v_x \sim 0.5 \cdot 10^{-3} \longrightarrow M_x \gtrsim 3 \cdot 10^{-22}$ eV

for fermions

Pauli blocking: $M_x \gtrsim 750$ eV

$$f(\mathbf{p},\mathbf{x}) = \frac{\rho_x(\mathbf{x})}{M_x} \cdot \frac{1}{\left(\sqrt{2\pi} M_x v_x\right)^3} \cdot e^{-\frac{\mathbf{p}^2}{2M_x^2 v_x^2}}\Big|_{\mathbf{p}=0} \leq \frac{g_x}{(2\pi)^3}$$

# Key observable: matter perturbations

- CMB is isotropic, but “up to corrections, of course...”
  1. Earth movement with respect to CMB
     $$\frac{\Delta T_{\text{dipole}}}{T} \sim 10^{-3}$$
  2. More complex anisotropy!
     $$\frac{\Delta T}{T} \sim 10^{-4} - 10^{-5}$$
- There were matter inhomogenities $\Delta\rho/\rho \sim \Delta T/T$ at the stage of recombination ($e + p \to \gamma + H^*$)
- Jeans instability in the system of gravitating particles at rest $\Longrightarrow \Delta\rho/\rho \nearrow \Longrightarrow$ galaxies (CDM halos)












# Astrophysical and cosmological data are in agreement




$$\left(\frac{\dot{a}}{a}\right)^2 = H^2(t) = \frac{8\pi}{3} G \rho^{\text{energy}}_{\text{density}}$$

$$\rho^{\text{energy}}_{\text{density}} = \rho_{\text{radiation}} + \rho^{\text{ordinary}}_{\text{matter}} + \rho^{\text{dark}}_{\text{matter}} + \rho_\Lambda$$

$$\rho_{\text{radiation}} \propto 1/a^4(t) \propto T^4(t)\,, \quad \rho_{\text{matter}} \propto 1/a^3(t)$$

$$\rho_\Lambda = \text{const}$$

$$\frac{3H_0^2}{8\pi G} = \rho^{\text{energy}}_{\text{density}}(t_0) \equiv \rho_c \approx 0.53 \times 10^{-5} \frac{\text{GeV}}{\text{cm}^3}$$

radiation: $\Omega_\gamma \equiv \frac{\rho_\gamma}{\rho_c} = 0.5 \times 10^{-4}$

Baryons (H, He): $\Omega_B \equiv \frac{\rho_B}{\rho_c} = 0.046$

Neutrino: $\Omega_\nu \equiv \frac{\sum \rho_{\nu_i}}{\rho_c} < 0.01$

Dark matter: $\Omega_{DM} \equiv \frac{\rho_{DM}}{\rho_c} = 0.28$

Dark energy: $\Omega_\Lambda \equiv \frac{\rho_\Lambda}{\rho_c} = 0.68$


2.7 K
TODAY
14 by
accelerated expansion
4.1 K
matter domination
7.1 by
0.26 eV
recombination
370 ty
$e+p \to H+\gamma$
matter domination
0.76 eV
radiation domination
57 ty
80 keV
3 min
$^3H+^4He \to ^7Li+\gamma$
primordial nucleosynthesis
$^2H+^2H \to n+^3He$
1 MeV
1 s
$p+p \to ^2H+\gamma$
2.5 MeV
neutrino decoupling
0.1 s
200 MeV
QCD phase transition
10 μs
confinement↔free quarks
100 GeV
Electroweak phase transition
0.1 ns
baryogenesis
hot Universe
dark matter production
reheating
inflation

# Friedmann equation for the present Universe

$$H^2 \equiv \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi}{3} G(\rho_{\rm M} + \rho_{rad} + \rho_\Lambda + \rho_{\rm curv})$$

$$\frac{8\pi}{3} G\rho_{\rm curv} = -\frac{\varkappa}{a^2}\,, \quad \rho_c \equiv \frac{3}{8\pi G} H_0^2$$

$$\rho_c = \rho_{\rm M,0} + \rho_{rad,0} + \rho_{\Lambda,0} = \rho_c = 0.53 \cdot 10^{-5} \frac{\rm GeV}{\rm cm^3}\,,$$

$$\Omega_X \equiv \frac{\rho_{X,0}}{\rho_c}$$

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi}{3} G\rho_c \left[\Omega_{\rm M}\left(\frac{a_0}{a}\right)^3 + \Omega_{rad}\left(\frac{a_0}{a}\right)^4 + \Omega_\Lambda\right]$$

# Simple tasks to be solved



1. Refine the estimate of the age of our Universe

$$t_0 = \frac{1}{H_0} = 14 \times 10^9 \,\text{years}$$

2. When ($z_{acc}$, $t_{acc}$, $T_\gamma$ – ?) did deceleration-acceleration transition happen?
3. When ($z_{EQ}$, $t_{EQ}$, $T_\gamma$ – ?) did matter-radiation transition (Equality) happen?

   *Hint: neutrino contribution to radiation energy density is 70% of photon contribution*
4. Find the time of Electroweak phase transition, $T \sim 100\,\text{GeV}$

   *The early Universe to be tested at LHC . . .*

   *Hint: all SM particles contribute to energy density as about 50 photon species*

# Space-dependent quantities in GR are tensors

All coordinate systems are equivalent

$$x^\mu \to x'^\mu = x'^\mu(x^\lambda) \quad \text{but} \quad df(x) = \partial_\mu f\, dx^\mu \quad \textit{is invariant}$$

co-vector: $dx^\mu \to dx'^\mu = \dfrac{\partial x'^\mu}{\partial x^\nu}\, dx^\nu$

contra-vector: $\partial_\mu f \equiv \dfrac{\partial f}{\partial x^\mu} \to \dfrac{\partial f}{\partial x'^\mu} = \dfrac{\partial f}{\partial x^\nu}\dfrac{\partial x^\nu}{\partial x'^\mu} = \partial_\nu f \dfrac{\partial x^\nu}{\partial x'^\mu}$

tensors:

$$A_{\lambda\rho\ldots}{}^{\mu\nu\ldots} \to A'_{\lambda\rho\ldots}{}^{\mu\nu\ldots}$$

invariant interval $ds$ with metric tensor $g_{\mu\nu}$ $\qquad g^{\mu\nu}g_{\nu\lambda} = \delta^\mu_\lambda$

$$ds^2 = dt^2 - \mathbf{dx}^2 = \eta_{\mu\nu}dx^\mu dx^\nu \quad \to \quad g_{\mu\nu}(x)dx^\mu dx^\nu$$

$g \equiv \det g_{\mu\nu}$, invariant measure is $d\Omega^{(4)} \equiv \sqrt{-g}\, d^4x$

$$S = \int \sqrt{-g}\, d^4x \left(\frac{1}{2}g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - \frac{1}{2}m^2\phi^2 - \frac{\lambda}{4}\phi^4\right)$$

# Convention of GR description

## metric $ds^2 = dt^2 - dx^2$

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu \,, \qquad \mu,\nu = 0,1,2,3 \qquad \text{signature } (+,-,-,-)$$

$$x^\mu \to x'^\mu(x^\mu) : \qquad B'^{\mu}_{\nu\lambda}(x') = \frac{\partial x'^\mu}{\partial x^\sigma}\frac{\partial x^\tau}{\partial x'^\nu}\frac{\partial x^\rho}{\partial x'^\lambda}B^\sigma_{\tau\rho}(x)$$

## invariant volume $\int d^4x\sqrt{-g}$

$$\int d^4x \;\longrightarrow\; \int d^4x\sqrt{-g}$$

## $g_{\mu\nu}(x)$ is the dynamical variable $T_{\mu\nu} \propto g_{\mu\nu}\Lambda$

$$\int d^4x\sqrt{-g}\;\Lambda$$

# Dark Energy: nonclumping matter?




- estimates of Matter contribution confined in galaxies and clusters
  $\rho_c - \rho_M \neq 0$ but the Universe is flat, so $\rho_{curv} \simeq 0$
- corrections to the Hubble law : red shift – brightness curves for standard candles (SN Ia)
- The age of the Universe
- CMB anisotropy, large scale structures (galaxy clusters formation), etc

$$\rho_\Lambda = 0.73\rho_c$$

$$\rho_\Lambda \sim 10^{-5}\ \text{GeV/cm}^3 \sim \left(10^{-11.5}\ \text{GeV}\right)^4$$

# Convention of GR description

**metric**

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu\,, \qquad \mu,\nu = 0,1,2,3 \qquad \text{signature } (+,-,-,-)$$

$$x^\mu \to x'^\mu(x^\mu)\,: \qquad B'^\mu_{\nu\lambda}(x') = \frac{\partial x'^\mu}{\partial x^\sigma}\frac{\partial x^\tau}{\partial x'^\nu}\frac{\partial x^\rho}{\partial x'^\lambda}B^\sigma_{\tau\rho}(x)$$

**Christoffel symbols:**

$$\tilde{A}^\mu(\tilde{x}) = A^\mu(x) - \Gamma^\mu_{\nu\lambda}(x)A^\nu(x)dx^\lambda$$

**Covarinat derivative**

$$A^\mu(\tilde{x}) - \tilde{A}^\mu(\tilde{x}) = \nabla_\nu A^\mu \cdot dx^\nu$$

$$\nabla_\nu A^\mu(x) = \partial_\nu A^\mu(x) + \Gamma^\mu_{\nu\lambda}(x)A^\lambda$$

If $\Gamma^\mu_{\nu\lambda} = \Gamma^\mu_{\lambda\nu}$ and $\nabla_\nu g_{\mu\lambda} = 0$ $\qquad A_\mu \equiv A^\nu g_{\mu\nu}$

$$\Gamma^\mu_{\nu\lambda} = \frac{1}{2}g^{\mu\rho}(\partial_\nu g_{\rho\lambda} + \partial_\lambda g_{\rho\nu} - \partial_\rho g_{\nu\lambda})$$

# FLRW metric $g_{\mu\nu}$

$$ds^2 = g_{\mu\nu}dx^\mu dx^\nu = dt^2 - a^2(t)dl^2 = dt^2 - a^2(t)\gamma_{ij}dx^i dx^j \ ,$$

Special frame: different parts look similar

it is comoving: world lines of particles at rest are geodesics,

derive it 

$$\frac{du^\mu}{ds} + \Gamma^\mu_{\nu\lambda} u^\nu u^\lambda = 0$$

particle trajectory $x^\mu(\tau)$, choosing $\tau = s$, we define $u^\mu = dx^\mu/ds$

$$S = -m\int ds = -m\int \sqrt{g_{\mu\nu}\frac{dx^\mu}{d\tau}\frac{dx^\nu}{d\tau}}\,d\tau$$

$$\gamma_{ij} \approx \delta_{ij}$$

# Photons in the expanding Universe

$$S = -\frac{1}{4}\int d^4x\sqrt{-g}g^{\mu\nu}g^{\lambda\rho}F_{\mu\lambda}F_{\nu\rho}$$

$dt = ad\eta$ — conformally flat metric

$$ds^2 = dt^2 - a^2(t)\delta_{ij}dx^i dx^j \longrightarrow ds^2 = a^2(\eta)[d\eta^2 - \delta_{ij}dx^i dx^j]$$

$$S = -\frac{1}{4}\int d^4x\,\eta^{\mu\nu}\eta^{\lambda\rho}F_{\mu\lambda}F_{\nu\rho}\,, \qquad A_\mu^{(\alpha)} = e_\mu^{(\alpha)}e^{ik\eta - i\mathbf{kx}}\,, \quad k = |\mathbf{k}|$$

$$\Delta x = 2\pi/k\,, \quad \Delta\eta = 2\pi/k$$

$$\lambda(t) = a(t)\Delta x = 2\pi\frac{a(t)}{k}\,, \quad T = a(t)\Delta\eta = 2\pi\frac{a(t)}{k}$$

# Redshift and the Hubble law $\lambda_0 = \lambda_i \frac{a_0}{a(t_i)} \equiv \lambda_i(1+z(t_i))$

$$\mathbf{p}(t) = \frac{\mathbf{k}}{a(t)}\,, \quad \omega(t) = \frac{k}{a(t)}$$

**for not very distant objects** $1\,\text{pc} \approx 3\,\text{ly}$

$$a(t_i) = a_0 - \dot{a}(t_0)(t_0 - t_i) \;\longrightarrow\; a(t_i) = a_0[1 - H_0(t_0 - t_i)]$$

$$z(t_i) = H_0(t_0 - t_i) = H_0 r\,, \quad z \ll 1$$

$$H_0 = h \cdot 100\,\frac{\text{km}}{\text{s}\cdot\text{Mpc}}\,, \quad h = 0.675 \pm 0.005$$

**similar reddening for other relativistic particles (small $H$, $\dot{H}$, etc.)**

$\mathbf{p} = \frac{\mathbf{k}}{a(t)}$ is true for massive particles as well

# Gas of free particles in the expanding Universe

homogeneous gas $dN = f(\mathbf{p},t)d^3\mathbf{X}d^3\mathbf{p} = \text{const}$

in comoving coordinates:

$$dN = f(\mathbf{k},\eta)d^3\mathbf{x}d^3\mathbf{k} = \text{const}\,, \quad d^3\mathbf{x} = \text{const}\,, \quad d^3\mathbf{k} = \text{const}$$

Hence $f(\mathbf{k},\eta) = f(\mathbf{k}) = \text{const}$

comoving volume equals physical volume

$$d^3\mathbf{x}d^3\mathbf{k} = d^3(a\mathbf{x})d^3\left(\frac{\mathbf{k}}{a}\right) = d^3\mathbf{X}d^3\mathbf{p}$$

$$f(\mathbf{p},t) = f(\mathbf{k}) = f(a(t)\cdot\mathbf{p})\,.$$

$$t = t_i\ :\quad f_i(\mathbf{p}) = f_i\left(\frac{\mathbf{k}}{a(t_i)}\right) \longrightarrow f(\mathbf{p},t) = f_i\left(\frac{\mathbf{k}}{a(t_i)}\right) = f_i\left(\frac{\mathbf{k}}{a(t_i)}\frac{a(t)}{a(t)}\right)$$

## Massless bosons (photons) fermions

$$f_i(\mathbf{p}) = f_{\rm Pl}\left(\frac{|\mathbf{p}|}{T_i}\right) = \frac{1}{(2\pi)^3}\frac{1}{e^{|\mathbf{p}|/T_i}-1}$$

$$f(\mathbf{p},t) = f_{\rm Pl}\left(\frac{a(t)|\mathbf{p}|}{a_i T_i}\right) = f_{\rm Pl}\left(\frac{|\mathbf{p}|}{T_{eff}(t)}\right)$$

$$T_{eff}(t) = \frac{a_i}{a(t)}T_i$$

decoupling at $T \gg m$ : neutrinos, hot(warm) dark matter

decoupling at $T \ll m$: thermally produced cold dark matter (e.g. WIMPs)

$$f(\mathbf{p}) = \frac{1}{(2\pi)^3}\exp\left(-\frac{m-\mu_i}{T_i}\right)\exp\left(-\frac{a^2(t)\mathbf{p}^2}{2ma_i^2 T_i}\right)$$

 calculate $f_0(p)$ for neutrinos

$$f(\mathbf{p},t) = \frac{1}{(2\pi)^3}\exp\left(-\frac{m-\mu_{eff}}{T_{eff}}\right)\exp\left(-\frac{\mathbf{p}^2}{2mT_{eff}}\right)$$

$$T_{eff}(t) = \left(\frac{a_i}{a(t)}\right)^2 T_i\,, \qquad \frac{m-\mu_{eff}(t)}{T_{eff}} = \frac{m-\mu_i}{T_i}$$

# Convention of GR description

**metric** inv: $\int d^4x\sqrt{-g}$

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu\,, \qquad \mu,\nu = 0,1,2,3 \qquad \text{signature } (+,-,-,-)$$

$$x^\mu \to x'^\mu(x^\mu)\,: \qquad B'^{\mu}_{\nu\lambda}(x') = \frac{\partial x'^\mu}{\partial x^\sigma}\frac{\partial x^\tau}{\partial x'^\nu}\frac{\partial x^\rho}{\partial x'^\lambda}B^\sigma_{\tau\rho}(x)$$

**Christoffel symbols:** $\nabla_\nu A^\mu(x) = \partial_\nu A^\mu(x) + \Gamma^\mu_{\nu\lambda}(x)A^\lambda(x)$

$$\Gamma^\mu_{\nu\lambda} = \tfrac{1}{2}g^{\mu\rho}(\partial_\nu g_{\rho\lambda} + \partial_\lambda g_{\rho\nu} - \partial_\rho g_{\nu\lambda})$$

## Riemann tensor

$$\nabla_\mu\nabla_\nu A^\lambda - \nabla_\nu\nabla_\mu A^\lambda = A^\sigma R^\lambda{}_{\sigma\mu\nu}\,, \quad R^\mu{}_{\nu\lambda\rho} = \partial_\lambda\Gamma^\mu_{\nu\rho} - \partial_\rho\Gamma^\mu_{\nu\lambda} + \Gamma^\mu_{\sigma\lambda}\Gamma^\sigma_{\nu\rho} - \Gamma^\mu_{\sigma\rho}\Gamma^\sigma_{\nu\lambda}$$

## Ricci tensor and scalar

$$R_{\mu\nu} \equiv R^\lambda{}_{\mu\lambda\nu}\,, \qquad R \equiv g^{\mu\nu}R_{\mu\nu} = R^{\lambda\mu}{}_{\lambda\mu}$$

# Einstein equations

$$ds^2 = dt^2 - a^2(t)\gamma_{ij}dx^i dx^j \ ,$$

$$S_{EH} = -\frac{1}{16\pi G}\int d^4x\sqrt{-g}R \ : \quad R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G T_{\mu\nu}$$

$$R_{00} = -3\frac{\ddot{a}}{a} \ , \quad R_{0i} = 0 \ , \quad R_{ij} = (\ddot{a}a + 2\dot{a}^2 + 2\varkappa)\gamma_{ij} \ ,$$

$$R = g^{\mu\nu}R_{\mu\nu} = -6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2} + \frac{\varkappa}{a^2}\right)$$

$T_{\mu\nu}$: macroscopic description

$T_{\mu\nu} = (p+\rho)u_\mu u_\nu - g_{\mu\nu}p$

$\frac{1}{2}\int d^4x\sqrt{-g}T_{\mu\nu}\delta g^{\mu\nu}$

liquid with $\rho(t)$ and $p(t)$

in the comoving frame $u^0 = 1, \mathbf{u} = 0$

$$T_{\mu\nu} = diag(\rho, \ a^2(t)\gamma_{ij}p)$$

# Friedmann equation for the present Universe

$$H^2 \equiv \left(\frac{\dot a}{a}\right)^2 = \frac{8\pi}{3} G(\rho_{\rm M} + \rho_{rad} + \rho_\Lambda + \rho_{\rm curv})$$

$$\frac{8\pi}{3} G\rho_{\rm curv} = -\frac{\varkappa}{a^2}\,, \quad \rho_c \equiv \frac{3}{8\pi G} H_0^2$$

$$\rho_c = \rho_{{\rm M},0} + \rho_{rad,0} + \rho_{\Lambda,0} = \rho_c = 0.53 \cdot 10^{-5} \frac{\rm GeV}{\rm cm^3}\,,$$

$$\Omega_X \equiv \frac{\rho_{X,0}}{\rho_c}\,, \qquad \frac{a_0}{a} = 1 + z$$

$$\left(\frac{\dot a}{a}\right)^2 = H_0^2 \left[\Omega_{\rm M}\left(\frac{a_0}{a}\right)^3 + \Omega_{rad}\left(\frac{a_0}{a}\right)^4 + \Omega_\Lambda + \Omega_{curv}\left(\frac{a_0}{a}\right)^2\right]$$

Obs: Equality ($\Omega_M = \Omega_{rad}$, $z \approx 3000$), Acceleration ($\ddot a = 0$); Age (14 Gy)

# Homogeneous and isotropic 3d manifolds

$$dl^2 = d\rho^2 + r^2(\rho)(d\theta^2 + \sin^2\theta d\phi^2)$$

$$r(\rho) = \begin{cases} R\sin(\rho/R)\,, & \text{3-sphere} \\ \rho\,, & \text{3-plane} \\ R\sinh(\rho/R)\,, & \text{3-hyperboloid} \end{cases}$$

$\rho$ is a geodesic distance; $S = 4\pi r^2(\rho)$; $\Delta\theta = \frac{l}{r(\rho)}$

# Brightness–redshift dependence in the Universe

$$ds^2 = dt^2 - a^2(t)\left[d\chi^2 + \sinh^2\chi\left(d\theta^2 + \sin^2\theta d\phi^2\right)\right]$$

coordinate distance $\chi = \int_{t_i}^{t_0} \frac{dt}{a(t)}$ $\qquad z(t) = \frac{a_0}{a(t)} - 1$

$$\chi(z) = \int_0^z \frac{dz'}{a_0 H_0} \frac{1}{\sqrt{\Omega_M (z'+1)^3 + \Omega_\Lambda + \Omega_{curv}(z'+1)^2}}$$

$$a_0^2 H_0^2 \Omega_{curv} = 1\,, \quad \Omega_M + \Omega_\Lambda + \Omega_{curv} = 1$$

$$S(z) = 4\pi r^2(z)\,, \quad r(z) = a_0 \sinh\chi(z)$$

detector: $N_\gamma \propto S^{-1}$, $\omega = \omega_i/(1+z)$, $dt_0 = (1+z)dt_i$

brightness (energy flux measured by a detector) is

$$J = \frac{L}{(1+z)^2 S(z)} \equiv \frac{L}{4\pi r_{ph}^2}\,, \quad r_{ph} = (1+z)\cdot r(z)$$

# Astrophysical and cosmological data are in agreement




$$\left(\frac{\dot{a}}{a}\right)^2 = H^2(t) = \frac{8\pi}{3} G \rho^{\text{energy}}_{\text{density}}$$

$$\rho^{\text{energy}}_{\text{density}} = \rho_{\text{radiation}} + \rho^{\text{ordinary}}_{\text{matter}} + \rho^{\text{dark}}_{\text{matter}} + \rho_\Lambda$$

$$\rho_{\text{radiation}} \propto 1/a^4(t) \propto T^4(t)\,, \quad \rho_{\text{matter}} \propto 1/a^3(t)$$

$$\rho_\Lambda = \text{const}$$

$$\frac{3H_0^2}{8\pi G} = \rho^{\text{energy}}_{\text{density}}(t_0) \equiv \rho_c \approx 0.53 \times 10^{-5} \frac{\text{GeV}}{\text{cm}^3}$$

radiation: $\Omega_\gamma \equiv \frac{\rho_\gamma}{\rho_c} = 0.5 \times 10^{-4}$

Baryons (H, He): $\Omega_B \equiv \frac{\rho_B}{\rho_c} = 0.046$

Neutrino: $\Omega_\nu \equiv \frac{\sum \rho_{\nu_i}}{\rho_c} < 0.01$

Dark matter: $\Omega_{DM} \equiv \frac{\rho_{DM}}{\rho_c} = 0.28$

Dark energy: $\Omega_\Lambda \equiv \frac{\rho_\Lambda}{\rho_c} = 0.68$

# Initial singularity (Bang!)

$$\left(\frac{\dot a}{a}\right)^2 = \frac{8\pi}{3} G\rho\,, \qquad p = w\rho\,, \qquad \textit{repeat}$$



**dust:** $p = 0$ — singular at $t = t_s$

$$\rho = \frac{\text{const}}{a^3}\,, \quad a(t) = const\cdot(t-t_s)^{2/3}\,, \quad \rho(t) = \frac{\text{const}}{(t-t_s)^2}$$

$$t_s = 0\,, \quad H(t) = \frac{\dot a}{a}(t) = \frac{2}{3t}\,, \quad \rho = \frac{3}{8\pi G}H^2 = \frac{1}{6\pi G}\frac{1}{t^2}$$

**radiation:** $p = \frac{1}{3}\rho$ — singular at $t = t_s$

$$\rho = \frac{\text{const}}{a^4}\,, \quad a(t) = const\cdot(t-t_s)^{1/2}\,, \quad \rho(t) = \frac{\text{const}}{(t-t_s)^2}$$

$$t_s = 0\,, \quad H(t) = \frac{\dot a}{a}(t) = \frac{1}{2t}\,, \quad \rho = \frac{3}{8\pi G}H^2 = \frac{3}{32\pi G}\frac{1}{t^2}$$

# Entropy

$$T^{\nu}_{\mu} = (p+\rho)u_{\mu}u^{\nu} - \delta^{\nu}_{\mu}p$$

$$\nabla_{\mu}T^{\mu}_{0} = 0 \quad \longrightarrow \quad \dot{\rho} + 3\frac{\dot{a}}{a}(\rho + p) = 0$$

for equation of state

$$p = p(\rho)$$

of the primordial plasma we obtain check $s(\rho, p)$ 

$$-3d(\ln a) = \frac{d\rho}{p+\rho} = d(\ln s)$$

entropy is conserved in a comoving volume

$$sa^3 = \text{const}$$

For the visible part of the Universe: $S \sim s_{\gamma,0} \cdot l_H^3 \sim 10^{88}$

# Horizon problem $l_H(t)$

a distance covered by photon emitted at $t = 0$

size of the causally connected part,
i.e. the visible part of the Universe ("inside horison")

$$ds^2 = dt^2 - a^2(t)\,dx^2 = a^2(\eta)\left(d\eta^2 - dx^2\right) \qquad ds^2 = 0$$

$$l_H(t) = a(t)\int dx = a(t)\int d\eta = a(t)\int_0^t \frac{c\,dt'}{a(t')} \propto t \quad \propto 1/H(t)$$




$a(t) \propto t^\alpha$, $0 < \alpha < 1$, $L_{phys} \propto a$




$$l_{H_0}/l_{H,r}(t_0) \sim l_{H_0}/l_{H,r}(t_r)\,a(t_r)/a_0 \sim H_r/H_0\,a(t_r)/a_0 \sim \sqrt{1+z_r} \simeq 30$$


2.7 K
TODAY
14 by
accelerated expansion
4.1 K
matter domination
7.1 by
0.26 eV
recombination
370 ty
$e+p \to H+\gamma$
matter domination
0.76 eV
radiation domination
57 ty
80 keV
3 min
$^3H+^4He \to ^7Li+\gamma$
primordial nucleosynthesis
$^2H+^2H \to n+^3He$
1 MeV
1 s
$p+p \to ^2H+\gamma$
2.5 MeV
neutrino decoupling
0.1 s
200 MeV
QCD phase transition
10 μs
confinement↔free quarks
100 GeV
Electroweak phase transition
0.1 ns
baryogenesis
hot Universe
dark matter production
reheating
inflation

# Ultra-relativistic particles

radiation: $p = \frac{1}{3}\rho$ singular at $t = 0$

$$\rho = \frac{\text{const}}{a^4}\,, \quad a(t) \propto t^{1/2}\,, \quad \rho(t) = \frac{\text{const}}{t^2}$$

$$H(t) = \frac{\dot{a}}{a}(t) = \frac{1}{2t}\,, \quad \rho = \frac{3}{8\pi G}H^2 = \frac{3}{32\pi G}\frac{1}{t^2}$$

$$l_H(t) = a(t)\int_0^t \frac{dt'}{a(t')} = 2t = \frac{1}{H(t)}\,.$$

In case of thermal equilibrium $T = \text{const}/(a g_*^{1/3}(T)) \approx \text{const}/a$

$$\rho_b = \frac{\pi^2}{30} g_b T^4\,, \quad \rho_f = \frac{7}{8}\frac{\pi^2}{30} g_f T^4$$ find $\frac{7}{8}$ ✍

$$\rho = \frac{\pi^2}{30} g_* T^4\,, \quad g_* = \sum_b g_b + \frac{7}{8}\sum_f g_f = g_*(T)\,, \quad s = \frac{\pi^2}{45} g_* T^3$$

# Last scattering: $\gamma e \to \gamma e$

$$\sigma_T = \frac{8\pi}{3}\frac{\alpha^2}{m_e^2} \approx 0.67 \cdot 10^{-24}\ \text{cm}^2\,, \qquad \tau_\gamma = \frac{1}{\sigma_T \cdot n_e(T)}$$

**last scattering:** $\tau_\gamma(T_f) \simeq H^{-1}(T_f) \simeq t_f$

$$T_f = 0.26\ \text{eV}\,, \quad z = 1100\,, \quad t_f = 370\,000\ \text{yr}$$

**for general processes one should solve kinetic equations**

$$\frac{dn_{X_i}}{dt} + 3Hn_{X_i} = \int (production - destruction)$$

Boltzmann equation in a comoving volume: $\frac{d}{dt}\left(n a^3\right) = a^3 \int \ldots$

# Recombination: horizon

**matter domination:** $l_{H,r} = 2H_r^{-1}$

$$H_r^2 = \frac{8\pi}{3} G\rho_M(t_r) = \frac{8\pi}{3} G\rho_{M,0} \left(\frac{a_0}{a_r}\right)^3 = \frac{8\pi}{3} G\rho_c \Omega_{M,0}(1+z_r)^3 .$$

**at recombination:** $l_{H,r} = \frac{2}{H_0\sqrt{\Omega_M}} \frac{1}{(1+z_r)^{3/2}}$

today: $l_{H,r}(t_0) = \frac{2}{H_0\sqrt{\Omega_M}} \frac{1}{\sqrt{1+z_r}}$

$$\frac{l_{H_0}}{l_{H,r}(t_0)} \sim \sqrt{1+z_r} \simeq 30$$

# Recombination: angle

$$\chi_r = \int_{t_r}^{t_0} \frac{dt}{a(t)}\,, \quad \Delta\theta_r = \frac{l_{H,r}}{r_a(z_r)}$$

$$r_a(z_r) = (1+z_r)^{-1} \cdot a_0 \cdot \sinh\chi_r$$

$$\Delta\theta_r = \frac{1}{\sqrt{z_r+1}}\,, \quad \Omega_{curv} = \Omega_\Lambda = 0\,.$$

$$\Delta\theta_r = \frac{1}{\sqrt{z_r+1}} \frac{2\sqrt{\Omega_{curv}/\Omega_M}}{\sinh\left(2\sqrt{\Omega_{curv}/\Omega_M}\, I\right)}\,.$$

$$I = \int_0^1 \frac{dy}{\sqrt{1+\frac{\Omega_\Lambda}{\Omega_M} y^6}}$$







Acoustic oscillations in relativistic plasma:
What matters is the sound horizon:

$$l_{s,r} = l_{H,r} \cdot v_s \approx$$

# Recombination: angle

$$\chi_r = \int_{t_r}^{t_0} \frac{dt}{a(t)}\,, \quad \Delta\theta_r = \frac{l_{H,r}}{r_a(z_r)}$$

$$r_a(z_r) = (1+z_r)^{-1} \cdot a_0 \cdot \sinh\chi_r$$

$$\Delta\theta_r = \frac{1}{\sqrt{z_r+1}}\,, \quad \Omega_{curv} = \Omega_\Lambda = 0\,.$$

$$\Delta\theta_r = \frac{1}{\sqrt{z_r+1}} \frac{2\sqrt{\Omega_{curv}/\Omega_M}}{\sinh\left(2\sqrt{\Omega_{curv}/\Omega_M}\, I\right)}\,.$$

$$I = \int_0^1 \frac{dy}{\sqrt{1+\frac{\Omega_\Lambda}{\Omega_M} y^6}}$$







Acoustic oscillations in relativistic plasma:
What matters is the sound horizon:

$$l_{s,r} = l_{H,r} \cdot v_s \approx l_{H,r}/\sqrt{3}$$

Then $\Delta\theta_{r,s} =$

$$\frac{1}{\sqrt{3}} \frac{1}{\sqrt{1+z}} \times \frac{180^\circ}{\pi} \simeq 1^\circ$$

- Hubble crisis: $67.4 \pm 0.5$ (Planck) vs $73.0 \pm 1.0$ (SH0ES)
- $\sigma_8$ tension... 3-2-1 $\sigma$
- Various tensions between different sets of data...


Distance Ladder (Cepheids)
ΛCDM
KP
SH0ES
CHP
SH0ES
SH0ES
SH0ES
SH0ES
SH0ES
WMAP1
WMAP3
WMAP5
WMAP7
WMAP9
P13
P15
BAO
P18
$H_0$ [km s$^{-1}$ Mpc$^{-1}$]
85
80
75
70
65
2000
2005
2010
2015
2020
Year

2105.05208




2105.05208




1707.06547







$$110/0.7\,\mathrm{Mpc} \simeq l_{H,r}(t_0) \times \sqrt{v_s^2} \simeq l_{H_0}/\sqrt{3}/\sqrt{1+z_r}$$

# Baryon Acoustic Oscillations (BAO), Sakharov




peak at $r_s \approx 147\,\mathrm{Mpc}$

$$\xi_z(s) \propto \frac{\langle n_z(\vec{x})\, n_z(\vec{x}+\vec{z})\rangle_{\vec{x}}}{n_z\, n_z}$$



# Impact of BAO: Galaxies vs Ly-$\alpha$

1707.06547









# Results of DESI: DE evolution $p = \omega(z)\rho$

2503.14738




$H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi}{3}\rho$

$\dot{\rho} + 3H(\rho + p) = 0$

$\omega = -1$: $p = -\rho$
gives $\rho = \text{const}$

$\omega > -1$ gives $\rho \searrow$

$\omega < -1$ gives $\rho \nearrow$

$\omega = 0$ is an oscillator:
$\langle E_k \rangle = \langle E_p \rangle$
dark matter

$\delta p = c_s^2 \delta\rho$

$\frac{\frac{1}{2}\dot{\phi}^2 + V(\phi)}{\frac{1}{2}\dot{\phi}^2 - V(\phi)} < -1$

# Perturbations blow up: $c_s^2 \to \infty$, $v \to \infty$







$$p = w\rho\,, \quad w(a) = w_0 + w_a\left(1 - \frac{a}{a_0}\right), \quad c_s^2 = \frac{\delta p}{\delta \rho} = w - \frac{w'}{3H(1+w)}$$

2505.24732


2.7 K
TODAY
14 by
accelerated expansion
4.1 K
matter domination
7.1 by
0.26 eV
recombination
370 ty
$e+p \to H+\gamma$
matter domination
0.76 eV
radiation domination
57 ty
80 keV
3 min
$^3H+^4He \to ^7Li+\gamma$
primordial nucleosynthesis
$^2H+^2H \to n+^3He$
1 MeV
1 s
$p+p \to ^2H+\gamma$
2.5 MeV
neutrino decoupling
0.1 s
200 MeV
QCD phase transition
10 μs
confinement↔free quarks
100 GeV
Electroweak phase transition
0.1 ns
baryogenesis
hot Universe
dark matter production
reheating
inflation

# Neutrino freeze-out

$T > m_e$ $\qquad e^+e^- \leftrightarrow \nu\bar{\nu},\ e\nu \leftrightarrow e\nu$

$$\sigma_\nu \sim G_F^2 E^2$$

neutrino interaction rate

$$\tau_\nu = \frac{1}{\langle \sigma_\nu n v \rangle} \sim \frac{1}{G_F^2 T^5}$$

$$\tau_\nu(T) \sim H^{-1}(T) = \frac{M_{Pl}^*}{T^2}$$

$$T_{\nu,f} \sim \left( \frac{1}{G_F^2 M_{Pl}^*} \right)^{1/3} \sim 2 \div 3 \text{ MeV}$$

# Neutron decoupling $p + e \longleftrightarrow n + \nu_e$

typical energy scales

$$T \gtrsim \Delta m = 1.3 \text{ MeV}, \qquad T \gtrsim m_e = 0.5 \text{ MeV}$$

## neutron interaction rate

$$\tau_{n\leftrightarrow p} = \frac{1}{\Gamma_{n\leftrightarrow p}} = \frac{1}{C_n G_F^2 T^5}$$

## neutron decoupling

$\Gamma_{n\leftrightarrow p}(T) \sim H(T) = T^2/M_{Pl}^*$

$$T_n = \frac{1}{\left(C_n M_{Pl}^* G_F^2\right)^{1/3}} \approx 1.4 \text{ MeV}$$

$g_* = 2 + \frac{7}{8}\cdot 4 + \frac{7}{8}\cdot 2 \cdot N_\nu$ $\qquad$ $t = \frac{1}{2H(T_n)} = \frac{M_{Pl}^*}{2T_n^2} = 1.2 \text{ s}$ $\qquad$ $T_n \approx 0.8 \text{ MeV}$

# Neutron density after decoupling

$$n_n = g_n \left(\frac{m_n T}{2\pi}\right)^{3/2} e^{\frac{\mu_n - m_n}{T}} \qquad \mu_n + \mu_{\nu_e} = \mu_p + \mu_e$$

$$\frac{n_n}{n_p} = e^{-\frac{m_n - m_p}{T}} e^{\frac{\mu_n - \mu_p}{T}}$$

for relativistic $e^+$ and $e^-$ Why ?

$$n_{e^-} - n_{e^+} \sim \mu_e T^2 \longrightarrow \frac{\mu_e}{T} \sim \frac{n_{e^-} - n_{e^+}}{T^3} = \frac{n_p}{T^3} \sim \eta_B \sim 10^{-9}$$

✍ find coefficient ⇑

$$\frac{n_n}{n_p} = e^{-\frac{m_n - m_p}{T_n}} \equiv e^{-\frac{\Delta m}{T_n}} \approx \frac{1}{5} e^{-\frac{\mu_{\nu_e}}{T_n}}$$

# $p + n \longrightarrow \mathrm{D} + \gamma$ Saha equation

$$n_n = 2\left(\frac{m_p T}{2\pi}\right)^{3/2} e^{\frac{\mu_n - m_n}{T}}, \qquad n_p = 2\left(\frac{m_p T}{2\pi}\right)^{3/2} e^{\frac{\mu_p - m_p}{T}},$$

## Chemical equilibrium: $\mu_D = \mu_p + \mu_n$

✍ prove that: $$n_D = n_p n_n g_D 2^{-1/2}\left(\frac{2\pi}{m_p T}\right)^{\frac{3}{2}} e^{\frac{\Delta_D}{T}}$$

Then $n_D \sim n_n$ at $T \simeq 70\,\mathrm{keV}$

## Interesting quantity

$$n_\mathrm{B} = \eta_\mathrm{B} \cdot n_\gamma = 0.24\eta_\mathrm{B} T^3$$

# Helium abundance (NO chemical equilibrium)

Neutrons remain mostly in helium

$$n_{^4\mathrm{He}}(T_{NS}) = \frac{1}{2} n_n(T_{NS}) \,,$$

## neutron-to-proton ratio

$\tau_n \approx 886$ s

$$\frac{n_n(T_{NS})}{n_p(T_{NS})} \approx \frac{1}{5} \cdot \mathrm{e}^{-\frac{t_{NS}}{\tau_n}} \cdot \mathrm{e}^{-\frac{\mu_\nu}{T_n}} \approx \frac{1}{7} \,,$$

$$Y_p \equiv X_{^4\mathrm{He}} = \frac{m_{^4\mathrm{He}} \cdot n_{^4\mathrm{He}}(T_{NS})}{m_p\,(n_p(T_{NS}) + n_n(T_{NS}))} = \frac{2}{\frac{n_p(T_{NS})}{n_n(T_{NS})} + 1} \approx 25\% \,.$$

from observations of relic helium abundance:

$$\Delta N_{\nu,eff} \leq 0.3 \,, \qquad \left|\frac{\mu_\nu}{T_n}\right| \lesssim 0.01$$

# Main nuclear reactions

1. $p(n,\gamma)$D — deuterium production, BBN starts.
2. D$(p,\gamma)^3$He, D(D,$n$)$^3$He, D(D,$p$)T, $^3$He$(n,p)$T — intermediate stage.
3. T(D,$n$)$^4$He, $^3$He(D,$p$)$^4$He — production of $^4$He.
4. T$(\alpha,\gamma)^7$Li, $^3$He$(\alpha,\gamma)^7$Be, $^7$Be$(n,p)^7$Li — production of the heaviest baryonic relics.
5. $^7$Li$(p,\alpha)^4$He — $^7$Li burning.

One has to compare reaction rates to the expansion rate

$$H(T_{NS} = 70 \text{ keV}) = 4 \cdot 10^{-3} \text{ s}^{-1}$$

to obtain nonequilibrium concentrations

# Neutron burning $p + n \longrightarrow \mathrm{D} + \gamma$

## @ $T = T_{NS} = 65$ keV

$$(\sigma v)_{p(n,\gamma)D} \approx 6 \cdot 10^{-20} \, \frac{\mathrm{cm}^3}{\mathrm{s}} \, .$$

for the rate of neutron disappearance (it meets proton!)

$$\Gamma_{p(n,\gamma)D} = n_p \cdot (\sigma v)_{p(n,\gamma)D} = \eta_B \cdot 2 \frac{\zeta(3)}{\pi^2} T^3 \cdot (\sigma v)_{p(n,\gamma)D} = 0.31 \ \mathrm{s}^{-1}$$

for $\eta_B = 6.15 \cdot 10^{-10}$ and $T = T_{NS}$

## So, neutrons disappear very rapidly

$$\Gamma_{p(n,\gamma)D} \gg H(T_{NS}) = 4 \cdot 10^{-3} \ \mathrm{s}^{-1}$$

# Deuterium burning $\mathrm{D}(\mathrm{D},n)^3\mathrm{He}$, $D(D,\mathrm{p})T$

Coloumb barier: tunneling $T_9 \equiv T/(10^9\ \mathrm{K}) = T/(86\ \mathrm{keV})$

$$\langle \sigma v \rangle_{DD} = 3 \cdot 10^{-15} \frac{\mathrm{cm}^3}{\mathrm{s}} \cdot T_9^{-2/3} \cdot \mathrm{e}^{-4.26 \cdot T_9^{-1/3}} .$$

deuterium stops burning when $T = T_{NS}(T_9 = 0.75)$

$$\Gamma_{DD} = n_D(T) \cdot \langle \sigma v \rangle_{DD}(T) \sim H(T) .$$

Then relic deuterium abundance is estimated as

$$\frac{n_D}{n_p} = \frac{1}{0.75 \eta_{\mathrm{B}}} \cdot \frac{n_D}{n_\gamma(T_{NS})} = 0.3 \cdot 10^{-4}$$

for $\eta_{\mathrm{B}} = 6.15 \cdot 10^{-10}$




Lack of Lithium...

$$Y_p = 0.2581 \pm 0.025\,,$$

$$D/H\big|_p = (2.87 \pm 0.21) \times 10^{-5}$$

1103.1261




similar results from other recent studies including structure formation

1001.4440, 1001.5218, 1202.2889

$N_\nu < 4.2$ @ 95%CL

$N_\nu < 4.3$ with shorter neutron's life...


2.7 K
TODAY
14 by
accelerated expansion
4.1 K
matter domination
7.1 by
0.26 eV
recombination
370 ty
$e+p \to H+\gamma$
matter domination
0.76 eV
radiation domination
57 ty
80 keV
3 min
$^3H+^4He \to ^7Li+\gamma$
primordial nucleosynthesis
$^2H+^2H \to n+^3He$
1 MeV
1 s
$p+p \to ^2H+\gamma$
2.5 MeV
neutrino decoupling
0.1 s
200 MeV
QCD phase transition
10 μs
confinement↔free quarks
100 GeV
Electroweak phase transition
0.1 ns
baryogenesis
hot Universe
dark matter production
reheating
inflation

# BACKUP SLIDES

# Galactic dark halos: flat rotation curves

$$v(R) = \sqrt{G\frac{M(R)}{R}}$$

$$M(R) = 4\pi \int_0^R \rho(r) r^2 dr$$






**observations:** $v(R) \simeq$ const

visible matter:

internal regions $v(R) \propto \sqrt{R}$

external ("empty") regions $v(R) \propto 1/\sqrt{R}$

# Dark Matter in clusters

## $X$-rays from hot gas in clusters

$$\frac{dP}{dR} = -\mu n_e(R) m_p \frac{GM(R)}{R^2}\,, \quad M(R) = 4\pi \int_0^R \rho(r) r^2 dr\,, \quad P(R) = n_e(R) T_e(R)$$

## galaxies in clusters

## virial theorem

$$U + 2E_k = 0$$

$$3M\langle v_r^2 \rangle = G\frac{M^2}{R}$$





## Milky Way: Virgo infall

# Gravitational lensing in GR: $\alpha = 4GM/(c^2 b)$



## Einstein Cross






$$\vec{\eta} = \frac{D_s}{D_l}\vec{\xi} - D_{ls}\,\vec{\alpha}\left(\vec{\xi}\right)$$

common lens with specific refraction coefficient

$$\vec{\alpha}\left(\vec{\xi}\right) = \frac{4G}{c}\int \frac{\vec{\xi}-\vec{\xi}'}{\left|\xi-\vec{\xi}'\right|^2}\, d^2\xi' \int \rho\left(\vec{\xi}', z\right) dz$$

source: quasar $D_s = 2.4$ Gpc

lens: galaxy $D_l = 120$ Mpc

# Dark Matter in clusters

gravitational lensing $\rho_B \approx 0.25\rho_{DM}$





# Colliding clusters (Bullet clusters 1E0657-558)







gravitational lensing

Observations in X-rays

$M \simeq 10 \times m$

scale is 200 kpc

clusters are at 1.5 Gpc