Elliptic Flow in BM@N

M.Mamaev


Tracks
Track selection: Nhits > 4 chi2/NDF < 5
Bad tracks
Good tracks
Primary selection DCA < 3 cm
Secondary
Primary
In acceptance
FHCal Acceptance
Out of Acceptance
Charge; kinematics
PID
protons
other
T-
T+
Acceptance Corrections
u-vectors
Monte-Carlo
reweighting
Acceptance Corrections
FHCal Modules
Subevent division
F{1, 2, …}
Acceptance Corrections
F{1, 2, … } x F{1, 2, … }
F{1, 2, … } x T{+,-}
Resolution
u_n,F{1, 2, … }
V_n


p_T (GeV/c)
2.5
2
1.5
1
0.5
0
−2.5 −2 −1.5 −1 −0.5 0 0.5 1 1.5 2 2.5
y_{cm}
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0

v_2 of protons wrt Reaction Plane (PCA h4)





v_2 of protons wrt FHCal symmetry planes (PCA h4)







v_2 of protons with different components (PCA h4)







v_2 of protons wrt FHCal symmetry planes (PCA h4)







v_2 of protons wrt FHCal symmetry planes (PCA h4)








q > 0
h1_chi2
Entries 53
Mean
Std Dev
hExpPos
Entries 725854
Mean 1.423
Std Dev 1.116
6000
5000
4000
3000
2000
1000
0
0.5
1
1.5
2
2.5
3
3.5
4
4.5
5
χ²
χ² / NDF

v_2 of protons wrt Reaction Plane





v_2 of protons wrt FHCal symmetry planes







Correction comparison: v1

plain



recentered



twist+resc.



pca (2 harm)



Correction comparison: v2

plain



recentered



twist+resc.



pca (2 harm)



Correction comparison: v1

2 harm



4 harm



8 harm



16 harm



Correction comparison: v2

2 harm



4 harm



8 harm



16 harm



Correction comparison: v2

2 harm



4 harm



8 harm



16 harm




N_{hits}
12
10
8
6
4
2
0
0
1
2
3
4
5
6
7
8
9
10
p (GeV/c)
0.3
0.25
0.2
0.15
0.1
0.05
0
Momentum resolution vs p and Nhits (prim protons)

Momentum resolution vs momentum (primary protons in TOF acceptance)







Momentum resolution vs momentum (primary protons in TOF acceptance)

ProjectionX of biny=8 [y=7.0..8.0]



ProjectionX of biny=9 [y=8.0..9.0]



ProjectionX of biny=10 [y=9.0..10.0]



DCA distribution for protons in TOF





CDF for proton DCA (prim / all)



Preliminary cuts on tracks

- p/q: [1.0, 8.0] GeV/c
- R_{dca}<3 cm (95% probability of proton being primary)
- Track not in X_{fhcal}: [-30, 160] cm, Y_{fhcal}:[-60; 60] cm (to reject autocorrelations)

Result after applying track-cuts





Efficiency


ProjectionY of binx=63 [x=-0.02..0.02]
slice_py_of_h2_pT_y_2212_good
Entries 351303
Mean 0.2766
Std Dev 0.2862
p_T (GeV/c)


ProjectionY of binx=76 [x=0.50..0.54]
slice_py_of_h2_pT_y_2212_good
Entries 86498
Mean 1.031
Std Dev 0.727
p_T (GeV/c)


ProjectionY of binx=89 [x=1.02..1.06]
slice_py_of_h2_pT_y_2212_good
Entries 364785
Mean 0.8419
Std Dev 0.3644
p_T (GeV/c)

2D efficiency



R_1 correction factor for F{1,...,4} (Monte-Carlo)

$x_2X_1X_1$
$x_2Y_1Y_1$
$y_2X_1Y_1$
$y_2Y_1X_1$









v_2 wrt Reaction Plane (Monte-Carlo)







Momentum resolution vs pT in bin of y



v_2 wrt Reaction Plane (Monte-Carlo, true momenta)




Browser File Edit View Options Tools Help
Files
Draw Option:
root
PROOF Sessions
ROOT Files
bmn.qa.jam2.md2.2026.09.28.root
h1_px;1
h1_py;1
h1_pz;1
h1_pT;1
h1_eta;1
h1_dca;1
h1_dca_prim;1
h1_dca_2212_tof;1
h1_dca_2212_prim_tof;1
h2_sim_pT_y_2212;1
h2_sim_pT_y_prim_2212;1
h2_pT_y_2212;1
h2_pT_y_2212_good;1
h2_pT_y_2212_good_400;1
h2_pT_y_2212_good_700;1
h3_phi_pT_y_2212_good;1
p1_dp_p;1
p2_dp_p_nhits_2212_prim;1
p2_dp_p_chi2_2212_good;1
p2_dp_pT_y_2212_good;1
home
mikhail
AmneziaVPN_4.8.9.2_linux_x64
Desktop
Documents
Downloads
Flow
GZ2025
Music
Pictures
Public
QnTools
Templates
Videos
bevybonk-template-master
bmn_environment
bmn_identification
bmn_monte_carlo
bmn_run8
correlation
flow
qa
CnSn.h8.root
Filter: All Files (*.*)
Canvas_1
Editor 1
Command
Command (local):


File Edit View Options Tools Help
ProjectionYZ of binx=15 [x=0.9..1]
p_T (GeV/c)
ϕ (rad)


File Edit View Options Tools Help
ProjectionX of biny=10 [y=0.90..1.00]
Entries 267602
Mean −0.09428
Std Dev 1.786
Underflow 0
Overflow 0
Integral 2.676e+05
ϕ (rad)


Browser File Edit View Options Tools Help
Files
Draw Option:
root
PROOF Sessions
ROOT Files
bmn.qa.jam2.md2.2026.09.28.root
h1_px;1
h1_py;1
h1_pz;1
h1_pT;1
h1_eta;1
h1_dca;1
h1_dca_prim;1
h1_dca_2212_tof;1
h1_dca_2212_prim_tof;1
h2_sim_pT_y_2212;1
h2_sim_pT_y_prim_2212;1
h2_pT_y_2212;1
h2_pT_y_2212_good;1
h2_pT_y_2212_good_400;1
h2_pT_y_2212_good_700;1
h3_phi_pT_y_2212_good;1
p1_dp_p;1
p2_dp_p_nhits_2212_prim;1
p2_dp_p_chi2_2212_good;1
p2_dp_pT_y_2212_good;1
home
mikhail
AmneziaVPN_4.8.9.2_linux_x64
Desktop
Documents
Downloads
Flow
GZ2025
Music
Pictures
Public
QnTools
Templates
Videos
bevybonk-template-master
bmn_environment
bmn_identification
bmn_monte_carlo
bmn_run8
correlation
flow
qa
CnSn.h8.root
Filter: All Files (*.*)
Canvas_1
Editor 1
Command
Command (local):
A special TPaveText to draw histogram st
stats
560,243
x=2.22388, y=0.553151


File Edit View Options Tools Help
ProjectionYZ of binx=15 [x=0.9..1]
p_T (GeV/c)
ϕ (rad)


File Edit View Options Tools Help
ProjectionX of biny=4 [y=0.30..0.40]
Number of Entries
Entries 1530916
Mean 0.2258
Std Dev 1.355
Underflow 0
Overflow 0
Integral 1.531e+06
ϕ (rad)


Browser File Edit View Options Tools Help
Files
Draw Option:
root
PROOF Sessions
ROOT Files
bmn.qa.jam2.md2.2026.09.28.root
h1_px;1
h1_py;1
h1_pz;1
h1_pT;1
h1_eta;1
h1_dca;1
h1_dca_prim;1
h1_dca_2212_tof;1
h1_dca_2212_prim_tof;1
h2_sim_pT_y_2212;1
h2_sim_pT_y_prim_2212;1
h2_pT_y_2212;1
h2_pT_y_2212_good;1
h2_pT_y_2212_good_400;1
h2_pT_y_2212_good_700;1
h3_phi_pT_y_2212_good;1
p1_dp_p;1
p2_dp_p_nhits_2212_prim;1
p2_dp_p_chi2_2212_good;1
p2_dp_pT_y_2212_good;1
home
mikhail
AmneziaVPN_4.8.9.2_linux_x64
Desktop
Documents
Downloads
Flow
GZ2025
Music
Pictures
Public
QnTools
Templates
Videos
bevybonk-template-master
bmn_environment
bmn_identification
bmn_monte_carlo
bmn_run8
correlation
flow
qa
CnSn.h8.root
Filter: All Files (*.*)
Canvas_1
Editor 1
Command
Command (local):
Canvas_1_n3
594,522
x=2.41876, y=-0.0952906


File Edit View Options Tools Help
ProjectionYZ of binx=6 [x=0.0..0]
p_T (GeV/c)
ϕ (rad)


File Edit View Options Tools Help
ProjectionX of biny=4 [y=0.30..0.40]
Entries 280505
Mean 0.08719
Std Dev 2.423
Underflow 0
Overflow 0
Integral 2.805e+05
ϕ (rad)


Browser File Edit View Options Tools Help
Files
Draw Option:
root
PROOF Sessions
ROOT Files
bmn.qa.jam2.md2.2026.09.28.root
h1_px;1
h1_py;1
h1_pz;1
h1_pT;1
h1_eta;1
h1_dca;1
h1_dca_prim;1
h1_dca_2212_tof;1
h1_dca_2212_prim_tof;1
h2_sim_pT_y_2212;1
h2_sim_pT_y_prim_2212;1
h2_pT_y_2212;1
h2_pT_y_2212_good;1
h2_pT_y_2212_good_400;1
h2_pT_y_2212_good_700;1
h3_centrality_sim_pT_y_prim_2212;
h3_phi_sim_pT_y_prim_2212;1
h3_phi_pT_y_2212_good;1
h3_centrality_pT_y_2212_good;1
p1_dp_p;1
p2_dp_p_nhits_2212_prim;1
p2_dp_p_chi2_2212_good;1
p2_dp_pT_y_2212_good;1
home
mikhail
AmneziaVPN_4.8.9.2_linux_x64
Desktop
Documents
Downloads
Flow
GZ2025
Music
Pictures
Public
QnTools
Templates
Videos
bevybonk-template-master
bmn_environment
bmn_identification
bmn_monte_carlo
bmn_run8
correlation
flow
qa
CnSn.h4.root
CnSn.h8.root
bmn.jam2.2026.07.10.x
bmn.qa.jam2.md2.2026
bmn.qa.jam2.md2.2026
Filter: All Files (*.*)
Canvas_1
Editor 1
Command
Command (local):
Pad graphics frame
TFrame
x=-0.985844, y=-8.47622


File Edit View Options Tools Help
ProjectionYZ of binx=11 [x=0.0..0]
h3_phi_pT_y_2212_good_yz
Entries 1213185
Mean x −0.01953
Mean y 0.3743
Std Dev x 2.305
Std Dev y 0.2791
p_T (GeV/c)
ϕ (rad)


File Edit View Options Tools Help
ProjectionX of biny=24 [y=1.15..1.20]
Entries 3307
Mean −2.839
Std Dev 0.8401
ϕ (rad)


Browser File Edit View Options Tools Help
Files
Draw Option:
root
PROOF Sessions
ROOT Files
bmn.qa.jam2.md2.2026.09.29.root
h1_px;1
h1_py;1
h1_pz;1
h1_pT;1
h1_eta;1
h1_dca;1
h1_dca_prim;1
h1_dca_2212_tof;1
h1_dca_2212_prim_tof;1
h2_sim_pT_y_2212;1
h2_sim_pT_y_prim_2212;1
h2_sim_px_pz;1
h2_sim_py_pz;1
h2_sim_px_py;1
h2_sim_px_pz_corr;1
h2_sim_py_pz_corr;1
h2_sim_px_py_corr;1
h2_pT_y_2212;1
h2_pT_y_2212_good;1
h2_pT_y_2212_good_400;1
h2_pT_y_2212_good_700;1
h2_px_pz_2212_good;1
h2_py_pz_2212_good;1
h2_px_pz_good;1
h2_py_pz_good;1
h2_px_py_good;1
h2_px_pz_corr_good;1
h2_py_pz_corr_good;1
h2_px_py_corr_good;1
h3_centrality_sim_pT_y_prim_2212;
h3_phi_sim_pT_y_prim_2212;1
h3_px_sim_pT_y_prim_2212;1
h3_py_sim_pT_y_prim_2212;1
h3_corr_phi_sim_pT_y_prim_2212;1
h3_phi_pT_y_2212_good;1
h3_corr_phi_pT_y_2212_good;1
h3_px_pT_y_2212_good;1
h3_py_pT_y_2212_good;1
h3_centrality_pT_y_2212_good;1
p1_dp_p;1
p1_dp_p_corr;1
p2_dp_p_nhits_2212_prim;1
p2_dp_p_chi2_2212_good;1
p2_dp_pT_y_2212_good;1
home
mikhail
AmneziaVPN_4.8.9.2_linux_x64
Desktop
Documents
Downloads
Filter: All Files (*.*)
Canvas_1
Editor 1
Command
Command (local):
ProjectionYZ of binx=11 [x=0.0..0]
h3_phi_sim
424,761
(x=5.86047, y=0.272349, binx=33, biny=6, binz=0 bine=0)
File Edit View Options Tools Help
ProjectionYZ of binx=11 [x=0.0..0]
Entries 1.335803e+07
Mean x 0.0005335
Mean y 0.7256
Std Dev x 1.79
Std Dev y 0.389
p_T (GeV/c)
ϕ (rad)
File Edit View Options Tools Help
ProjectionX of biny=11 [y=0.50..0.55]
Entries 689652
Mean −0.0002427
Std Dev 1.798
Before rotation
ϕ (rad)
1 2 3 4 5
Canvas_1_n3
us
jinr
04%
44°C
9.4G
70%
58%
09/30 14:00


Browser File Edit View Options Tools Help
Files
Draw Option:
root
PROOF Sessions
ROOT Files
bmn.qa.jam2.md2.2026.09.29.root
h1_px;1
h1_py;1
h1_pz;1
h1_pT;1
h1_eta;1
h1_dca;1
h1_dca_prim;1
h1_dca_2212_tof;1
h1_dca_2212_prim_tof;1
h2_sim_pT_y_2212;1
h2_sim_pT_y_prim_2212;1
h2_sim_px_pz;1
h2_sim_py_pz;1
h2_sim_px_py;1
h2_sim_px_pz_corr;1
h2_sim_py_pz_corr;1
h2_sim_px_py_corr;1
h2_pT_y_2212;1
h2_pT_y_2212_good;1
h2_pT_y_2212_good_400;1
h2_pT_y_2212_good_700;1
h2_px_pz_2212_good;1
h2_py_pz_2212_good;1
h2_px_pz_good;1
h2_py_pz_good;1
h2_px_py_good;1
h2_px_pz_corr_good;1
h2_py_pz_corr_good;1
h2_px_py_corr_good;1
h3_centrality_sim_pT_y_prim_2212;
h3_phi_sim_pT_y_prim_2212;1
h3_px_sim_pT_y_prim_2212;1
h3_py_sim_pT_y_prim_2212;1
h3_corr_phi_sim_pT_y_prim_2212;
h3_phi_pT_y_2212_good;1
h3_corr_phi_pT_y_2212_good;1
h3_px_pT_y_2212_good;1
h3_py_pT_y_2212_good;1
h3_centrality_pT_y_2212_good;1
p1_dp_p;1
p1_dp_p_corr;1
p2_dp_p_nhits_2212_prim;1
p2_dp_p_chi2_2212_good;1
p2_dpT_y_2212_good;1
home
mikhail
AmneziaVPN_4.8.9.2_linux_x64
Desktop
Documents
Downloads
Filter: All Files (*.*)
Canvas_1
Editor 1
Command
Command (local):
ProjectionYZ of binx=11 [x=0.0..0]
h3_corr_pT
313,753
(x=3.27907, y=0.293139, binx=33, biny=6, binc=0 bine=0)
File Edit View Options Tools Help
ProjectionYZ of binx=11 [x=0.0..0]
Entries 1.335851e+07
Mean x 0.0007073
Mean y 0.7253
Std Dev x 1.816
Std Dev y 0.3889
p_T (GeV/c)
ϕ (rad)
File Edit View Options Tools Help
ProjectionX of biny=12 [y=0.55..0.60]
Entries 691618
Mean −0.003152
Std Dev 1.816
After rotation
ϕ (rad)

original



Cov from true momentum



Cov from rec momentum



original



Cov from true momentum



Cov from rec momentum



R_1 correction factor for F{1,...,4}

$x_2X_1X_1$
$x_2Y_1Y_1$
$y_2X_1Y_1$
$y_2Y_1X_1$









Momentum resolution vs nhits in CT



Momentum resolution vs chi2/NDF (nhits =4, 5, 6)







Momentum resolution vs chi2/NDF (nhits > 5)



DCA-R comparison between prim and non-prim





Covariances for x_2 as a function of y_{cm} (n=1, 2)







Covariances for x_2 as a function of y_{cm} (n=2, 4)







Covariances for y_2 as a function of y_{cm} (n=1, 2)







Covariances for y_2 as a function of y_{cm} (n=2, 4)







v_2 wrt different planes (PCA with 1-4 harmonics)





$$v_2 = \frac{4\langle x_2^i X_1^j X_1^k \rangle}{R_1^j R_1^k}$$

v_2 with different components (PCA with 1-4 harmonics)











$$v_2 = \frac{4\langle x_2^i X_1^j X_1^k \rangle}{R_1^j R_1^k}$$


Tracks
Track selection: nhits chi2/NDF
Bad tracks
Good tracks
Primary selection
Secondary
Primary
FHCal Acceptance
In acceptance
Out of Acceptance
Charge; kinematics
T-
T+
PID
protons
other
u-vectors
Monte-Carlo
reweiting
Acceptance Corrections
Acceptance Corrections
FHCal Modules
Subevent division
F{1, 2, …}
F{1, 2, … } x F{1, 2, … }
F{1, 2, … } x T{+,-}
Resolution
u_n,F{1, 2, … }
V_n

v_2 comparison with STAR: pT (PCA 1-4)







v_2 comparison with STAR: ycm (PCA 1-4)








Xe+CsI 3.8A GeV Plain
0-10%; -0.1<y_cm<0.5
v2
pT (GeV/c)


Xe+CsI 3.8A GeV Plain
10-40%; -0.1<y_cm<0.5
BM@N Xe+CsI √sNN=3.26 GeV
STAR Au+Au √sNN=3.2 GeV
STAR Au+Au √sNN=3.5 GeV
v2
pT (GeV/c)


Xe+CsI 3.8A GeV Plain
40-60%; -0.1<y_cm<0.5
v2
pT (GeV/c)


Xe+CsI 3.8A GeV Plain
0-10%; 0.4<p_T<2 (GeV/c)
v_2
y_cm


Xe+CsI 3.8A GeV Plain
10-40%; 0.4<p_T<2 (GeV/c)
BM@N Xe+CsI √s_NN=3.26 GeV
STAR Au+Au √s_NN=3.2 GeV
STAR Au+Au √s_NN=3.5 GeV
v_2
y_cm


Xe+CsI 3.8A GeV Plain
40-60%; 0.4<p_T<2 (GeV/c)
v_2
y_cm

Proton v_2: pT, y, centrality and run-Id (de-bugged FHCal numeration)









OLD: Proton v_2



UPD: Proton v_2







UPD: Proton v_2 after track reweighting







OLD: Proton v_2







UPD: Proton v_2







UPD: Proton v_2







UPD: Proton v_2







Analysis setup

- JAM MD2 + coalescence for Xe+Cs at E_{kin}=3.8A GeV as an input
- Full reconstruction
- Event selection:
  - N tracks for VTX > 2
- Track selection (as in data)
  - Nhits > 5
  - Chi2/NDF for CT-track < 5
  - R_{dca}<5
- Identification via associated MC-track + hit in any of the TOF

Issue in question:

Previously even v_2 was in agreement, check the article in Particles





Issue fixed: Tw+Resc Qn-Tools like: v1





Issue fixed: Tw+Resc Qn-Tools like: v2





Basic flow measurements scheme



Recentering, twist and rescaling corrections

Twist and rescaling:

Given u_n= [cos nφ, sin nφ]

$$C = \begin{bmatrix} 1 + c_{2n} & s_{2n} \\ s_{2n} & 1 - c_{2n} \end{bmatrix}$$ mixing matrix for double harm.

$$u'_n = C^{-1}(u_n - r_n)$$ Recentering, twist and rescaling

$$r = \begin{bmatrix} c_n \\ s_n \end{bmatrix}$$ recentering vector

In QnTools there are no u_n-vectors, for all vectors sum is calculated according to:

$$\vec{q}_n = \frac{\sum_k w_k \vec{u}_n}{\sum_k w_k}$$ In each event

$$\begin{bmatrix} c_n \\ s_n \end{bmatrix} = \frac{1}{M} \sum_{m=1}^{M} \vec{q}_n^{\,m}$$ Sum over events

Then recentering, twist and rescaling is applied event-by-event to each q_n. But:

$$\begin{bmatrix} c_n \\ s_n \end{bmatrix} = \frac{\sum_{m=1}^{M} \sum_k w_k \vec{u}_n}{\sum_{m=1}^{M} \sum_k w_k}$$ ←this value bias flow,

not this→ $$\begin{bmatrix} c_n \\ s_n \end{bmatrix} = \frac{1}{M} \sum_{m=1}^{M} \frac{\sum_k w_k \vec{u}_n}{\sum_k w_k}$$

The subtle difference in calculating corrections

$$\begin{bmatrix} c_n \\ s_n \end{bmatrix} = \frac{\sum_{m=1}^{M} \sum_k w_k \vec{u}_n}{\sum_{m=1}^{M} \sum_k w_k}$$ ←this value bias flow,

not this→ $$\begin{bmatrix} c_n \\ s_n \end{bmatrix} = \frac{1}{M} \sum_{m=1}^{M} \frac{\sum_k w_k \vec{u}_n}{\sum_k w_k}$$

- Averaging over all particles in all events — characteristic of a detector
- Correcting particle distribution: moves the fourier moments to zero
- Work for all multiplicities

- Averaging in a single event and then averaging over all events
- Correcting event-by-event distribution of q_n
- Probably should work the same for large number of particles in one event (>1000)

After correction x and y components should give the same result (by design)

$$r = \begin{bmatrix} c_n \\ s_n \end{bmatrix}$$ recentering vector

$$C = \begin{bmatrix} 1 + c_{2n} & s_{2n} \\ s_{2n} & 1 - c_{2n} \end{bmatrix}$$ mixing matrix for double harm.

Removes the constant: shifts mean to zero

Untangles and then equalizes cos/sin

Example from the data: u_2 F F correlation







Still large contribution from uncorrected
$\langle x_2 \rangle \langle Y_1 Y_1 \rangle$ and $\langle y_2 \rangle \langle Y_1 Y_1 \rangle$

Correction comparison: v1

QnTools-like Rec+Tw+Resc



Corrected rec+tw+resc



Correction comparison: v2

QnTools-like Rec+Tw+Resc



Corrected rec+tw+resc



Basic flow measurements scheme



Efficiencies used



3D efficiency: p, η, φ



Efficiency comparison, QnTools Tw+Resc: v1

2D efficiency: p, η



3D efficiency: p, η, φ



Efficiency comparison, QnTools Tw+Resc: v2

2D efficiency: p, η



3D efficiency: p, η, φ



Efficiency comparison, corrected Tw+Resc: v1

2D efficiency: p, η



3D efficiency: p, η, φ



Efficiency comparison, corrected Tw+Resc: v2

2D efficiency: p, η



3D efficiency: p, η, φ



Efficiency comparison, corrected PCA: v1

2D efficiency: p, η



3D efficiency: p, η, φ



Efficiency comparison, corrected PCA: v2

2D efficiency: p, η



3D efficiency: p, η, φ



Directed by

ROBERT B. WEIDE

Correction comparison: v1

Rec+Tw+Resc



PCA



Correction comparison: v2

Rec+Tw+Resc



PCA



Efficiency comparison: Tw+Resc + v1

Corrected rec+tw+resc





Upd: v1 vs pT and y for MC simulation (η<3.0)





Old: v1 vs pT and y for MC simulation (tr not pointing to FHCAl)





PCA 5 harmonics: v1





PCA 2 harmonics: v1





PCA 5 harmonics: v2





PCA 2 harmonics: v2





R1 vs centrality for MC simulation (Tw+Resc)







Twist+Rescaling performance on v1





Twist+Rescaling performance on v2





v1 for protons without corrections on u





v2 for protons without corrections on u





The role of detector acceptance in the flow measurements

$$v_n = \frac{1}{2\pi} \int d\varphi d\Psi A(\varphi) \left(1 + 2\sum_{n=1}^{\infty} v_n(\cos n\varphi \cos n\Psi + \sin n\varphi \sin n\Psi)\right) \cos n\varphi \cos n\Psi$$

detector acceptance/ efficiency

distribution due to physics

In the ideal case scenario A(φ) = 1 / 2π, probability to measure any φ is the same

In the real world probability to measure different φ varies with φ:

$$A(\varphi) = \frac{1}{2\pi}\left[1 + 2\sum_{m=1}^{\infty}(c_m \cos m\varphi + s_m \sin m\varphi)\right]$$

Interaction of the sine and cosine with acceptance

$$x_n = A(\varphi)\cos n\varphi = \frac{1}{2\pi}\left[1 + 2\sum_{m=1}^{\infty}(c_m \cos m\varphi + s_m \sin m\varphi)\right]\cos n\varphi =$$

$$\frac{1}{2\pi}\left[c_n + (1 + c_{2n})\cos n\varphi + s_{2n}\sin n\varphi + \sum_{k=1;k\neq n}^{\infty}((c_{k+n} + c_{k-n})\cos k\varphi + (s_{k+n} + s_{k-n})\sin k\varphi)\right]$$

$$y_n = A(\varphi)\sin n\varphi = \frac{1}{2\pi}\left[1 + 2\sum_{m=1}^{\infty}(c_m \cos m\varphi + s_m \sin m\varphi)\right]\sin n\varphi =$$

$$\frac{1}{2\pi}\left[s_n + s_{2n}\cos n\varphi + (1 - c_{2n})\sin n\varphi + \sum_{k=1;k\neq n}^{\infty}((s_{k+n} - s_{k-n})\cos k\varphi + (c_{k-n} - c_{k+n})\sin k\varphi)\right]$$

The same goes with Q_n-vectors. In attempt to measure v_n we get:
$v_n = p_0 + p_1 v_1 + p_2 v_2 + \ldots + p_n v_n + \ldots$

Rescentering, Twist and Rescaling procedure



One of the ways to calculate the corrections is via the covariance matrix:

$$\vec{u}_n = (x_n, y_n) = (\cos n\varphi, \sin n\varphi)$$

$$\Sigma = \begin{bmatrix} cov(x_n, x_n) & cov(y_n, x_n) \\ cov(x_n, y_n) & cov(y_n, y_n) \end{bmatrix} = \begin{bmatrix} \vec{w}_1 \\ \vec{w}_2 \end{bmatrix}^T \times \begin{bmatrix} \sigma_1 & 0 \\ 0 & \sigma_2 \end{bmatrix} \times \begin{bmatrix} \vec{w}_1 \\ \vec{w}_2 \end{bmatrix}$$

Limitations of the twist+rescaling

$$\vec{u}_n = (x_n, y_n) = (\cos n\varphi, \sin n\varphi)$$

$$\Sigma = \begin{bmatrix} cov(x_n, x_n) & cov(y_n, x_n) \\ cov(x_n, y_n) & cov(y_n, y_n) \end{bmatrix} = \begin{bmatrix} \vec{w}_1 \\ \vec{w}_2 \end{bmatrix}^T \times \begin{bmatrix} \sigma_1 & 0 \\ 0 & \sigma_2 \end{bmatrix} \times \begin{bmatrix} \vec{w}_1 \\ \vec{w}_2 \end{bmatrix}$$

Before the correction:
$v_n = p_0 + p_1 v_1 + p_2 v_2 + \ldots + p_n v_n + \ldots$

With the correction we might untangle sine and cosine of the same harmonic:
$v_n = 0 + p_1 v_1 + p_2 v_2 + \ldots + 1 \times v_n + \ldots$

This works if the v_n >> rest (for v_1 for example)
But for v_2 the rest sum might be still larger

Proposed correction

Why stop on the harmonic n, when we can build the covariance matrix for arbitrary number of harmonics:

$$\Sigma = \begin{bmatrix} cov(x_1,x_1) & cov(y_1,x_1) & cov(x_2,x_1) & \dots & cov(y_n,x_1) \\ cov(x_1,y_1) & cov(y_1,y_1) & cov(x_2,y_1) & \dots & cov(y_n,y_1) \\ \dots & & & & \\ cov(x_1,y_n) & cov(y_1,y_n) & cov(x_2,y_n) & \dots & cov(y_n,y_n) \end{bmatrix} = \begin{bmatrix} \vec{w}_1 \\ \dots \\ \vec{w}_{2n} \end{bmatrix}^T \begin{bmatrix} \sigma_1 & 0 & \dots & 0 \\ 0 & \sigma_2 & \dots & 0 \\ \dots & & & \\ 0 & 0 & \dots & \sigma_{2n} \end{bmatrix} \begin{bmatrix} \vec{w}_1 \\ \dots \\ \vec{w}_{2n} \end{bmatrix}$$

Directions that are really measured

Amplitude of changes in this directions

Before the correction:

$v_n = p_0 + p_1 v_1 + p_2 v_2 + \dots + p_n v_n + \dots$

Now, applying the correction we get:

$v_n = v_n + \dots$

PCA performance on v1





PCA performance on v2





R1 vs centrality for MC simulation (PCA)







v1 vs pT and y for MC simulation (Tw+Resc)





v1 vs pT and y for MC simulation (PCA)





New correction performance on v1





New correction performance on v2





BM@N Monte-Carlo simulation

- JAM v2 with MD2 EOS + coalescence was used to generate Xe^{124}+Cs^{133} collisions at E_{kin}=3.8A GeV
- Approximately 3M of events were put through the realistic chain of simulation
- Same track selection criteria as in data
- Identification via the PDG and Mother-Id of associated model particle + hit in either of TOF400 or TOF700



R1 vs centrality for MC simulation









Comparison of v1 wrt RP for reconstructed and model





Comparison of v2 wrt RP for reconstructed and model





BM@N v_1 of p vs model calculations





BM@N v_2 of p vs model calculations



BM@N v_1, v_2 of p vs model calculations





BM@N v_1 of p vs JAM MD2 with vs no coalescence





BM@N v_1 of p vs JAM Skyrme (no coalescence)





BM@N v_1 of p vs UrQMD hard-skyrme with vs no coalescence





BM@N v_2 of p vs JAM MD2 with vs no coalescence





BM@N v_2 of p vs JAM Skyrme (no coalescence)





BM@N v_1 of p vs JAM MD2 calculations





BM@N v_2 of p vs JAM MD2 calculations




y_{cm}
0.7
0.6
0.5
0.4
0.3
0.2
0.1
0
6800
7000
7200
7400
7600
7800
8000
8200
run id


p_T (GeV/c)
0.5
0.4
0.3
0.2
0.1
0
6800
7000
7200
7400
7600
7800
8000
8200
run id


Phi
(rad)
0.35
0.3
0.25
0.2
0.15
0.1
0.05
0
6800
7000
7200
7400
7600
7800
8000
8200
run id

Relative run-by-run efficiency (target run 7500)

#7128, 2 < η < 3



#8188, 2 < η < 3



4D-efficiency was built: η, p_T, ϕ, run-id. $E = N_{curr}/N_{target\ run\text{-}id}$

Proton y-distribution before/after reweighting





Proton p_T-distribution before/after reweighting





Proton p_T-distribution before/after reweighting





Proton v_1 after track reweighting









Proton v_2 after track reweighting



Proton v_2 after track reweighting



Proton v_2 after track reweighting







Todo list:

- ❏ Corrections: my recentering, my twist and rescaling, my decomposition steps
- ❏ <Nhits> vs run-id
- ❏ <chi2/NDF> vs run-id
- ❏ <DCA> vs run-id
- ❏ Run-id dependency of vtx for central events
- ❏ $\langle F^i F^j \rangle$ as a function of centrality and run-id for midcentral events and for different run-id's
- ❏ Run-id QA for only physical data
- ❏ <beta> vs p/q
- ❏ Run-id dependence for each matrix element (visualize corrections)
- ❏ TOF vs T0 width
- ❏ TOF vs RPC width
- ❏ TOF QA strip-by-strip run-by-run
- ❏ Why 2 GeV/c cut for time-shifts
- ❏ p_T-y distributions for negatively charged particles and for identified as pions


N_{hits}
0
1
2
3
4
5
6
6800
7000
7200
7400
7600
7800
8000
8200
run id


χ^2/NDF
1.4
1.2
1
0.8
0.6
0.4
0.2
0
6800
7000
7200
7400
7600
7800
8000
8200
run id


N_p
0
1
2
3
4
5
6
7
6800
7000
7200
7400
7600
7800
8000
8200
run id

Charged particles multiplicity vs run-id



Normalized multiplicity



Squared mass of protons with TOF400 and TOF700





Vertex position







TOF-400



TOF-700



TOF-400



TOF-700



New FHCal-subevents layout

$$A(\varphi) = \frac{1}{2\pi}\left[1 + 2\sum_{m=1}^{\infty}(c_m \cos m\varphi + s_m \sin m\varphi)\right]$$

Acceptance function acts as a weight A(φ) u_n and the measured vector x is:

$$x = Mu + c$$

$$u = [x_1, y_1, x_2, y_2]^T$$

$$c = [c_1, s_1, c_2, s_2]^T$$

$$M = \begin{bmatrix} 1+c_2 & s_2 & c_1+c_3 & s_1+s_3 \\ s_2 & 1-c_2 & s_3-s_1 & c_1-c_3 \\ c_1+c_3 & s_3-s_1 & 1+c_4 & s_4 \\ s_1+s_3 & c_1-c_3 & s_4 & 1-c_4 \end{bmatrix}$$



$$u = M^{-1}(x - c)$$

Acceptance Fourier-moments for protons







Acceptance Fourier-moments for protons (corrected)







Acceptance Fourier-moments for protons (corrected & zoomed)







Acceptance Fourier-moments for FHCal









Acceptance Fourier-moments for FHCal



Result with only recentering applied









Result with only recentering applied: y-dep vs STAR







Result with only recentering applied: pT-dep vs STAR







Result with decomposition applied



Result with full decomposition: y-dep vs STAR







Result with full decomposition: pT-dep vs STAR







Additional check: comparison with v_1-preliminary









Proton N-sigma distributions

TOF-400



TOF-700



Proton p_T-y acceptance







Further notations

In the case of azimuthal isotropy of detector acceptance and no magnetic field:

$$\langle x_2 \rangle = 0 \quad \langle Y_1 \rangle = 0$$

$$v_2 = 2\frac{\langle x_2 Y_1^a Y_1^b \rangle}{\langle Y_1^a Y_1^b \rangle}$$

v_2 is the resolution-corrected cumulant of u_2 and two Q_1-vectors



$$cum(x_2, Y_1^a, Y_1^b) = \langle x_2 Y_1^a Y_1^b \rangle - \langle x_2 \rangle \langle Y_1^a Y_1^b \rangle - \langle x_2 Y_1^a \rangle \langle Y_1^b \rangle - \langle x_2 Y_1^b \rangle \langle Y_1^a \rangle + 2\langle x_2 \rangle \langle Y_1^a \rangle \langle Y_1^b \rangle$$

$$v_2 = 2\frac{cum(x_2, Y_1^a, Y_1^b)}{\langle Y_1^a Y_1^b \rangle}$$


20 cm
15 cm
35
36
37
38
39
40
41
42
43
44
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
45
46
47
48
49
50
51
52
53
54
100 cm
105 cm
2,5 cm
75 cm
155 cm

New FHCal-subevents layout

$$A(\varphi) = \frac{1}{2\pi}\left[1 + 2\sum_{m=1}^{\infty}(c_m \cos m\varphi + s_m \sin m\varphi)\right]$$

Acceptance function acts as a weight A(φ) u_n and the measured vector x is:

$$x = Mu + c$$

$$u = [x_1, y_1, x_2, y_2]^T$$

$$c = [c_1, s_1, c_2, s_2]^T$$

$$M = \begin{bmatrix} 1+c_2 & s_2 & c_1+c_3 & s_1+s_3 \\ s_2 & 1-c_2 & s_3-s_1 & c_1-c_3 \\ c_1+c_3 & s_3-s_1 & 1+c_4 & s_4 \\ s_1+s_3 & c_1-c_3 & s_4 & 1-c_4 \end{bmatrix}$$



$$u = M^{-1}(x - c)$$

Correlation of recentered Q_1-vectors from FHCal







Plain protons with recentered F1 and F2







Plain protons with recentered F1 and F3







Plain protons with recentered F2 and F3







Plain proton v_2 wrt different symmetry planes







Plain protons with plain F2 and F3







Plain Q1-vectors

Correlation of plain Q_1-vectors from FHCal







Plain protons with plain F1 and F2







Plain protons with plain F1 and F3







Plain protons with plain F2 and F3







Plain proton v_2 wrt different symmetry planes







Plain protons with plain F2 and F3







plain u_2-vector with rec.-corr. F1 and F2







plain u_2-vector with rec.-corr. F1 and F3







plain u_2-vector with rec.-corr. F2 and F3







v_2 as a function of y_{cm}, p_T and centrality







Proton v_2 vs deuteron v_2







Comparison with STAR-FXT: y-dependence







Comparison with STAR-FXT: y-dependence








Xe+CsI 3.8A GeV Plain
0-10%; 0.4<p_T<2 (GeV/c)
c_n
s_n
v_2
y_cm


Xe+CsI 3.8A GeV Plain
0-10%; 0.0<y_cm<0.6
1
2
3
4
v_2
p_T (GeV/c)


Xe+CsI 3.8A GeV Plain
0<p_T<2 (GeV/c); 0.0<y_cm<0.6
v_2
centrality (%)


Xe+CsI 3.8 A GeV
F1×F2
F1
F1×F2
centrality (%)


Xe+CsI 3.8 A GeV
F1×F3
⟨x1⟩
⟨y1⟩
⟨x2⟩
⟨y2⟩
F2
F1×F3
centrality (%)


Xe+CsI 3.8 A GeV
F2×F3
F3
F2×F3
centrality (%)


Xe+CsI 3.8A GeV Plain
0-10%; -0.1<y_cm<0.5
v2
p_T (GeV/c)


Xe+CsI 3.8A GeV Plain
10-40%; -0.1<y_cm<0.5
BM@N Xe+CsI √s_NN=3.26 GeV
STAR Au+Au √s_NN=3.2 GeV
STAR Au+Au √s_NN=3.5 GeV
v2
p_T (GeV/c)


Xe+CsI 3.8A GeV Plain
40-60%; -0.1<y_cm<0.5
v2
p_T (GeV/c)


Au+Au, JAM MF MD2
protons, 0.4 < p_T < 2.0 GeV/c, |y_cm| < 0.1
0-10%
20-30%
50-60%
V2
√s_NN, GeV


Au+Au, JAM MF MD2
protons, 0.4 < p_T < 2.0 GeV/c, |y_cm| < 0.4
0-10%
20-30%
50-60%
V2
√s_NN, GeV

Decomposition correction

Average of FHCal $Q_{1,2}$-vector components







Average of proton $u_{1,2}$-vector components







The core idea of correction

In the case of ideal detector acceptance, distribution is described as:

$$\rho(\varphi - \Psi) = \frac{1}{2\pi}(1 + 2\sum_{n} v_n \cos n(\varphi - \Psi))$$

In the case of azimuthal acceptance non-uniformity this transforms into:

$$P(\varphi - \Psi) = \frac{1}{2\pi}\left(1 + 2\sum_{m}(c_m \cos m\varphi + s_m \sin m\varphi)\right) \times \left(1 + 2\sum_{n} v_n \cos n(\varphi - \Psi)\right)$$

Q_n-vector components in the case of ideal detector acceptance:

$$X_n^{\rho} \equiv \langle x_n \rangle_{\rho}(\Psi) = \int d\varphi \frac{1}{2\pi}(1 + \sum_{n=1}^{\infty} v_n \cos n(\varphi - \Psi)) \cos n\varphi = V_n \cos n\Psi$$

$$Y_n^{\rho} \equiv \langle y_n \rangle_{\rho}(\Psi) = \int d\varphi \frac{1}{2\pi}(1 + \sum_{n=1}^{\infty} v_n \cos n(\varphi - \Psi)) \sin n\varphi = V_n \sin n\Psi$$

In the case of azimuthal non-uniformity:

$$X_n^{P} = \int d\varphi P(\varphi - \Psi) \cos n\varphi = X_n^{\rho} + \sum_{m=1}^{\infty}(c_m X_{n+m}^{\rho} + c_m X_{m-n}^{\rho} + s_m Y_{m+n}^{\rho} + s_m Y_{m-n}^{\rho})$$

$$Y_n^{P} = \int d\varphi P(\varphi - \Psi) \sin n\varphi = Y_n^{\rho} + \sum_{m=1}^{\infty}(c_m Y_{m+n}^{\rho} - c_m Y_{m-n}^{\rho} - s_m X_{m+n}^{\rho} + s_m X_{m-n}^{\rho})$$

Acceptance decomposition

$$\langle x_n \rangle = \int d\varphi \frac{1}{2\pi} \left(1 + 2\sum_m (c_m \cos m\varphi + s_m \sin m\varphi)\right) \cos n\varphi = c_n$$

$$\langle y_n \rangle = \int d\varphi \frac{1}{2\pi} \left(1 + 2\sum_m (c_m \cos m\varphi + s_m \sin m\varphi)\right) \sin n\varphi = s_n$$

$$cov(x_n x_l) = \langle x_n x_l \rangle - c_n c_l$$

First, for vector s = [X_1, Y_1, X_2, Y_2] we bild covariation matrix M, where $M_{i,j}$ = cov(s_i, s_j). Then we apply the whitening procedure:

x = $M^{-1/2}$s
After this procedure: cov(x_i, x_i) = 1, cov(x_i, x_j)$_{i!=j}$=0
This supposed to decompose acceptance so that:

$$x = [\cos\varphi, \sin\varphi, \cos 2\varphi, \sin 2\varphi]$$

Cov(x_1, x_n/y_n) for protons before/after whitening:







$Cov(y_1, x_n/y_n)$ for protons before/after whitening:







Cov(x_2, x_n/y_n) for protons before/after whitening:







Cov(y_2, x_n/y_n) for protons before/after whitening:







$Cov(x_1, x_n/y_n)$ for FHCal before/after whitening:







$Cov(y_1, x_n/y_n)$ for protons before/after whitening:







Cov(x_2, x_n/y_n) for protons before/after whitening:







Cov(y_2, x_n/y_n) for protons before/after whitening:







Xe+CsI at 3.8A GeV

Protons x Rec. F1 x Rec. F2







Protons x Rec. F1 x Rec. F3







Protons x Rec. F2 x Rec. F3







Corrected proton v_2







Corrected proton v_2







Xe+CsI at 3.0A GeV

Protons x Rec. F1 x Rec. F2







Protons x Rec. F1 x Rec. F3







Protons x Rec. F2 x Rec. F2







Protons x Rec. F2 x Rec. F2







Correlation with F1 and F2

Checking the corrections on u-vector, FHCal vectors are Rescaled-corrected

- After applying recentering on u-vector, correlation drops 10 times



Correlation with F2 and F3

Checking the corrections on u-vector, FHCal vectors are Rescaled-corrected

- After applying recentering on u-vector, correlation drops 10 times



Correlation with F1 and F3

Checking the corrections on u-vector, FHCal vectors are Rescaled-corrected

- After applying recentering on u-vector, correlation drops 10 times



Plain proton with single F1







Plain proton with single F2







Plain proton with single F3







Recentered proton with single F1







Recentered proton with single F2







Recentered proton with single F3







Plain double proton







Recentered double







Rescaled double proton







Correlations with recentered F-vectors

Plain proton v_2







Recentered proton v_2







Rescaled proton v_2







Correlations with rescaled F-vectors

Plain proton v_2







Recentered proton v_2







Rescaled proton v_2







Plain proton v_2







Recentered proton v_2







Rescaled proton v_2







Plain proton correlation with F1







Recentered proton correlation with F1







Rescaled proton correlation with F1







Plain proton correlation with F2







Recentered proton correlation with F2







Rescaled proton correlation with F2







Plain proton correlation with F3







Recentered proton correlation with F3







Rescaled proton correlation with F3







Calculating the effective rotation angle

α ~ 31.717474







Symmetry plane resolution in Xe+Cs(I) collisions







All the estimations for symmetry plane resolutions are in a good agreement