

# On moduli space of the Wigner quasiprobability distributions for N-dimensional quantum systems

Vahagn Abgaryan, Arsen Khvedelidze, Astghik Torosyan

Laboratory of Information Technologies  
Joint Institute for Nuclear Research  
Dubna, Russia



# Content

- 1 Objective
- 2 Introduction
  - Wigner function
  - Stratonovich-Weyl kernel
  - Master equations
- 3 Moduli space parametrization
- 4 Examples: Qubit, Qutrit, Quatrit
- 5 Conclusions

## Context

Recently an ambiguity in specification of the Wigner quasiprobability distribution for a finite-dimensional quantum system has been studied.

It was shown that for an  $N$ -level quantum system one can construct  $N - 2$  parametric family of unitary non-equivalent Wigner quasiprobability distributions.

## The main objective

In the report the moduli space of the Wigner quasiprobability distributions for  $N$ -dimensional quantum systems will be discussed and exemplified for low dimensional cases: for a single qubit, qutrit and quatrit.

# Introduction

## Defining a quantum state

A state of an  $N$ -level quantum-mechanical system is described by a **density operator**  $\rho$  acting on the  $\mathbb{C}^N$  Hilbert space <sup>a</sup>, satisfying the conditions:

- Hermicity :  $\rho^\dagger = \rho$  ;
- Completeness :  $tr(\rho) = 1$  ;
- Semi-positivity :  $\langle \psi | \rho | \psi \rangle \geq 0$  ;

---

<sup>a</sup>The states with  $\rho^2 = \rho$  are called pure states.

## The Wigner function

is constructed from the density matrix  $\rho$  describing a quantum state and the **Stratonovich-Weyl self-dual kernel**  $\Delta(\Omega_N)$  defined over the symplectic manifold  $\Omega_N$ :

$$W_\rho(\Omega) = tr(\rho \Delta(\Omega)) .$$

**Stratonovich-Weyl:** operators( $\mathcal{H}$ )  $\xleftrightarrow{\Delta(\Omega)}$  functions( $\Omega$ ), at that

- ① *Reconstruction* of the state:

$$\varrho = \int_{\Omega} d\Omega \Delta(\Omega) W_{\varrho}(\Omega);$$

- ② *Hermicity* of the kernel:

$$\Delta(\Omega) = \Delta(\Omega)^{\dagger};$$

- ③ *Finite norm*:

$$\text{tr}(\varrho) = \int_{\Omega} d\Omega W_{\varrho}(\Omega), \quad \int_{\Omega} d\Omega \Delta(\Omega) = 1;$$

- ④ *Covariance*: the unitary symmetry  $\varrho' = U(\alpha) \varrho U^{\dagger}(\alpha)$  induces the kernel transformation <sup>a</sup>

$$\Delta(\Omega') = U(\alpha)^{\dagger} \Delta(\Omega) U(\alpha).$$

<sup>a</sup>C. Brif, A. Mann, Phys.Rev.A, 59, 2, 1999

## Master equations:

$$\text{tr}(\Delta(\Omega)) = 1, \quad \text{tr}(\Delta(\Omega)^2) = N.$$

1

---

<sup>1</sup>V. Abgaryan, A. Khvedelidze, *On families of Wigner functions for N-level quantum systems*, arXiv:1708.05981v3, 2018

## The Stratonovich-Weyl kernel

$$\Delta(\Omega|\nu) = \frac{1}{N} U(\Omega) \left[ I + \kappa \sum_{\lambda \in H} \mu_s(\nu) \lambda_s \right] U(\Omega)^\dagger, \quad \kappa = \sqrt{N(N^2 - 1)/2},$$

where

- $H$  is the **Cartan subalgebra** in  $SU(N)$ ,
- parameter  $\nu = (\nu_1, \dots, \nu_{N-2})$  labels members of the WF family,
- coefficients  $\left[ \sum_{s=2}^N \mu_{s^2-1}^2(\nu) = 1 \right]$ .

## A density matrix of an $N$ -dimensional quantum system

$$\varrho_\xi = \frac{1}{N} \left[ I + \sqrt{\frac{N(N-1)}{2}} (\xi, \lambda) \right],$$

where

- $\xi$  is an  $(N^2 - 1)$ -dimensional Bloch vector,
- $\lambda = \{\lambda_1, \dots, \lambda_{N^2-1}\}$  is  $\mathfrak{su}(N)$  algebra basis.



$$W = \text{tr}(\varrho \Delta)$$

## A family of the Wigner functions

$$W_{\xi}^{(\nu)}(\Omega_N) = \frac{1}{N} \left[ 1 + \frac{N^2 - 1}{\sqrt{N + 1}} (\mathbf{n}, \xi) \right],$$

where

- $\mathbf{n} = \mu_3 \mathbf{n}^{(3)} + \dots + \mu_{N^2-1} \mathbf{n}^{(N^2-1)},$
- $\mathbf{n}^{(s^2-1)} = \frac{1}{2} \text{tr} (U \lambda_{s^2-1} U^\dagger \lambda_{\mu}), \quad s = \overline{2, N}.$

## The spectrum $\{\pi_1, \dots, \pi_N\}$ of the Stratonovich-Weyl kernel:

$$\pi_i = \frac{1}{N} \left( 1 + \sqrt{2} \kappa \sum_{s=i+1}^N \frac{\mu_{s^2-1}}{\sqrt{s(s-1)}} - \kappa \sqrt{\frac{2(i-1)}{i}} \mu_{i^2-1} \right).$$

# Constraints on the spherical angles

## The spherical $(N - 2)$ angles:

$$\mu_3 = \sin \psi_1 \cdots \sin \psi_{N-2},$$

$$\vdots$$

$$\mu_{i^2-1} = \sin \psi_1 \cdots \sin \psi_{N-i} \cos \psi_{N-i+1},$$

$$\vdots$$

$$\mu_{N^2-1} = \cos \psi_1, \quad i = \overline{2, N}.$$

## For decreasing order $\pi_1 \geq \cdots \geq \pi_N$

$$\mu_3 \geq 0, \quad \mu_{(i+1)^2-1} \geq \sqrt{\frac{i-1}{i+1}} \mu_{i^2-1}, \quad i = \overline{2, N-1}.$$

# Examples: qubit, qutrit and quatrit

# The Wigner function of a single qubit

A generic **qubit** quantum state is parameterized in a standard way

$$\rho_{\text{qubit}} = \frac{1}{2} (I + \mathbf{r} \cdot \boldsymbol{\sigma})$$

by the Bloch vector  $\mathbf{r} = (r \sin \psi \cos \phi, r \sin \psi \sin \phi, r \cos \psi)$ .

The master equations determine the spectrum:

$$\text{spec} \left( P^{(2)} \right) = \left\{ \frac{1 + \sqrt{3}}{2}, \frac{1 - \sqrt{3}}{2} \right\}.$$

The **Wigner function** for a single qubit is

$$W_{\mathbf{r}}(\alpha, \beta) = \frac{1}{2} + \frac{\sqrt{3}}{2} (\mathbf{r}, \mathbf{n}),$$

where  $\mathbf{n} = (-\cos \alpha \sin \beta, \sin \alpha \sin \beta, \cos \beta)$  is the unit 3-vector.

## Qutrit kernel and its fundamental region

A generic **qutrit** state is given by the density matrix

$$\rho_{\text{qutrit}} = \frac{1}{3} \left( I + \sqrt{3} \sum_{\nu=1}^8 \xi_{\nu} \lambda_{\nu} \right).$$

The **Stratonovich-Weyl** kernel

$$\Delta(\Omega_3) = U(\Omega_3) \frac{1}{3} \left[ I + 2\sqrt{3} (\mu_3 \lambda_3 + \mu_8 \lambda_8) \right] U(\Omega_3)^{\dagger},$$

where the coefficients

$$\mu_3(\nu) = \frac{\sqrt{3}}{4} \sqrt{(1+\nu)(5-3\nu)}, \quad \mu_8(\nu) = \frac{1}{4}(1-3\nu)$$

are functions of the parameter  $\nu = \frac{1}{3} - \frac{4}{3} \cos(\zeta)$  with  $\zeta \in [0, \pi/3]$  being the moduli parameter of the unitary nonequivalent WF of a qutrit.

The **Wigner function** of a single qutrit

$$W_{\xi}^{(\nu)}(\Omega_3) = \frac{1}{3} + \frac{4}{3} [\mu_3(\mathbf{n}^{(3)}, \xi) + \mu_8(\mathbf{n}^{(8)}, \xi)],$$

with two orthogonal unit 8-vectors

$$n_{\nu}^{(3)} = \frac{1}{2} \text{tr} [U \lambda_3 U^{\dagger} \lambda_{\nu}], \quad n_{\nu}^{(8)} = \frac{1}{2} \text{tr} [U \lambda_8 U^{\dagger} \lambda_{\nu}].$$

The **master equations**

$$\text{tr}(\Delta(\Omega)) = 1, \quad \text{tr}(\Delta(\Omega)^2) = 3$$

determine one-parametric family of kernels  $P^{(3)}(\nu)$ .

# One-parametric $P^{(3)}(\nu)$ -family

- The spectrum of **generic** kernels:

$$\text{spec} \left( P^{(3)}(\nu) \right) = \left\{ \frac{1 - \nu + \delta}{2}, \frac{1 - \nu - \delta}{2}, \nu \right\},$$

where  $\delta = \sqrt{(1 + \nu)(5 - 3\nu)}$  and  $\nu \in (-1, -\frac{1}{3})$ .

- Two **degenerate** kernels:

$$\text{spec} \left( P^{(3)}(-1) \right) = \{1, 1, -1\}, \quad \text{spec} \left( P^{(3)}(-1/3) \right) = \left\{ \frac{5}{3}, -\frac{1}{3}, -\frac{1}{3} \right\}.$$


- The spectrum of **singular** kernel:

$$\text{spec} \left( P_{\det=0}^{(3)} \right) = \left\{ \frac{1 + \sqrt{5}}{2}, 0, \frac{1 - \sqrt{5}}{2} \right\}, \quad \text{tr} \left( [P_{\det=0}^{(3)}]^m \right) = \mathcal{L}_m,$$

where the  $m$ -th **Lucas number**  $\mathcal{L}_m = \phi^m + (-\phi)^{-m}$  and  $\phi = \frac{1 + \sqrt{5}}{2}$ .


The ordering of the **density matrix** eigenvalues  $1 \geq r_1 \geq r_2 \geq r_3 \geq 0$  and condition  $\sum r_i = 1$  lead to

$$\xi_3 \geq 0, \quad \xi_8 \geq \frac{\xi_3}{\sqrt{3}}.$$



The ordering of the **SW kernel** eigenvalues  $\pi_1 \geq \pi_2 \geq \pi_3$  and condition  $\sum \mu_i^2 = 1$  lead to

$$\mu_3 = \sin \zeta, \quad \mu_8 = \cos \zeta, \quad 0 \leq \zeta \leq \frac{\pi}{3}.$$





## Quatrit kernel and its fundamental region

A generic **quatrit** ( $N = 4$ ) state is given by the density matrix

$$\rho_{\text{quatrit}} = \frac{1}{4} \left( I + \sqrt{6} \sum_{\nu=1}^{15} \xi_{\nu} \lambda_{\nu} \right).$$

The **Stratonovich-Weyl kernel**

$$\Delta(\Omega_N | \nu) = U(\Omega_N) \frac{1}{4} \left[ I + \sqrt{30} (\mu_3 \lambda_3 + \mu_8 \lambda_8 + \mu_{15} \lambda_{15}) \right] U(\Omega_N)^{\dagger}.$$

The **Wigner function** of a quatrit


$$W_{\xi}^{(\nu)}(\Omega_4) = \frac{1}{4} + \frac{3\sqrt{5}}{4} \left[ \mu_3(\mathbf{n}^{(3)}, \xi) + \mu_8(\mathbf{n}^{(8)}, \xi) + \mu_{15}(\mathbf{n}^{(15)}, \xi) \right],$$

with

$$n_{\nu}^{(3,8,15)} = \frac{1}{2} \text{tr} \left[ U \lambda_{3,8,15} U^{\dagger} \lambda_{\nu} \right].$$

# Quatrit density matrix

In a quatrit case, there are 24 ways of the spec  $(\rho_{quatrit}) = \{r_1, r_2, r_3, r_4\}$  ordering.



The fixed order of the eigenvalues

$$1 \geq r_1 \geq r_2 \geq r_3 \geq r_4 \geq 0,$$

$$0 \leq r_i \leq 1, \quad \sum r_i = 1,$$

leads to

$$0 \leq \xi_3 \leq \sqrt{2/3},$$

$$\frac{\xi_3}{\sqrt{3}} \leq \xi_8 \leq \sqrt{2/3},$$

$$\frac{\xi_8}{\sqrt{2}} \leq \xi_{15} \leq 1/3.$$

## The master equations

$$\text{tr}(\Delta(\Omega)) = 1, \quad \text{tr}(\Delta(\Omega)^2) = 4$$


determine two-parametric family of kernels  $P^{(4)}$  with  $\pi_1 \geq \pi_2 \geq \pi_3 \geq \pi_4$ :

- **Generic** kernel:

$$\text{spec} \left( P^{(4)}(\nu_1, \nu_2) \right) = \left\{ \frac{1 - \nu_1 - \nu_2 + \delta}{2}, \frac{1 - \nu_1 - \nu_2 - \delta}{2}, \nu_1, \nu_2 \right\},$$

where

$$\delta = \sqrt{7 + 2\nu_1 - 3\nu_1^2 + 2\nu_2 - 2\nu_1\nu_2 - 3\nu_2^2}.$$



Note that

$$\mathcal{R}_m = \mathcal{R}_{m-1} + \frac{3}{2}\mathcal{R}_{m-2}, \quad \mathcal{R}_1 = 1, \mathcal{R}_2 = 4;$$

$$\mathcal{L}_m = \mathcal{L}_{m-1} + \mathcal{L}_{m-2}, \quad \mathcal{L}_1 = 2, \mathcal{L}_2 = 1.$$

- **Degenerate kernels:**

- Double degenerate

$$1|234 : \pi_1 = \pi_2 \neq \pi_3 \neq \pi_4,$$

$$12|34 : \pi_1 \neq \pi_2 = \pi_3 \neq \pi_4,$$

$$123|4 : \pi_1 \neq \pi_2 \neq \pi_3 = \pi_4,$$

$$1|23|4 : \pi_1 = \pi_2 \neq \pi_3 = \pi_4.$$

- Triple degenerate

$$12|3|4 : \pi_1 \neq \pi_2 = \pi_3 = \pi_4,$$

$$1|2|34 : \pi_1 = \pi_2 = \pi_3 \neq \pi_4.$$

- **Singular kernels**

$$1034 : \pi_1 \neq \pi_2 = 0 \neq \pi_3 \neq \pi_4,$$

$$1204 : \pi_1 \neq \pi_2 = 0 \neq \pi_3 \neq \pi_4,$$

$$1004 : \pi_1 \neq \pi_2 \neq \pi_3 = 0 \neq \pi_4,$$

$$\text{with } \text{tr} \left( [P_{1004}^{(4)}]^m \right) = \mathcal{R}_m.$$

Parameterizing  $\mu$  by two spherical coordinates

$$\mu_3 = \sin \psi_1 \sin \psi_2, \quad \mu_8 = \sin \psi_1 \cos \psi_2, \quad \mu_{15} = \cos \psi_1$$

and using the constraints coming from the requirement of a decreasing order of the SW kernel's eigenvalues

$$\mu_3 \geq 0, \quad \mu_8 \geq \frac{\mu_3}{\sqrt{3}}, \quad \mu_{15} \geq \frac{\mu_8}{\sqrt{2}},$$

we have:

$$\left[ \begin{array}{l} \left\{ \begin{array}{l} \psi_2 \in (0, \frac{\pi}{3}] , \\ 0 < \psi_1 \leq \operatorname{arccot} (\cos \psi_2 / \sqrt{2}) ; \end{array} \right. \\ \\ \left\{ \begin{array}{l} \psi_2 = 0 , \\ 0 < \psi_1 \leq \operatorname{arccot} (1 / \sqrt{2}) ; \end{array} \right. \\ \\ \psi_1 = 0 . \end{array} \right. \quad (\text{See Figure 1})$$

**Girard's theorem:** the spherical excess of a triangle determines the solid angle

$$\pi/2 + \pi/3 + \pi/3 - \pi = 4\pi/24.$$

Any fixed order of eigenvalues corresponds to one of 24 possible ways to tessellate a sphere.




Figure 1: Möbius (2, 3, 3) triangle with  $(\pi/2, \pi/3, \pi/3)$  angles.

# Conclusions

An ambiguity in the master equation's solution for Stratonovich-Weyl kernel is analyzed and the corresponding moduli spaces of the Wigner QPDF is determined for  $N = 2, 3, 4$  quantum systems:

- for the qubit the moduli space is the single point,
- for the qutrit the moduli space is the  $\frac{\pi}{3}$  arc of the unit circle,
- for the quatrit the moduli space is  $(2, 3, 3)$  Möbius triangle.

The basic goal of our further studies is

- to understand a **physical meaning** of the Wigner function moduli space;
- to clarify the role these unitary invariant moduli parameters play in **dynamics** of classical and quantum systems.

Thank you <sup>2</sup> you

---

<sup>2</sup>V. Abgaryan, A. Khvedelidze, A. Torosyan, *On moduli space of the Wigner quasiprobability distributions for N-dimensional quantum systems*, Zap. Nauch. Sem. POMI, 468, 177-201, 2018 [↗](#)