Contribution ID: 10

Type: not specified

Self-consistent study of isobaric analog resonances in nuclei with pairing

Wednesday, 17 July 2019 15:20 (20 minutes)

Fully self-consistent calculation of the IAR resonances is performed for the long chain of Ni isotopes with Tz \neq 0. The recently established DF3-f parametrization [1] of the Fayans energy density functional [2] is used. Its performance in "reference" neutron-rich Sn and Pb isotopes with developed pairing correlations is also studied. The main aim is to figure out the influence of T=1 pn-pairing on IAR characteristics and to check how the self-consistency is preserved in the calculations for long isotopic chains. Comparison with the available calculations using relativistic Hartree-Bogolubov plus D3C* functional [3] and the HF+BCS with SAMi functional [4] is performed,

[1] I.N. Borzov, S.V. Tolokonnikov, Physics of Atomic Nuclei v.82 № 6 (2019)

[2] S. A. Fayans, JETP Letters 68, 161 (1998)

[3] N.Paar, T. Niksic, D. Vretenar and P. Ring, Physical Review C 69, 054303 (2004)

[4] Li-Gang Cao, X. Roca-Mazo, G. Colo, H. Sagawa, Phys.Rev.92, 034308 (2015)

Primary author: Dr BORZOV, Ivan (NRC KI, BLTP JINR)

Co-author: Dr TOLOKONNIKOV, Sergei (NRC KI)

Presenter: Dr BORZOV, Ivan (NRC KI, BLTP JINR)

Session Classification: Modern problems in nuclear and elementary particle physics