

γ emission from neutron-unbound states in ^{133}Sn

Wednesday, 17 April 2019 14:15 (15 minutes)

The study of ^{133}Sn provides excellent conditions to investigate single-particle transitions relevant in the neutron-rich ^{132}Sn region due to the simplicity of its nuclear structure. After many experimental activities employing one-neutron transfer reactions [1–4], traditional β -decay studies are an attractive technique to refine our knowledge on ^{133}Sn . Since the positions of neutron single-particle states in ^{133}Sn were established and confirmed in many measurements [1–5], our focus moves to single-hole states expected at higher excitation energies. Because of the low neutron-separation energy of ^{133}Sn , $S_n=2.4$ MeV [6], all of them are supposed to be neutron-unbound. β -decay studies are therefore a natural choice to investigate their nature since there is a large energy window for their population in the β decay of ^{133}In ($Q_\beta=13.4(2)$ MeV [6]).

Our experiment was performed at the ISOLDE Decay Station, where excited states in ^{133}Sn were investigated via the β decay of ^{133}In . Isomer-selective ionization using the ISOLDE RILIS enabled the β decays of ^{133}gIn ($I\pi=9/2^+$) and ^{133}mIn ($I\pi=1/2^-$) to be studied independently for the first time. Thanks to the large spin difference of those two β -decaying states, it is possible to investigate separately the lower- and higher-spin states in the daughter ^{133}Sn and thus to probe independently different single-particle and single-hole levels. We identified new γ transitions following the $^{133}\text{In}\rightarrow^{133}\text{Sn}$ decay. Single-hole states in ^{133}Sn were found at energies exceeding S_n up to 3.7 MeV [7]. Due to centrifugal barrier hindering the neutron from leaving the nucleus, the contribution of electromagnetic decay of those unbound states was found to be significant.

- [1] K. L. Jones et al., Nature (London) 465, 454 (2010).
- [2] K. L. Jones et al., Phys. Rev. C 84, 034601 (2011).
- [3] J. M. Allmond et al., Phys. Rev. Lett. 112, 172701 (2014).
- [4] V. Vaquero et al., Phys. Rev. Lett. 118, 202502 (2017).
- [5] P. Hoff et al., Phys. Rev. Lett. 77, 1020 (1996).
- [6] M. Wang et al., Chin. Phys. C 41, 030003 (2017).
- [7] M. Piersa et al., Phys. Rev. C 99, 024304 (2019).

Primary author: Ms PIERSA, Monika (Faculty of Physics University of Warsaw)

Presenter: Ms PIERSA, Monika (Faculty of Physics University of Warsaw)

Session Classification: Nuclear Physics

Track Classification: Nuclear Physics