

Chiral heat-vortical wave in cold Fermi liquid and the deformation of zero sound

David Frenklakh

August 24, 2017

The Helmholtz International Summer School "Hadron Structure, Hadronic Matter, and Lattice QCD"

The plan

- Anomalous transport effects
- Chiral waves
- Chiral kinetic theory formalism for Fermi liquid
- Results

- blah-blah

- Anomalous transport effects should reveal themselves in heavy ion collisions

- Anomalous transport effects should reveal themselves in heavy ion collisions
- Namely, chiral waves should cause generation of quadrupole moment in quark-gluon plasma

- Chiral magnetic effect

$$\mathbf{j}_V = \frac{\mu_A \mathbf{B}}{2\pi^2}$$

Anomalous transport effects

- Chiral magnetic effect

$$\mathbf{j}_V = \frac{\mu_A \mathbf{B}}{2\pi^2}$$

- Chiral separation effect

$$\mathbf{j}_A = \frac{\mu_V \mathbf{B}}{2\pi^2}$$

- Chiral magnetic effect

$$\mathbf{j}_V = \frac{\mu_A \mathbf{B}}{2\pi^2}$$

- Chiral separation effect

$$\mathbf{j}_A = \frac{\mu_V \mathbf{B}}{2\pi^2}$$

- Chiral vortical effect

$$\mathbf{j}_A = \left[\frac{(\mu_V^2 + \mu_A^2)}{2\pi^2} + \frac{T^2}{6} \right] \boldsymbol{\Omega}$$

$$\mathbf{j}_V = \frac{\mu_V \mu_A}{\pi^2} \boldsymbol{\Omega}$$

Chiral magnetic wave

- Chiral effects + axial and vector current conservation laws

Chiral magnetic wave

- Chiral effects + axial and vector current conservation laws
- Chiral magnetic wave:

$$\mathbf{j}_{V/A} = \frac{\mu_{A/V} \mathbf{B}}{2\pi^2},$$
$$\partial_t \rho_{V/A} + \boldsymbol{\partial} \cdot \mathbf{j}_{V/A} = 0$$

Chiral magnetic wave

- Chiral effects + axial and vector current conservation laws
- Chiral magnetic wave:

$$\mathbf{j}_{V/A} = \frac{\mu_{A/V} \mathbf{B}}{2\pi^2},$$
$$\partial_t \rho_{V/A} + \boldsymbol{\partial} \cdot \mathbf{j}_{V/A} = 0$$

- Consider small fluctuations above neutral background ($\mu_V = \mu_A = 0$)

$$\delta \rho_{V/A} = \chi \delta \mu_{V/A}$$

Chiral magnetic wave

- Chiral effects + axial and vector current conservation laws
- Chiral magnetic wave:

$$\mathbf{j}_{V/A} = \frac{\mu_{A/V} \mathbf{B}}{2\pi^2},$$
$$\partial_t \rho_{V/A} + \boldsymbol{\partial} \cdot \mathbf{j}_{V/A} = 0$$

- Consider small fluctuations above neutral background ($\mu_V = \mu_A = 0$)

$$\delta \rho_{V/A} = \chi \delta \mu_{V/A}$$
$$\omega \chi \delta \mu_{V/A} + \frac{(\mathbf{k} \cdot \mathbf{B}) \delta \mu_{A/V}}{2\pi^2} = 0$$

Dispersion relation

$$v_{CMW} = \frac{B}{2\pi^2 \chi}$$

Chiral vortical wave

- Chiral vortical effect is quadratic in chemical potentials \Rightarrow
- We need finite vector charge density in the background

$$\mathbf{j}_A = \left[\frac{(\mu_V^2 + \mu_A^2)}{2\pi^2} + \frac{T^2}{6} \right] \boldsymbol{\Omega}$$
$$\mathbf{j}_V = \frac{\mu_V \mu_A}{\pi^2} \boldsymbol{\Omega}$$

Chiral vortical wave

- Chiral vortical effect is quadratic in chemical potentials \Rightarrow
- We need finite vector charge density in the background

$$\mathbf{j}_A = \left[\frac{(\mu_V^2 + \mu_A^2)}{2\pi^2} + \frac{T^2}{6} \right] \boldsymbol{\Omega}$$

$$\mathbf{j}_V = \frac{\mu_V \mu_A}{\pi^2} \boldsymbol{\Omega}$$

$$\partial_t \rho_{V/A} + \boldsymbol{\partial} \cdot \mathbf{j}_{V/A} = 0$$

- Assume that temperature is kept constant

$$\delta \rho_{V/A} = \chi \delta \mu_{V/A}$$

Chiral vortical wave

- Chiral vortical effect is quadratic in chemical potentials \Rightarrow
- We need finite vector charge density in the background

$$\mathbf{j}_A = \left[\frac{(\mu_V^2 + \mu_A^2)}{2\pi^2} + \frac{T^2}{6} \right] \boldsymbol{\Omega}$$

$$\mathbf{j}_V = \frac{\mu_V \mu_A}{\pi^2} \boldsymbol{\Omega}$$

$$\partial_t \rho_{V/A} + \boldsymbol{\partial} \cdot \mathbf{j}_{V/A} = 0$$

- Assume that temperature is kept constant

$$\delta \rho_{V/A} = \chi \delta \mu_{V/A}$$

Dispersion relation

$$v_{CVW} = \frac{\mu_V \Omega}{2\pi^2 \chi}$$

- Anomalous energy transport due to rotation:

$$\mathbf{j}_E = \frac{\mu_A}{3} \left[\frac{3\mu_V^2 + \mu_A^2}{\pi^2} + T^2 \right] \boldsymbol{\Omega}$$

- Anomalous energy transport due to rotation:

$$\mathbf{j}_E = \frac{\mu_A}{3} \left[\frac{3\mu_V^2 + \mu_A^2}{\pi^2} + T^2 \right] \boldsymbol{\Omega}$$

- Energy conservation law:

$$\partial_t \epsilon + \boldsymbol{\partial} \cdot \mathbf{j}_E = 0$$

- Anomalous energy transport due to rotation:

$$\mathbf{j}_E = \frac{\mu_A}{3} \left[\frac{3\mu_V^2 + \mu_A^2}{\pi^2} + T^2 \right] \boldsymbol{\Omega}$$

- Energy conservation law:

$$\partial_t \epsilon + \boldsymbol{\partial} \cdot \mathbf{j}_E = 0$$

- For small fluctuations of vector charge and energy density above equilibrium

$$\delta \rho_V = \chi \delta \mu_V + \alpha \delta T$$

$$\delta \epsilon = C \delta T + \gamma \delta \mu_V$$

Dispersion relation

$$v_{CHW} = \sqrt{\frac{T^3}{C\chi - \alpha\gamma} \frac{\Omega}{3}}$$

Mixed chiral heat-vortical wave

- Non-zero background charge density
- Temperature, vector and axial charges fluctuating

Mixed chiral heat-vortical wave

- Non-zero background charge density
- Temperature, vector and axial charges fluctuating

Dispersion relation

$$v_{CHVW} = \sqrt{\frac{\Omega^2 |9\mu_V^2(C - \alpha\mu_V) - 3\mu_V T \pi^2(\alpha T + \gamma - \chi\mu_V) + \pi^2 \chi T^3|}{9\pi^4 \chi |C\chi - \alpha\gamma|}}$$

- System of strongly interacting chiral particles

Berry Fermi liquid

- System of strongly interacting chiral particles
- Not far from equilibrium is described in terms of quasiparticles

- System of strongly interacting chiral particles
- Not far from equilibrium is described in terms of quasiparticles
- The anomalous effects are captured by Berry connection in momentum space

$$\frac{\partial n_{R/L}}{\partial t} + \dot{\mathbf{x}}_{R/L} \cdot \frac{\partial n_{R/L}}{\partial \mathbf{x}} + \dot{\mathbf{p}}_{R/L} \cdot \frac{\partial n_{R/L}}{\partial \mathbf{p}} = C_{R/L}[n_R, n_L]$$

- $n_{R/L}(t, \mathbf{x}, \mathbf{p})$ are right and left quasiparticles distribution functions
- $C_{R/L}$ are respective collision integrals

$$\begin{aligned}\sqrt{G_{R/L}}\dot{\mathbf{x}}_{R/L} &= \mathbf{v}_{R/L} + 2\epsilon_{R/L}\boldsymbol{\Omega}(\mathbf{v}_{R/L} \cdot \mathbf{b}_{R/L}) + \boldsymbol{\mathcal{E}}_{R/L} \times \mathbf{b}_{R/L} \\ \sqrt{G_{R/L}}\dot{\mathbf{p}}_{R/L} &= \boldsymbol{\mathcal{E}}_{R/L} + 2\epsilon_{R/L}\mathbf{v}_{R/L} \times \boldsymbol{\Omega} + (\boldsymbol{\mathcal{E}}_{R/L} \cdot \boldsymbol{\Omega})2\epsilon_{R/L}\mathbf{b}_{R/L}\end{aligned}$$

- Here $\epsilon_{R/L}$ are quasiparticles energy functionals,
- $\mathbf{v}_{R/L} = \frac{\partial \epsilon_{R/L}}{\partial \mathbf{p}}$, $\boldsymbol{\mathcal{E}}_{R/L} = -\frac{\partial \epsilon_{R/L}}{\partial \mathbf{x}}$
- $\mathbf{b}_{R/L} = \pm \frac{\hat{\mathbf{p}}}{2p^3}$ are Berry connections curvature in momentum space ($p = |\mathbf{p}|$)
- $\sqrt{G_{R/L}} = 1 + 2\epsilon_{R/L}(\mathbf{b}_{R/L} \cdot \boldsymbol{\Omega})$ modify phase space volume

- We are considering the small fluctuations above the equilibrium distribution functions:

$$n_{R/L} = n_{R/L}^0 + \delta n_{R/L} e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}$$

where

$$n_{R/L}^0 = \frac{1}{e^{\beta(\epsilon^0 - \mu)} + 1}$$

- Here $\beta = T^{-1}$ and we assume $\mu \gg T$

- We are considering the small fluctuations above the equilibrium distribution functions:

$$n_{R/L} = n_{R/L}^0 + \delta n_{R/L} e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)}$$

where

$$n_{R/L}^0 = \frac{1}{e^{\beta(\epsilon^0 - \mu)} + 1}$$

- Here $\beta = T^{-1}$ and we assume $\mu \gg T$
- Two regimes: $\omega \ll \tau^{-1}$ (hydrodynamic regime) $\omega \gg \tau^{-1}$ (collisionless regime). Here τ is characteristic relaxation time of collision integral

Dispersion relation

$$\omega = \pm \frac{\mu(\mathbf{k} \cdot \boldsymbol{\Omega})}{2\pi^2\chi} \sqrt{F_1 F_2} \left[1 + \frac{T^2 \pi^2}{6\mu^2 F_1} (1 - 4A + 4A^2) \right]$$

- $A = \frac{\mu}{v_{FPF}}$
- F_1 and F_2 are taken from linearised energy functionals

Non-modified implicit zero sound dispersion relation

$$\operatorname{arccotanh} s_0 = \frac{1}{s_0} \left(\frac{1}{2(F_S \pm F_A)} + 1 \right)$$

- Here $s_0 = \frac{\omega}{v_f k}$, F_S and F_A are taken from linearised energy functionals

Modification due to the chiral heat-vortical wave

$$\delta s = s - s_0 \approx \mp \frac{\omega^2 \mu^2}{p_F^4} \left(1 - \frac{2\pi^2 T^2}{3v_F^2 p_F^2} \right).$$

$$\frac{F_S L_2(s_0) + [L_1(s_0)^2 - 2L_0(s_0)L_2(s_0)](F_S^2 - F_A^2)}{F_A [\text{arccotanh } s_0 - \frac{s_0}{2((s_0)^2 - 1)}]}$$

$$L_0(s_0) = s_0 \text{arccotanh } s_0 - 1$$

$$L_1(s_0) = 3s_0(s_0 \text{arccotanh } s_0 - 1)$$

$$L_2(s_0) = 2s_0[-3s_0 + (3s_0^2 - 1) \text{arccotanh } s_0]$$

- The result for velocity of chiral heat-vortical wave coincides with the one known from hydrodynamics

- The result for velocity of chiral heat-vortical wave coincides with the one known from hydrodynamics
- There turn out to be two branches of modified zero sound with the correction to velocity being quadratic in angular velocity

THANK YOU FOR YOUR ATTENTION